

The BKL Conjectures for Spatially Homogeneous Spacetimes

Michael Reiterer, Eugene Trubowitz

Department of Mathematics, ETH Zurich, Switzerland

Abstract: We rigorously construct and control a generic class of spatially homogeneous (Bianchi VIII and Bianchi IX) vacuum spacetimes that exhibit the oscillatory BKL phenomenology. We investigate the causal structure of these spacetimes and show that there is a “particle horizon”.

1. Introduction

The goal of this paper is to rigorously construct and explicitly control a generic class of solutions $\Phi = \alpha \oplus \beta : [0, \infty) \rightarrow \mathbb{R}^3 \oplus \mathbb{R}^3$, with independent variable $\tau \in [0, \infty)$ and with¹ $(\alpha_1 + \alpha_2 + \alpha_3)|_{\tau=0} < 0$, to the autonomous system of six ordinary differential equations

$$0 = -\frac{d}{d\tau}\alpha_i - (\beta_i)^2 + (\beta_j)^2 + (\beta_k)^2 - 2\beta_j\beta_k \quad (1.1a)$$

$$0 = -\frac{d}{d\tau}\beta_i + \beta_i\alpha_i \quad (1.1b)$$

for all $(i, j, k) \in \mathcal{C} \stackrel{\text{def}}{=} \{(1, 2, 3), (2, 3, 1), (3, 1, 2)\}$, subject to the quadratic constraint²

$$0 = \alpha_2\alpha_3 + \alpha_3\alpha_1 + \alpha_1\alpha_2 - (\beta_1)^2 - (\beta_2)^2 - (\beta_3)^2 + 2\beta_2\beta_3 + 2\beta_3\beta_1 + 2\beta_1\beta_2 \quad (1.1c)$$

Here, $\alpha = (\alpha_1, \alpha_2, \alpha_3)$, $\beta = (\beta_1, \beta_2, \beta_3)$. The system (1.1) are the vacuum Einstein equations for spatially homogeneous (Bianchi) spacetimes, see Proposition 2.1.

The pioneering calculations and heuristic picture of Belinskii, Khalatnikov, Lifshitz³ [BKL1] and Misner [Mis] suggest that a generic class of solutions to (1.1) are oscillatory as $\tau \rightarrow +\infty$ and that the dynamics of one degree of freedom is closely related to the discrete dynamics of the Gauss map $G(x) = \frac{1}{x} - \lfloor \frac{1}{x} \rfloor$, a non-invertible map

¹ If $\tau \mapsto \Phi(\tau)$ is a solution to (1.1), so is $\tau \mapsto -\Phi(-\tau)$. The condition $(\alpha_1 + \alpha_2 + \alpha_3)|_{\tau=0} < 0$ breaks this symmetry. Solutions to (1.1) with $(\alpha_1 + \alpha_2 + \alpha_3)|_{\tau=0} < 0$ do not break down in finite positive time, that is, they extend to $[0, \infty)$. A proof of this fact is given later in this introduction.

² As a quadratic form on $\mathbb{R}^3 \oplus \mathbb{R}^3$, the right hand side of (1.1c) has signature $(+, +, -, -, -, -)$.

³ The work of Belinskii, Khalatnikov, Lifshitz concerns general (inhomogeneous) spacetimes, but relies on intuition about the homogeneous case.

from $(0, 1) \setminus \mathbb{Q}$ to itself. Every element of $(0, 1) \setminus \mathbb{Q}$ admits a unique infinite continued fraction expansion

$$\langle k_1, k_2, k_3, \dots \rangle = \frac{1}{k_1 + \frac{1}{k_2 + \frac{1}{k_3 + \dots}}} \quad (1.2)$$

where $(k_n)_{n \geq 1}$ are strictly positive integers. The Gauss map is the left-shift,

$$G(\langle k_1, k_2, k_3, \dots \rangle) = \langle k_2, k_3, k_4, \dots \rangle \quad (1.3)$$

Rigorous results about spatially homogeneous spacetimes have been obtained by Rendall [Ren] and Ringström [Ri1], [Ri2]. See also Heinzle and Uggla [HU2]. We refer to the very readable paper [HU1] for a detailed discussion.

The first rigorous proofs that there exist spatially homogeneous vacuum spacetimes whose asymptotic behavior is related, in a precise sense, to iterates of the Gauss map, have been obtained recently by Béguin [Be] and by Liebscher, Härterich, Webster and Georgi [LHWG]. These theorems apply to a dense subset of $(0, 1) \setminus \mathbb{Q}$. A basic restriction of both these works is that the sequence $(k_n)_{n \geq 1}$ has to be bounded, a condition fulfilled only by a Lebesgue measure zero subset of $(0, 1) \setminus \mathbb{Q}$. The results of the present paper apply to any sequence $(k_n)_{n \geq 1}$ that grows at most polynomially. The corresponding subset of $(0, 1) \setminus \mathbb{Q}$ has full Lebesgue measure one.

We point out some properties of the system (1.1a), (1.1b), not assuming (1.1c):

- (i) The right hand side of (1.1c) is a conserved quantity.
- (ii) If $\tau \mapsto \Phi(\tau)$ is a solution, so is $\tau \mapsto p\Phi(p\tau + q)$, for all $p, q \in \mathbb{R}$.
- (iii) The signatures $(\text{sgn } \beta_1, \text{sgn } \beta_2, \text{sgn } \beta_3)$ are constant.
- (iv) $\frac{d}{d\tau} |\beta_1 \beta_2 \beta_3|^2 = 2(\alpha_1 + \alpha_2 + \alpha_3) |\beta_1 \beta_2 \beta_3|^2$.
- (v) We have⁴ $\frac{d}{d\tau} (\alpha_1 + \alpha_2 + \alpha_3) \geq -3 |\beta_1 \beta_2 \beta_3|^{2/3}$.

If in addition we assume (1.1c), then:

- (vi) $\frac{d}{d\tau} (\alpha_1 + \alpha_2 + \alpha_3) = \alpha_2 \alpha_3 + \alpha_3 \alpha_1 + \alpha_1 \alpha_2 \leq \frac{1}{3} (\alpha_1 + \alpha_2 + \alpha_3)^2$.

Let $\Phi = \alpha \oplus \beta$ be any solution to (1.1), that is (1.1a), (1.1b), (1.1c), on the half-open interval $[0, \tau_1)$ with $0 < \tau_1 < \infty$. Set $\phi = \alpha_1 + \alpha_2 + \alpha_3$ and suppose $\phi(0) < 0$. Then

$$\phi(\tau) \leq -|\phi(0)| / \left(1 + \frac{1}{3} |\phi(0)| \tau\right) < 0 \quad \text{for all } \tau \in [0, \tau_1) \quad (1.4)$$

by (vi). Consequently, $|\beta_1 \beta_2 \beta_3|$ is bounded, by (iv), and ϕ is bounded below, by (v), on $[0, \tau_1)$. The constraint (1.1c) implies that⁵ $(\alpha_1)^2 + (\alpha_2)^2 + (\alpha_3)^2 \leq 6 |\beta_1 \beta_2 \beta_3|^{2/3} + \phi^2$ is bounded. Now (1.1b) implies that $(\beta_1)^2 + (\beta_2)^2 + (\beta_3)^2$ is bounded. Therefore, solutions to (1.1) with $\phi(0) < 0$ can be extended to $[0, \infty)$. The solutions considered in this paper belong to this general class. We are especially interested in their $\tau \rightarrow +\infty$ asymptotics.

⁴ $(\beta_1)^2 + (\beta_2)^2 + (\beta_3)^2 - 2\beta_2\beta_3 - 2\beta_3\beta_1 - 2\beta_1\beta_2 + 3|\beta_1\beta_2\beta_3|^{2/3} \geq 0$ holds for all $\beta_1, \beta_2, \beta_3 \in \mathbb{R}$, see [HU1]. The only nontrivial cases are $\beta_1, \beta_2, \beta_3 > 0$ or $\beta_1, \beta_2, \beta_3 < 0$. In these cases, the inequality is a direct consequence of the polynomial identity

$$x^6 + y^6 + z^6 - 2y^3z^3 - 2z^3x^3 - 2x^3y^3 + 3x^2y^2z^2 = \frac{1}{2} (x^2 + y^2 + z^2 + yz + zx + xy) \left((y-z)^2(y+z-x)^2 + (z-x)^2(z+x-y)^2 + (x-y)^2(x+y-z)^2 \right)$$


⁵ Use $2(\alpha_2\alpha_3 + \alpha_3\alpha_1 + \alpha_1\alpha_2) = \phi^2 - (\alpha_1)^2 - (\alpha_2)^2 - (\alpha_3)^2$.

For every solution to (1.1) with $\phi(0) < 0$, as in the last paragraph, the half-infinite interval $[0, \infty)$ actually corresponds to a *finite* physical duration of the associated spatially homogeneous vacuum spacetime (given in Proposition 2.1). In fact, an increasing affine parameter along the timelike geodesics orthogonal to the level sets of τ is given by $\tau \mapsto \int_0^\tau \exp(\frac{1}{2} \int_0^s \phi) ds$, with uniform upper bound $6|\phi(0)|^{-1}$, by (1.4).

In this paper, we consider only solutions to (1.1) for which $\beta_1, \beta_2, \beta_3 \neq 0$ (also called Bianchi VIII or IX models). We now give an informal description of the solutions that we construct, the phenomenological picture of [BKL1]. The structure of each of these solutions is described by three sequences of compact subintervals $(\mathcal{I}_j)_{j \geq 1}$, $(\mathcal{B}_j)_{j \geq 1}$, $(\mathcal{S}_j)_{j \geq 1}$ of $[0, \infty)$, for which:

- (a.1) The left endpoint of $\mathcal{I}_1$ is the origin, and the right endpoint of $\mathcal{I}_j$, henceforth denoted τ_j , coincides with the left endpoint of $\mathcal{I}_{j+1}$, for all $j \geq 1$. Set $\tau_0 = 0$.
- (a.2) $\bigcup_{j \geq 1} \mathcal{I}_j = [0, \infty)$, that is, $\lim_{j \rightarrow +\infty} \tau_j = +\infty$.
- (a.3) $\mathcal{B}_j$ is contained in the interior of $\mathcal{I}_j$, and $0 < |\mathcal{B}_j| \ll |\mathcal{I}_j|$, for all $j \geq 1$.
- (a.4) $\mathcal{S}_j$ is the closed interval of all points between $\mathcal{B}_j$ and $\mathcal{B}_{j+1}$, for all $j \geq 1$.

Here is a picture:



- (c.4) By the last three items and (b.4), there is at most one point in $\mathcal{S}_j$ where $|\beta_{\mathbf{a}(j)}| = |\beta_{\mathbf{a}(j+1)}|$. By (b.1), (b.2), there is such a point, because $|\beta_{\mathbf{a}(j)}|$ is going from not small to small on $\mathcal{S}_j$, and $|\beta_{\mathbf{a}(j+1)}|$ is going from small to not small on $\mathcal{S}_j$. By convention, this point is τ_j .

Separately on each interval $\mathcal{I}_j$, $j \geq 1$ (in particular on $\mathcal{B}_j \subset \mathcal{I}_j$):

- (d.1) $\alpha_{\mathbf{a}(j)} + \alpha_{\mathbf{b}(j)}$ and $\alpha_{\mathbf{a}(j)} + \alpha_{\mathbf{c}(j)}$ are essentially constant, by (1.1a), (b.1), and they are both negative, by (c.1), (c.2). Also, $\log |\beta_{\mathbf{a}(j)}\beta_{\mathbf{b}(j)}|$, $\log |\beta_{\mathbf{a}(j)}\beta_{\mathbf{c}(j)}|$ are essentially linear functions with slopes $\alpha_{\mathbf{a}(j)} + \alpha_{\mathbf{b}(j)}$ and $\alpha_{\mathbf{a}(j)} + \alpha_{\mathbf{c}(j)}$, by (1.1b).
(d.2) Essentially $(\alpha_{\mathbf{a}(j)} + \alpha_{\mathbf{b}(j)})(\alpha_{\mathbf{a}(j)} + \alpha_{\mathbf{c}(j)}) = (\alpha_{\mathbf{a}(j)})^2 + (\beta_{\mathbf{a}(j)})^2$, by (1.1c). Since the left hand side is essentially constant by (d.1), so is the right hand side.
(d.3) By (d.1), it only remains to understand the behavior of $\alpha_{\mathbf{a}(j)}$, $\beta_{\mathbf{a}(j)}$. By (1.1a), we essentially have

$$\frac{d}{d\tau}\alpha_{\mathbf{a}(j)} = -(\beta_{\mathbf{a}(j)})^2 \quad \frac{d}{d\tau}\beta_{\mathbf{a}(j)} = \alpha_{\mathbf{a}(j)}\beta_{\mathbf{a}(j)} \quad (1.5)$$

A special solution is $\alpha_{\mathbf{a}(j)} = -\tanh \tau$ and $\beta_{\mathbf{a}(j)} = \pm \operatorname{sech} \tau = \pm(\cosh \tau)^{-1}$. The general solution is obtained from the special solution by applying the affine symmetry transformation (ii) above, with $p > 0$. Since $\mathcal{B}_j$ is essentially the interval on which $|\beta_{\mathbf{a}(j)}|$ is not small, see (b.2), we must have $p \sim |\mathcal{B}_j|^{-1}$ (here $\sim$ means “same order of magnitude”). See [BKL1], Section 3, in particular pages 534 and 535.

- (d.4) Recall (c.1). By (d.3), we have $\alpha_{\mathbf{a}(j)}|_{\mathcal{S}_{j-1}} = -\alpha_{\mathbf{a}(j)}|_{\mathcal{S}_j}$, since the hyperbolic tangent just flips the sign. Therefore, by (d.1), the net change across $\mathcal{B}_j$ of the components of α , from right to left, is given by

$$\begin{aligned} \alpha_{\mathbf{a}(j)}|_{\mathcal{S}_{j-1}} &= \alpha_{\mathbf{a}(j)}|_{\mathcal{S}_j} - 2\alpha_{\mathbf{a}(j)}|_{\mathcal{S}_j} \\ \alpha_{\mathbf{b}(j)}|_{\mathcal{S}_{j-1}} &= \alpha_{\mathbf{b}(j)}|_{\mathcal{S}_j} + 2\alpha_{\mathbf{a}(j)}|_{\mathcal{S}_j} \\ \alpha_{\mathbf{c}(j)}|_{\mathcal{S}_{j-1}} &= \alpha_{\mathbf{c}(j)}|_{\mathcal{S}_j} + 2\alpha_{\mathbf{a}(j)}|_{\mathcal{S}_j} \end{aligned}$$

These equations make sense only for $j \geq 2$, since $\mathcal{S}_0$ has not been defined.

In this paper, we turn the heuristic picture of [BKL1], sketched above, into a mathematically rigorous one, globally on $[0, \infty)$, for a generic class of solutions. The first step is to construct a discrete dynamical system, that maps the state $\Phi(\tau_j)$ to the state $\Phi(\tau_{j-1})$ at the earlier time $\tau_{j-1} < \tau_j$, for all $j \geq 1$. That is, the construction proceeds from right-to-left, beginning at $\tau = +\infty$. We refer to the discrete dynamical system maps as transfer maps.

For each $j \geq 0$, two components of $\beta(\tau_j)$ have the same absolute value, see (c.4), and $\Phi(\tau_j)$ satisfies the constraint (1.1c). Therefore, the states of the discrete dynamical system have 4 continuous degrees of freedom. By the symmetry (ii), the transfer maps commute with rescalings. Taking the quotient, one obtains a 3-dimensional discrete dynamical system. The three “dimensionless” quantities that we use to parametrize the discrete states are denoted $\mathbf{f}_j = (\mathbf{h}_j, w_j, q_j)$. Morally, they are interpreted as follows:

- $\mathbf{h}_j \sim |\mathcal{B}_j|/|\mathcal{I}_j| > 0$. In the billiard picture of [Mis], it is the dimensionless ratio of the collision and free-motion times. By (a.3), one has $0 < \mathbf{h}_j \ll 1$. In fact, $\mathbf{h}_j$ is the all-important small parameter in our construction. It goes to zero rapidly as $j \rightarrow \infty$. This is necessary for us to make a global construction on $[0, \infty)$. The precise rate depends on the sequence $(k_n)_{n \geq 1}$. The rate is the same as in Proposition 4.4, up to even smaller corrections.

- The components of α are essentially constant on $\mathcal{S}_j$ and subject to the reduced constraint equation in (c.2). Thus, modulo the scaling symmetry (ii), only one degree of freedom is required to parametrize $\alpha|_{\mathcal{S}_j}$. We use $w_j \approx -(\alpha_{\mathbf{b}(j)} / (\alpha_{\mathbf{a}(j)} + \alpha_{\mathbf{b}(j)}))|_{\mathcal{S}_j}$. By (c.2) and (c.3), we have $w_j > 0$. The left-to-right discrete dynamics of w_j (which is opposite to the right-to-left direction of our transfer maps) is closely related to a variant of the Gauss map, sometimes referred to as the *BKL map* or *Kasner map*.
- The meaning of q_j will be explained in a more indirect way. As pointed out above, the left-to-right dynamics of w_j is related to the Gauss map, which is a non-invertible left-shift, see (1.3). The non-invertibility of the Gauss map seems to be at odds with the invertible dynamics of the system of ordinary differential equations (1.1). The parameter q_j is introduced so that the *joint* left-to-right discrete dynamics of (w_j, q_j) is closely related to the left-shift on *two-sided* sequences $(k_n)_{n \in \mathbb{Z}}$ of strictly positive integers, which is invertible. Accordingly, the right-to-left transfer maps are related to the right shift on two-sided sequences $(k_n)_{n \in \mathbb{Z}}$.

This concludes the informal discussion. We emphasize that the notation used above is specific to the introduction. In particular, $(\mathcal{I}_j)_{j \geq 1}$, $(\mathcal{B}_j)_{j \geq 1}$, $(\mathcal{S}_j)_{j \geq 1}$ do not appear in the main text. Starting from Section 2, all the notation is introduced from scratch.

We now state simplified, self-contained versions of our results. References to their stronger counterparts are given. Here is a short guide:

Definition 1.1 (equivalent to Definition 3.12). Introduces the state vectors $\Phi_*(\pi, \mathbf{f}, \sigma_*)$ of the 3-dimensional discrete dynamical system. The dynamics of the signature vector σ_* is trivial, by (iii), but it affects the dynamics of $(\pi, \mathbf{f})$ in a non-trivial way.

Definition 1.2 (this is Definition 3.16). Introduces explicit maps $\mathcal{P}_L, \mathcal{Q}_L, \lambda_L$ that turn out to be very good approximations to the transfer maps. It is shown in Section 4 that iterates of $\mathcal{Q}_L$ can be understood in terms of the Gauss map / continued fractions and, by a change of variables, in terms of solutions to certain linear equations.

Definition 3.19 (only in the main text). The essential smallness condition on $\mathbf{h} > 0$ is quantitatively encoded in an open subset $\mathcal{F} \subset (0, 1) \times (0, \infty) \times ((0, \infty) \setminus \{1\})$. It determines the domain of definition of the transfer maps.

Proposition 1.1 (slimmed-down version of Proposition 3.3). It asserts the existence of transfer maps. The pair $(\mathcal{P}_L, \Pi)$ and the triple $(\mathcal{P}_L, \Pi, \Lambda)$ constitute the transfer maps for the 3-dimensional and 4-dimensional systems, respectively, and they are very close to $(\mathcal{P}_L, \mathcal{Q}_L)$ and $(\mathcal{P}_L, \mathcal{Q}_L, \lambda_L)$. Explicit error bounds and precise estimates for the transfer solution appear only in the full version, Proposition 3.3.

Theorem 1.1 (simplified version of Theorems 6.2, 6.3). Gives a generic class of iterates to $(\mathcal{P}_L, \Pi)$ that are super-exponentially close to iterates of $(\mathcal{P}_L, \mathcal{Q}_L)$. That is, it asserts the existence of solutions to the 3-dimensional discrete dynamical system.

The overview is as follows. Every solution to the 3-dimensional discrete dynamical system as in Theorem 1.1 can be lifted to a unique solution to the 4-dimensional discrete dynamical system, up to an overall scale, through the map Λ in Proposition 1.1. This solution corresponds to the sequence of states $(\Phi(\tau_j))_{j \geq 0}$ in the informal discussion. Proposition 1.1 gives solutions to (1.1) on compact intervals that connect next-neighbor states. Symmetry (ii) is used to translate these compact intervals and place them next to each other, beginning at $\tau = 0$, just like the $(\mathcal{I}_j)_{j \geq 1}$ in the informal discussion. As in (a.2) of the informal discussion, the union of these intervals is indeed $[0, \infty)$, and a semi-global solution to (1.1) is obtained. To see this, denote the states by $\lambda_j \Phi_*(\pi_j, \mathbf{g}_j, \sigma_*)$ with $\lambda_j > 0$ and $\pi_j \in S_3$ and $\mathbf{g}_j = (\mathbf{h}'_j, w'_j, q'_j) \in \mathcal{F}$, where $j \geq 0$. One has $\lambda_j = \Lambda[\pi_{j+1}, \sigma_*](\mathbf{g}_{j+1})\lambda_{j+1} \geq \lambda_{j+1}$ by the definition of Λ and

$\mathbf{h}'_j \in (0, 1)$ by the definition of $\mathcal{F}$. In particular, the sequence of products $(\lambda_j \mathbf{h}'_j)_{j \geq 0}$ is bounded from above by $\lambda_0 > 0$. By Proposition 1.1, the length of each of the intervals is bounded from below by $(2\lambda_0)^{-1} > 0$.

Definition 1.1 (State vectors). Let $\pi = (\mathbf{a}, \mathbf{b}, \mathbf{c}) \in S_3$ and $\sigma_* \in \{-1, +1\}^3$ and $\mathbf{f} = (\mathbf{h}, w, q) \in (0, \infty)^2 \times \mathbb{R}$. Let $\Phi_* = \Phi_*(\pi, \mathbf{f}, \sigma_*) = \alpha \oplus \beta \in \mathbb{R}^3 \oplus \mathbb{R}^3$ be the vector given by $(\text{sgn } \beta_1, \text{sgn } \beta_2, \text{sgn } \beta_3) = \sigma_*$ and by

$$\begin{aligned} \alpha_{\mathbf{a}} &= -1 & \mathbf{h} \log |\tfrac{1}{2}\beta_{\mathbf{a}}| &= -\frac{1+w}{1+2w}(1 + \mathbf{h} \log 2) \\ \alpha_{\mathbf{b}} &= \frac{w}{1+w} & \mathbf{h} \log |\tfrac{1}{2}\beta_{\mathbf{b}}| &= -\frac{1+w}{1+2w}(1 + \mathbf{h} \log 2) \\ \alpha_{\mathbf{c}} &= -w - \mu & \mathbf{h} \log |\tfrac{1}{2}\beta_{\mathbf{c}}| &= -(1+w)q - \frac{w(1+w)}{1+2w} - \frac{1+3w+w^2}{1+2w} \mathbf{h} \log 2 \end{aligned}$$

where $\mu = \mu(\pi, \mathbf{f}, \sigma_*) \in \mathbb{R}$ is uniquely determined by requiring that (1.1c) holds.

Definition 1.2 (Approximate transfer maps). Introduce three maps

$$\begin{aligned} \mathcal{P}_L : S_3 \times (0, \infty)^3 &\rightarrow S_3 & ((\mathbf{a}, \mathbf{b}, \mathbf{c}), \mathbf{f}) &\mapsto (\mathbf{a}', \mathbf{b}', \mathbf{c}') \\ \mathcal{Q}_L : (0, \infty)^3 &\rightarrow (0, \infty)^2 \times \mathbb{R} & \mathbf{f} &\mapsto (\mathbf{h}_L, w_L, q_L) \\ \lambda_L : (0, \infty)^3 &\rightarrow (0, \infty) & \mathbf{f} &\mapsto \lambda_L \end{aligned}$$

where $\mathbf{f} = (\mathbf{h}, w, q)$ and $q_L = \text{num}1_L / \text{den}_L$ and $\mathbf{h}_L = \text{num}2_L / \text{den}_L$, and:

• if $q \leq 1$:

$$\begin{aligned} (\mathbf{a}', \mathbf{b}', \mathbf{c}') &= (\mathbf{c}, \mathbf{a}, \mathbf{b}) & \text{num}1_L &= (1+w)(1-q) - \mathbf{h} \log 2 + \mathbf{h}w \log(2+w) \\ w_L &= \frac{1}{1+w} & \text{num}2_L &= \mathbf{h}(2+w) \\ \lambda_L &= 2+w & \text{den}_L &= (1+w)(q - \mathbf{h} \log 2) + \mathbf{h}(3+w) \log(2+w) \end{aligned}$$

• if $q > 1$:

$$\begin{aligned} (\mathbf{a}', \mathbf{b}', \mathbf{c}') &= (\mathbf{b}, \mathbf{a}, \mathbf{c}) & \text{num}1_L &= (1+w)(q-1 - \mathbf{h} \log 2) - \mathbf{h}w \log \frac{2+w}{1+w} \\ w_L &= 1+w & \text{num}2_L &= \mathbf{h}(2+w) \\ \lambda_L &= \frac{2+w}{1+w} & \text{den}_L &= (1+w) - \mathbf{h} \log 2 + \mathbf{h}(3+2w) \log \frac{2+w}{1+w} \end{aligned}$$

Observe that $\text{den}_L > 0$.

Proposition 1.1 (Transfer maps). Fix $\sigma_* \in \{-1, +1\}^3$ and $\pi = (\mathbf{a}, \mathbf{b}, \mathbf{c}) \in S_3$. There exist maps⁶

$$\Pi[\pi, \sigma_*] : \mathcal{F} \rightarrow (0, \infty)^2 \times \mathbb{R} \quad \text{and} \quad \Lambda[\pi, \sigma_*] : \mathcal{F} \rightarrow [1, \infty)$$

such that for every $\lambda > 0$ and $\mathbf{f} = (\mathbf{h}, w, q) \in \mathcal{F}$, the solution to (1.1) starting at $\lambda \Phi_*(\pi, \mathbf{f}, \sigma_*)$ at time 0 passes through $\lambda' \Phi_*(\pi', \mathbf{f}', \sigma_*)$ at an earlier time $\tau' < 0$, with $\frac{1}{2} \leq \mathbf{h} \lambda |\tau'| \leq 3$. Here $\mathbf{f}' = \Pi[\pi, \sigma_*](\mathbf{f})$ and $\lambda' = \lambda \Lambda[\pi, \sigma_*](\mathbf{f})$ and $\pi' = \mathcal{P}_L(\pi, \mathbf{f})$. Schematically, the transition is

$$\lambda \Lambda[\pi, \sigma_*](\mathbf{f}) \Phi_*(\mathcal{P}_L(\pi, \mathbf{f}), \Pi[\pi, \sigma_*](\mathbf{f}), \sigma_*) \quad \longleftarrow \quad \lambda \Phi_*(\pi, \mathbf{f}, \sigma_*)$$

Furthermore (informal): Π and Λ are approximated by the maps $\mathcal{Q}_L$ and λ_L , with errors that go to zero exponentially as $\mathbf{h} \downarrow 0$ (for fixed w, q). See Proposition 3.3

⁶ Caution: The maps Π cannot immediately be iterated / composed, because $(0, \infty)^2 \times \mathbb{R} \not\subset \mathcal{F}$.

Theorem 1.1. Fix $\sigma_* \in \{-1, +1\}^3$ and $\pi_0 \in S_3$. Fix constants $\mathbf{D} \geq 1$, $\gamma \geq 0$. Suppose the vector $\mathbf{f}_0 = (\mathbf{h}_0, w_0, q_0) \in (0, \infty)^3$ satisfies

- (i) $w_0 \in (0, 1) \setminus \mathbb{Q}$ and $q_0 \in (0, \infty) \setminus \mathbb{Q}$.
- (ii) $k_n \leq \mathbf{D} \max\{1, n\}^\gamma$ for all $n \geq -2$, where the two-sided sequence of strictly positive integers $(k_n)_{n \in \mathbb{Z}}$ is given by

$$(1 + q_0)^{-1} = \langle k_0, k_{-1}, k_{-2}, \dots \rangle \quad w_0 = \langle k_1, k_2, k_3, \dots \rangle$$

- (iii) $0 < \mathbf{h}_0 < \mathbf{A}^\#$ where $\mathbf{A}^\# = \mathbf{A}^\#(\mathbf{D}, \gamma) = 2^{-56} \mathbf{D}^{-4} (4(\gamma + 1))^{-4(\gamma + 1)}$.

Then $\mathbf{f}_0$ and π_0 are the first elements of a unique sequence $(\mathbf{f}_j)_{j \geq 0}$ in $\mathcal{F}$ and a unique sequence $(\pi_j)_{j \geq 0}$ in S_3 , respectively, with $\pi_j = \mathcal{P}_L(\pi_{j+1}, \mathbf{f}_{j+1})$ and $\mathbf{f}_j = \mathcal{Q}_L(\mathbf{f}_{j+1})$ for all $j \geq 0$. Furthermore, there exists a sequence $(\mathbf{g}_j)_{j \geq 0}$ in $\mathcal{F}$ such that for all $j \geq 0$,

$$\mathbf{g}_j = \Pi[\pi_{j+1}, \sigma_*](\mathbf{g}_{j+1}) \quad \text{and} \quad \pi_j = \mathcal{P}_L(\pi_{j+1}, \mathbf{g}_{j+1})$$

and, with $\rho_+ = \frac{1}{2}(1 + \sqrt{5})$,

$$\|\mathbf{g}_j - \mathbf{f}_j\|_{\mathbb{R}^3} \leq \exp\left(-\frac{1}{\mathbf{h}_0} \mathbf{A}^\# \rho_+^{((\mathbf{D}^{-1}j)^{1/(\gamma+1)})}\right)$$

If $\gamma > 1$ and $\mathbf{D} > \frac{1}{\log 2} \frac{\gamma}{\gamma-1}$, then the set of all vectors $\mathbf{f}_0 \in (0, \infty)^3$ that satisfy (i), (ii), (iii) has positive Lebesgue measure.

The class of solutions that we construct is generic in the sense of the last sentence of Theorem 1.1. It would be desirable to have a stronger genericity statement, namely a genericity statement for “the $\mathbf{g}_0$ rather than the $\mathbf{f}_0$ ”.

For the causal structure and particle horizons, see Proposition 2.2 and Section 7.

It is a pleasure to thank J. Fröhlich, G.M. Graf and T. Spencer for their support and encouragement.

2. Spatially homogeneous vacuum spacetimes

Proposition 2.1. Let $\alpha \oplus \beta : (\tau_0, \tau_1) \rightarrow \mathbb{R}^3 \oplus \mathbb{R}^3$ be a solution to (1.1) and let $\Omega \subset \mathbb{R}^3$ be open, with Cartesian coordinates $\mathbf{x} = (x^1, x^2, x^3)$. Fix any $\tau_* \in (\tau_0, \tau_1)$ and let

$$v_1 = \sum_{\mu=1}^3 v_1^\mu(\mathbf{x}) \frac{\partial}{\partial x^\mu} \quad v_2 = \sum_{\mu=1}^3 v_2^\mu(\mathbf{x}) \frac{\partial}{\partial x^\mu} \quad v_3 = \sum_{\mu=1}^3 v_3^\mu(\mathbf{x}) \frac{\partial}{\partial x^\mu}$$

be three smooth vector fields on Ω that are a frame at each point and satisfy

$$[v_j, v_k] = \beta_i(\tau_*) v_i \quad \text{on } \Omega$$

for all $(\mathbf{i}, \mathbf{j}, \mathbf{k}) \in \mathcal{C} \stackrel{\text{def}}{=} \{(1, 2, 3), (2, 3, 1), (3, 1, 2)\}$. Introduce

$$\begin{aligned} e_0 &= e^{\zeta(\tau)} \frac{\partial}{\partial \tau} & e_i &= e^{\zeta_i(\tau)} v_i & \mathbf{i} &= 1, 2, 3 \\ \zeta(\tau) &= \zeta_1(\tau) + \zeta_2(\tau) + \zeta_3(\tau) & \zeta_i(\tau) &= -\frac{1}{2} \int_{\tau_*}^{\tau} ds \alpha_i(s) & \mathbf{i} &= 1, 2, 3 \end{aligned}$$

on the domain $(\tau_0, \tau_1) \times \Omega \subset \mathbb{R}^4$. Then, the Lorentzian metric g with inverse

$$g^{-1} = -e_0 \otimes e_0 + e_1 \otimes e_1 + e_2 \otimes e_2 + e_3 \otimes e_3$$

is a solution to the vacuum Einstein equations $\text{Ric}(g) = 0$ on $(\tau_0, \tau_1) \times \Omega$.

Proof. In this proof, everywhere $(\mathbf{i}, \mathbf{j}, \mathbf{k}) \in \mathcal{C}$. It follows from $\frac{d}{d\tau}e^{-2\zeta_{\mathbf{i}}} = \alpha_{\mathbf{i}}e^{-2\zeta_{\mathbf{i}}}$ and $\zeta_{\mathbf{i}}(\tau_*) = 0$ and (1.1b) that $\beta_{\mathbf{i}}(\tau) = \beta_{\mathbf{i}}(\tau_*)e^{-2\zeta_{\mathbf{i}}(\tau)}$. Now, by direct calculation,

$$[e_0, e_{\mathbf{i}}] = -\frac{1}{2}e^{\zeta}\alpha_{\mathbf{i}}e_{\mathbf{i}} \quad [e_{\mathbf{j}}, e_{\mathbf{k}}] = e^{\zeta}\beta_{\mathbf{i}}e_{\mathbf{i}}$$

Let ∇ be the Levi-Civita connection associated to g . Then, for all $a, b, c = 0, 1, 2, 3$,

$$g(\nabla_{e_a}e_b, e_c) = \frac{1}{2}\left(g([e_a, e_b], e_c) - g([e_b, e_c], e_a) + g([e_c, e_a], e_b)\right)$$

By direct calculation,

$$\begin{aligned} \nabla_{e_0}e_0 &= 0 & \nabla_{e_{\mathbf{i}}}e_{\mathbf{i}} &= \frac{1}{2}e^{\zeta}\alpha_{\mathbf{i}}e_0 \\ \nabla_{e_0}e_{\mathbf{i}} &= 0 & \nabla_{e_{\mathbf{j}}}e_{\mathbf{k}} &= \frac{1}{2}e^{\zeta}(+\beta_{\mathbf{i}} - \beta_{\mathbf{j}} + \beta_{\mathbf{k}})e_{\mathbf{i}} \\ \nabla_{e_{\mathbf{i}}}e_0 &= \frac{1}{2}e^{\zeta}\alpha_{\mathbf{i}}e_{\mathbf{i}} & \nabla_{e_{\mathbf{k}}}e_{\mathbf{j}} &= \frac{1}{2}e^{\zeta}(-\beta_{\mathbf{i}} - \beta_{\mathbf{j}} + \beta_{\mathbf{k}})e_{\mathbf{i}} \end{aligned}$$

and

$$\begin{aligned} \text{Riem}(e_{\mathbf{i}}, e_{\mathbf{j}}, e_{\mathbf{i}}, e_{\mathbf{j}}) &= \frac{1}{4}e^{2\zeta}\left((+\beta_{\mathbf{i}} - \beta_{\mathbf{j}} - \beta_{\mathbf{k}})(+\beta_{\mathbf{i}} - \beta_{\mathbf{j}} + \beta_{\mathbf{k}}) + 2\beta_{\mathbf{k}}(+\beta_{\mathbf{i}} + \beta_{\mathbf{j}} - \beta_{\mathbf{k}}) + \alpha_{\mathbf{i}}\alpha_{\mathbf{j}}\right) \\ \text{Riem}(e_0, e_{\mathbf{a}}, e_{\mathbf{i}}, e_{\mathbf{j}}) &= \frac{1}{4}e^{2\zeta}\delta_{\mathbf{a}\mathbf{k}}\left((-\beta_{\mathbf{i}} + \beta_{\mathbf{j}} - \beta_{\mathbf{k}})\alpha_{\mathbf{i}} + (+\beta_{\mathbf{i}} - \beta_{\mathbf{j}} - \beta_{\mathbf{k}})\alpha_{\mathbf{j}} + 2\alpha_{\mathbf{k}}\beta_{\mathbf{k}}\right) \\ \text{Riem}(e_0, e_{\mathbf{a}}, e_0, e_{\mathbf{i}}) &= -\frac{1}{4}e^{2\zeta}\delta_{\mathbf{a}\mathbf{i}}\left(2\frac{d}{d\tau}\alpha_{\mathbf{i}} - (\alpha_{\mathbf{j}} + \alpha_{\mathbf{k}})\alpha_{\mathbf{i}}\right) \end{aligned}$$

Furthermore, $\text{Riem}(e_{\mathbf{a}}, e_{\mathbf{b}}, e_{\mathbf{c}}, e_{\mathbf{d}}) = 0$ unless $\{\mathbf{a}, \mathbf{b}\} = \{\mathbf{c}, \mathbf{d}\}$ with $\mathbf{a} \neq \mathbf{b}$. The Riemann curvature tensor is completely specified by these identities and by its algebraic symmetries. It follows that

$$\begin{aligned} \text{Ric}(e_0, e_0) &= -\frac{1}{2}e^{2\zeta}\frac{d}{d\tau}(\alpha_1 + \alpha_2 + \alpha_3) + \frac{1}{2}e^{2\zeta}(\alpha_2\alpha_3 + \alpha_3\alpha_1 + \alpha_1\alpha_2) \\ \text{Ric}(e_0, e_{\mathbf{i}}) &= 0 \\ \text{Ric}(e_{\mathbf{i}}, e_{\mathbf{i}}) &= +\frac{1}{2}e^{2\zeta}\frac{d}{d\tau}\alpha_{\mathbf{i}} + \frac{1}{2}e^{2\zeta}\left((\beta_{\mathbf{i}})^2 - (\beta_{\mathbf{j}})^2 - (\beta_{\mathbf{k}})^2 + 2\beta_{\mathbf{j}}\beta_{\mathbf{k}}\right) \\ \text{Ric}(e_{\mathbf{j}}, e_{\mathbf{k}}) &= 0 \end{aligned}$$

The right hand sides of the first and third equation vanish by (1.1a) and (1.1c). $\square$

Proposition 2.2. *In the context of Proposition 2.1, let $\gamma : (\tau'_0, \tau'_1) \rightarrow (\tau_0, \tau_1) \times \Omega$ be a smooth curve given by $\gamma(\tau) = (\tau, \gamma^\sharp(\tau))$, where $\gamma^\sharp$ is a curve on Ω . Let $g^\sharp$ be the Riemannian metric on Ω defined by $g^\sharp(v_{\mathbf{a}}, v_{\mathbf{b}}) = \delta_{\mathbf{a}\mathbf{b}}$ for all $\mathbf{a}, \mathbf{b} = 1, 2, 3$. If γ is non-spacelike with respect to g , then the length of $\gamma^\sharp$ with respect to $g^\sharp$ is bounded by*

$$\text{Length}_{g^\sharp}(\gamma^\sharp) \leq \int_{\tau'_0}^{\tau'_1} d\tau \max_{(\mathbf{i}, \mathbf{j}, \mathbf{k}) \in \mathcal{C}} e^{-\zeta_{\mathbf{j}} - \zeta_{\mathbf{k}}} \quad (2.1)$$

The integral on the right hand side may be divergent.

Proof. Write the velocity $\frac{d}{d\tau}\gamma$ as

$$\frac{\partial}{\partial\tau} + \sum_{i=1}^3 X^i v_i = e^{-\zeta} e_0 + \sum_{i=1}^3 X^i e^{-\zeta_i} e_i$$

with smooth coefficients $X^i = X^i(\tau)$. By assumption, γ is non-spacelike:

$$0 \geq g\left(\frac{d}{d\tau}\gamma, \frac{d}{d\tau}\gamma\right) = -e^{-2\zeta} + \sum_{i=1}^3 (X^i)^2 e^{-2\zeta_i}$$

Consequently, $\sum_{i=1}^3 (X^i)^2 \leq \max_{(i,j,k) \in \mathcal{C}} e^{-2\zeta_j - 2\zeta_k}$. Now, the claim follows from

$$\text{Length}_{g^\#}(\gamma^\#) = \int_{\tau'_0}^{\tau'_1} d\tau \sqrt{g^\#(\frac{d}{d\tau}\gamma^\#, \frac{d}{d\tau}\gamma^\#)} = \int_{\tau'_0}^{\tau'_1} d\tau \sqrt{\sum_{i=1}^3 (X^i)^2}$$

□

3. Construction of the transfer maps

Let $(\tau_0, \tau_1) \subset \mathbb{R}$ be a finite or infinite open interval, parametrized by $\tau \in (\tau_0, \tau_1)$ (“time”). In this paper, the unknown field is a vector valued map $\Phi \in C^\infty((\tau_0, \tau_1), \mathbb{R}^6)$:

$$\Phi = \alpha[\Phi] \oplus \beta[\Phi] : (\tau_0, \tau_1) \rightarrow \mathbb{R}^3 \oplus \mathbb{R}^3 \quad (3.1)$$

If no confusion can arise, we just write $\Phi = \alpha \oplus \beta$.

Definition 3.1. To every field $\Phi = \alpha \oplus \beta \in C^\infty((\tau_0, \tau_1), \mathbb{R}^6)$, every constant $\mathbf{h} > 0$ and every $n \in \mathbb{R}^3$, associate a field

$$\mathbf{a}[\Phi, \mathbf{h}, n] \oplus \mathbf{b}[\Phi, \mathbf{h}, n] \oplus c[\Phi, \mathbf{h}, n] : (\tau_0, \tau_1) \rightarrow \mathbb{R}^3 \oplus \mathbb{R}^3 \oplus \mathbb{R}$$

by

$$\mathbf{a}_i[\Phi, \mathbf{h}, n] = -\mathbf{h} \frac{d}{d\tau} \alpha_i - (n_i \beta_i)^2 + (n_j \beta_j - n_k \beta_k)^2 \quad (3.2a)$$

$$\mathbf{b}_i[\Phi, \mathbf{h}, n] = -\mathbf{h} \frac{d}{d\tau} \beta_i + \beta_i \alpha_i \quad (3.2b)$$

$$c[\Phi, \mathbf{h}, n] = \sum_{(i,j,k) \in \mathcal{C}} (\alpha_j \alpha_k - (n_i \beta_i)^2 + 2n_j n_k \beta_j \beta_k) \quad (3.2c)$$

for all $(i, j, k) \in \mathcal{C}$. For later use, it is convenient to introduce, for all $m, n \in \mathbb{R}^3$,

$$\begin{aligned} \mathbf{a}_i[\Phi, \mathbf{h}, n, m] &= \mathbf{a}_i[\Phi, \mathbf{h}, n] - \mathbf{a}_i[\Phi, \mathbf{h}, m] \\ &= -(n_i \beta_i)^2 + (n_j \beta_j - n_k \beta_k)^2 + (m_i \beta_i)^2 - (m_j \beta_j - m_k \beta_k)^2 \end{aligned} \quad (3.3)$$

Definition 3.2.

$$B_1 = (1, 0, 0) \quad B_2 = (0, 1, 0) \quad B_3 = (0, 0, 1) \quad Z = (1, 1, 1)$$

These vectors will play the role of the vector $n \in \mathbb{R}^3$ that appears in Definition 3.1.

Proposition 3.1 (Global symmetries). Let $\chi : (\tau_0, \tau_1) \rightarrow (\tau'_0, \tau'_1)$ be a linear diffeomorphism between finite or infinite intervals, $\chi(\tau) = p\tau + q$ with $p > 0$, and let $A > 0$ be a constant. Then

$$(\mathbf{a}, \mathbf{b}, c) \left[A(\Phi \circ \chi), \frac{1}{p} A \mathbf{h}, n \right] = A^2 \left((\mathbf{a}, \mathbf{b}, c)[\Phi, \mathbf{h}, n] \circ \chi \right)$$

for all fields $\Phi = \alpha \oplus \beta \in C^\infty((\tau'_0, \tau'_1), \mathbb{R}^6)$, all constants $\mathbf{h} > 0$ and all $n \in \mathbb{R}^3$.

Corollary 3.1. *In Proposition 3.1, the field $(\mathbf{a}, \mathbf{b}, c)[A(\Phi \circ \chi), \frac{1}{p}A\mathbf{h}, n]$ vanishes identically on (τ_0, τ_1) if and only if $(\mathbf{a}, \mathbf{b}, c)[\Phi, \mathbf{h}, n]$ vanishes identically on (τ'_0, τ'_1) .*

Remark 3.1. The equations $(\mathbf{a}, \mathbf{b}, c)[\Phi, 1, Z] = 0$ are identical to (1.1). The equations $(\mathbf{a}, \mathbf{b}, c)[\Phi, \mathbf{h}, Z] = 0$ are equivalent to (1.1), for any $\mathbf{h} > 0$, by Corollary 3.1.

Proposition 3.2. *Recall Definition 3.1. For all $\Phi = \alpha \oplus \beta \in C^\infty((\tau_0, \tau_1), \mathbb{R}^6)$, all $\mathbf{h} > 0$ and all $n \in \mathbb{R}^3$, we have*

$$0 = -\mathbf{h} \frac{d}{d\tau} c + \sum_{(i,j,k) \in \mathcal{C}} \left(-\alpha_j \mathbf{a}_k - \alpha_k \mathbf{a}_j + 2(n_i)^2 \beta_i \mathbf{b}_i - 2n_j n_k \beta_j \mathbf{b}_k - 2n_j n_k \beta_k \mathbf{b}_j \right) \quad (3.4)$$

with $(\mathbf{a}, \mathbf{b}, c) = (\mathbf{a}, \mathbf{b}, c)[\Phi, \mathbf{h}, n]$. In particular, if $(\mathbf{a}, \mathbf{b}) = 0$ identically on (τ_0, τ_1) , then c vanishes identically on (τ_0, τ_1) if and only if c vanishes at one point of (τ_0, τ_1) .

Proof. Replace all occurrences of $\mathbf{a}, \mathbf{b}$ and c on the right hand side of (3.4) by the respective right hand sides of (3.2). Then, verify that everything cancels. $\square$

Definition 3.3. *For all $\mathbf{h} > 0$ and all vectors $\Phi = \alpha \oplus \beta \in \mathbb{R}^3 \oplus \mathbb{R}^3$ with $\beta_1, \beta_2, \beta_3 \neq 0$, define $A_{\mathbf{m}}[\Phi] \in (0, \infty)$ and $\varphi_{\mathbf{m}}[\Phi] \in \mathbb{R}$ by*

$$A_{\mathbf{m}}[\Phi] = \sqrt{|\alpha_{\mathbf{m}}|^2 + |\beta_{\mathbf{m}}|^2} > |\alpha_{\mathbf{m}}| \geq 0 \\ \varphi_{\mathbf{m}}[\Phi] = -\operatorname{arcsinh} \frac{\alpha_{\mathbf{m}}}{|\beta_{\mathbf{m}}|}$$

for all $\mathbf{m} = 1, 2, 3$. Equivalently,

$$\alpha_{\mathbf{m}} = -A_{\mathbf{m}}[\Phi] \tanh \varphi_{\mathbf{m}}[\Phi] \quad (3.5a)$$

$$\beta_{\mathbf{m}} = (\operatorname{sgn} \beta_{\mathbf{m}}) A_{\mathbf{m}}[\Phi] \operatorname{sech} \varphi_{\mathbf{m}}[\Phi] \quad (3.5b)$$

Furthermore, define $\xi_{\mathbf{m}}[\Phi, \mathbf{h}] \in \mathbb{R}$ by

$$\xi_{\mathbf{m}}[\Phi, \mathbf{h}] = \mathbf{h} \log \left| \frac{1}{2} \beta_{\mathbf{m}} \right|$$

for all $\mathbf{m} = 1, 2, 3$. Furthermore, for all $\mathbf{m}, \mathbf{n} = 1, 2, 3$, introduce the abbreviations

$$\alpha_{\mathbf{m}, \mathbf{n}}[\Phi] = \alpha_{\mathbf{m}} + \alpha_{\mathbf{n}} \quad \xi_{\mathbf{m}, \mathbf{n}}[\Phi, \mathbf{h}] = \xi_{\mathbf{m}}[\Phi, \mathbf{h}] + \xi_{\mathbf{n}}[\Phi, \mathbf{h}]$$

If no confusion can arise, we drop the explicit dependence $[\Phi]$ or $[\Phi, \mathbf{h}]$. For instance, we write $A_{\mathbf{m}} = A_{\mathbf{m}}[\Phi]$. If Φ is not an element of $\mathbb{R}^3 \oplus \mathbb{R}^3$, but rather a function of the real variable τ with values in $\mathbb{R}^3 \oplus \mathbb{R}^3$, with $\beta_1, \beta_2, \beta_3 \neq 0$ everywhere, then $A_{\mathbf{m}}, \varphi_{\mathbf{m}}, \xi_{\mathbf{m}}, \xi_{\mathbf{m}, \mathbf{n}}, \alpha_{\mathbf{m}, \mathbf{n}}$, with $\mathbf{m}, \mathbf{n} = 1, 2, 3$, are functions of τ , too. In this case, we define the additional functions $\theta_{\mathbf{m}}[\Phi, \mathbf{h}]$, $\mathbf{m} = 1, 2, 3$, through

$$\varphi_{\mathbf{m}}[\Phi](\tau) = \frac{1}{\mathbf{h}} (\tau - \theta_{\mathbf{m}}[\Phi, \mathbf{h}](\tau)) A_{\mathbf{m}}[\Phi](\tau)$$

Remark 3.2. In the context of Definition 3.3, we have, for all $\mathbf{m} = 1, 2, 3$:

$$\mathbf{h} |\varphi_{\mathbf{m}}| = -\xi_{\mathbf{m}} + \mathbf{h} \log \left(\left| \frac{1}{2} \alpha_{\mathbf{m}} \right| + \sqrt{\left| \frac{1}{2} \alpha_{\mathbf{m}} \right|^2 + \exp(\frac{1}{\mathbf{h}} 2\xi_{\mathbf{m}})} \right)$$

Lemma 3.1. Recall Definitions 3.1, 3.2, 3.3. For all $\mathbf{h} > 0$ and all $\Phi = \alpha \oplus \beta \in C^\infty((\tau_0, \tau_1), \mathbb{R}^6)$ such that $\beta_1, \beta_2, \beta_3$ never vanish on (τ_0, τ_1) , we have

$$\frac{d}{d\tau} \begin{pmatrix} A_i \\ \theta_i \end{pmatrix} = \frac{1}{(A_i)^2} \begin{pmatrix} \frac{1}{\mathbf{h}}(A_i)^2 \tanh \varphi_i & \frac{1}{\mathbf{h}}(A_i)^2 \operatorname{sech} \varphi_i \\ \varphi_i \tanh \varphi_i - 1 & \sinh \varphi_i + \varphi_i \operatorname{sech} \varphi_i \end{pmatrix} \begin{pmatrix} \mathbf{a}_i[\Phi, \mathbf{h}, B_i] \\ -\sigma_i \mathbf{b}_i[\Phi, \mathbf{h}, B_i] \end{pmatrix}$$

for $i = 1, 2, 3$ and $\sigma_i = \operatorname{sgn} \beta_i \in \{-1, +1\}$. The matrix on the right hand side has determinant $\frac{1}{\mathbf{h}}(A_i)^2 \cosh \varphi_i \neq 0$.

Proof. We have $\mathbf{a}_i[\Phi, \mathbf{h}, B_i] = -\mathbf{h} \frac{d}{d\tau} \alpha_i - (\beta_i)^2$ and $\mathbf{b}_i[\Phi, \mathbf{h}, B_i] = -\mathbf{h} \frac{d}{d\tau} \beta_i + \alpha_i \beta_i$. Replace all occurrences of α_i and β_i by the right hand sides of (3.5), respectively. Use $\frac{d}{d\tau} \varphi_i = \frac{1}{A_i} (\frac{d}{d\tau} A_i) \varphi_i + \frac{1}{\mathbf{h}} A_i (1 - \frac{d}{d\tau} \theta_i)$. Now, solve for $\frac{d}{d\tau} A_i$ and $\frac{d}{d\tau} \theta_i$. $\square$

Remark 3.3. So far, we have stated all definitions and propositions for a C^∞ -field $\Phi = \alpha \oplus \beta$, defined on an open interval. This was just for convenience. We will, from now on, use these definitions and propositions even when the C^∞ -requirement is not met, or when the field is defined on, say, a closed interval rather than an open interval. It will be clear in each case, that the respective definition or proposition still makes sense.

Definition 3.4. Set $S_3 = \{(1, 2, 3), (2, 3, 1), (3, 1, 2), (3, 2, 1), (1, 3, 2), (2, 1, 3)\}$, the set of all permutations of $(1, 2, 3)$.

Definition 3.5. For all $\sigma_* \in \{-1, +1\}^3$ let $\mathcal{D}(\sigma_*)$ be the set of all $\Phi = \alpha \oplus \beta \in \mathbb{R}^3 \oplus \mathbb{R}^3$ with $(\operatorname{sgn} \beta_1, \operatorname{sgn} \beta_2, \operatorname{sgn} \beta_3) = \sigma_*$. For all $\tau_0, \tau_1 \in \mathbb{R}$ with $\tau_0 < \tau_1$ let $\mathcal{E}(\sigma_*; \tau_0, \tau_1)$ be the set of all continuous maps $\Phi : [\tau_0, \tau_1] \rightarrow \mathcal{D}(\sigma_*)$.

Definition 3.6. For all $\pi = (\mathbf{a}, \mathbf{b}, \mathbf{c}) \in S_3$ and $\mathbf{h} > 0$ and $\sigma_* \in \{-1, +1\}^3$ define two functions $\mathcal{D}(\sigma_*) \times \mathcal{D}(\sigma_*) \rightarrow [0, \infty)$ by

$$d_{\mathcal{D}(\sigma_*), (\pi, \mathbf{h})}(\Phi, \Psi) = \max \left\{ \begin{aligned} &|A_{\mathbf{a}}[\Phi] - A_{\mathbf{a}}[\Psi]|, \quad \left| \mathbf{h} \frac{\varphi_{\mathbf{a}}[\Phi]}{A_{\mathbf{a}}[\Phi]} - \mathbf{h} \frac{\varphi_{\mathbf{a}}[\Psi]}{A_{\mathbf{a}}[\Psi]} \right|, \\ &|\alpha_{\mathbf{b}, \mathbf{a}}[\Phi] - \alpha_{\mathbf{b}, \mathbf{a}}[\Psi]|, \quad |\xi_{\mathbf{b}, \mathbf{a}}[\Phi, \mathbf{h}] - \xi_{\mathbf{b}, \mathbf{a}}[\Psi, \mathbf{h}]|, \\ &|\alpha_{\mathbf{c}, \mathbf{a}}[\Phi] - \alpha_{\mathbf{c}, \mathbf{a}}[\Psi]|, \quad |\xi_{\mathbf{c}, \mathbf{a}}[\Phi, \mathbf{h}] - \xi_{\mathbf{c}, \mathbf{a}}[\Psi, \mathbf{h}]| \end{aligned} \right\}$$

and

$$\mathcal{d}_{\mathcal{D}(\sigma_*), \mathbf{h}}(\Phi, \Psi) = \max_{i=1,2,3} \left\{ |\alpha_i[\Phi] - \alpha_i[\Psi]|, |\xi_i[\Phi, \mathbf{h}] - \xi_i[\Psi, \mathbf{h}]| \right\}$$

Then $(\mathcal{D}(\sigma_*), d_{\mathcal{D}(\sigma_*), (\pi, \mathbf{h})})$ and $(\mathcal{D}(\sigma_*), \mathcal{d}_{\mathcal{D}(\sigma_*), \mathbf{h}})$ are metric spaces.

Definition 3.7. For all $\pi \in S_3$ and $\mathbf{h} > 0$ and $\sigma_* \in \{-1, +1\}^3$ and $\tau_0, \tau_1 \in \mathbb{R}$ with $\tau_0 < \tau_1$ define a function $\mathcal{E}(\sigma_*; \tau_0, \tau_1) \times \mathcal{E}(\sigma_*; \tau_0, \tau_1) \rightarrow [0, \infty)$ by

$$d_{\mathcal{E}(\sigma_*; \tau_0, \tau_1), (\pi, \mathbf{h})}(\Phi, \Psi) = \sup_{\tau \in [\tau_0, \tau_1]} d_{\mathcal{D}(\sigma_*), (\pi, \mathbf{h})}(\Phi(\tau), \Psi(\tau))$$

Then $(\mathcal{E}(\sigma_*; \tau_0, \tau_1), d_{\mathcal{E}(\sigma_*; \tau_0, \tau_1), (\pi, \mathbf{h})})$ is a metric space.

Lemma 3.2. Let $\pi = (\mathbf{a}, \mathbf{b}, \mathbf{c}) \in S_3$ and $\mathbf{h} > 0$ and $\sigma_* \in \{-1, +1\}^3$. Suppose $\mathbf{h} \leq 1$. Let $C, D \geq 1$ be constants. Then, for all $\Phi, \Psi \in \mathcal{D}(\sigma_*)$ such that

$$C^{-1} \leq A_{\mathbf{a}}[X] \leq C \quad D^{-1} \leq \mathbf{h} |\varphi_{\mathbf{a}}[X]| \leq D$$

for both $X = \Phi$ and $X = \Psi$ and such that $\operatorname{sgn} \varphi_{\mathbf{a}}[\Phi] = \operatorname{sgn} \varphi_{\mathbf{a}}[\Psi]$, we have:

- (a) $d_{\mathcal{D}}(\Phi, \Psi) \leq 2^3 C^2 D d_{\mathcal{D}}(\Phi, \Psi)$
(b) If $\exp(-\frac{1}{\mathbf{h}} C^{-2} D^{-1}) \leq 2^{-6} C^{-4} D^{-2}$, then $d_{\mathcal{D}}(\Phi, \Psi) \leq 2^5 C^3 D d_{\mathcal{D}}(\Phi, \Psi)$

Here, $d_{\mathcal{D}} = d_{\mathcal{D}(\sigma_*)}(\pi, \mathbf{h})$ and $d_{\mathcal{D}} = d_{\mathcal{D}(\sigma_*)}(\pi, \mathbf{h})$.

Proof. In this proof, A, B, α, ξ play the roles of $A_{\mathbf{a}}, \mathbf{h}\varphi_{\mathbf{a}}/A_{\mathbf{a}}, \alpha_{\mathbf{a}}, \xi_{\mathbf{a}}$, respectively. To show (a), let $P : (0, \infty) \times \mathbb{R} \rightarrow \mathbb{R}^2$, $(A, B) \mapsto (\alpha(A, B), \xi(A, B))$, where

$$\alpha(A, B) = -A \tanh(\frac{1}{\mathbf{h}} AB) \quad \xi(A, B) = \mathbf{h} \log(\frac{1}{2} A \operatorname{sech}(\frac{1}{\mathbf{h}} AB))$$

This is a diffeomorphism. The Jacobian J of P is given by

$$J = \begin{pmatrix} \frac{\partial \alpha}{\partial A} & \frac{\partial \alpha}{\partial B} \\ \frac{\partial \xi}{\partial A} & \frac{\partial \xi}{\partial B} \end{pmatrix} = \begin{pmatrix} -\frac{1}{\mathbf{h}} AB \operatorname{sech}^2(\frac{1}{\mathbf{h}} AB) - \tanh(\frac{1}{\mathbf{h}} AB) & -\frac{1}{\mathbf{h}} A^2 \operatorname{sech}^2(\frac{1}{\mathbf{h}} AB) \\ \frac{\mathbf{h}}{A} - B \tanh(\frac{1}{\mathbf{h}} AB) & -A \tanh(\frac{1}{\mathbf{h}} AB) \end{pmatrix}$$

Let $p_i = (A_i, B_i) \in (0, \infty) \times \mathbb{R}$ and set $(\alpha_i, \xi_i) = P(p_i)$, where $i = 0, 1$. Set $\gamma(t) = (A(t), B(t)) = (1-t)p_0 + tp_1$ where $t \in [0, 1]$. We have

$$\begin{pmatrix} \alpha_1 - \alpha_0 \\ \xi_1 - \xi_0 \end{pmatrix} = M \begin{pmatrix} A_1 - A_0 \\ B_1 - B_0 \end{pmatrix} \quad \text{with} \quad M = \begin{pmatrix} M_{00} & M_{01} \\ M_{10} & M_{11} \end{pmatrix} = \int_0^1 dt J(\gamma(t))$$

Suppose $C^{-1} \leq A_i \leq C$ and $(CD)^{-1} \leq |B_i| \leq CD$ and $\operatorname{sgn} B_0 = \operatorname{sgn} B_1$. Then, $C^{-1} \leq A(t) \leq C$ and $(CD)^{-1} \leq |B(t)| \leq CD$ for all $t \in [0, 1]$. Observe that $|\varphi \operatorname{sech}^2 \varphi| \leq \frac{1}{2}$ for all $\varphi \in \mathbb{R}$. We have $|M_{ij}| \leq 2C^2 D$ for all $i, j \in \{0, 1\}$. This implies (a).

We show that under the assumptions of (b), we have $|\det M| \geq 2^{-3} C^{-1}$, and therefore $|(M^{-1})_{ij}| \leq 2^4 C^3 D$ for all $i, j \in \{0, 1\}$. This would imply (b). We have $|\det M| \geq |M_{00}M_{11}| - |M_{01}M_{10}|$. Set $\varphi(t) = \frac{1}{\mathbf{h}} A(t)B(t)$. We have $|\varphi(t)| \geq \frac{1}{\mathbf{h}} C^{-2} D^{-1}$. By the assumption of (b), we have $e^{-|\varphi(t)|} \leq 2^{-6} C^{-4} D^{-2}$, for all $t \in [0, 1]$. We will also use the general inequalities $0 \leq 1 - \tanh |\varphi| \leq 2e^{-2|\varphi|}$ and $|\varphi \operatorname{sech}^2 \varphi| \leq 4|\varphi|e^{-2|\varphi|} \leq 4e^{-|\varphi|}$. We have $|\varphi \operatorname{sech}^2 \varphi - \tanh \varphi| \geq \tanh |\varphi| = 1 - (1 - \tanh |\varphi|) \geq 2^{-1}$. The last inequality holds for all $t \in [0, 1]$ and implies $|M_{00}| \geq 2^{-1}$, because φ has constant sign. We have $|M_{11}| \geq 2^{-1} C^{-1}$ and $|M_{10}| \leq 2CD$ and $|M_{01}| \leq 2^{-4} C^{-2} D^{-1}$. This implies $|\det M| \geq 2^{-3} C^{-1}$. $\square$

Definition 3.8. Let $\mathcal{X} = \mathcal{D}(\sigma_*)$ or $\mathcal{X} = \mathcal{E}(\sigma_*; \tau_0, \tau_1)$. For all $\delta \geq 0$ and $\Phi \in \mathcal{X}$ and $\pi \in S_3$ and $\mathbf{h} > 0$, set $B_{\mathcal{X}, (\pi, \mathbf{h})}[\delta, \Phi] = \{\Psi \in \mathcal{X} \mid d_{\mathcal{X}, (\pi, \mathbf{h})}(\Phi, \Psi) \leq \delta\}$.

Definition 3.9 (The reference field Φ_0). For all $\pi = (\mathbf{a}, \mathbf{b}, \mathbf{c}) \in S_3$, $\mathbf{f} = (\mathbf{h}, w, q) \in (0, \infty)^3$, $\sigma_* \in \{-1, +1\}^3$ let $\Phi_0 = \Phi_0(\pi, \mathbf{f}, \sigma_*) : \mathbb{R} \rightarrow \mathcal{D}(\sigma_*)$ be given by

$$A_{\mathbf{a}}[\Phi_0](\tau) = 1 \tag{3.6a}$$

$$\theta_{\mathbf{a}}[\Phi_0, \mathbf{h}](\tau) = 0 \tag{3.6b}$$

$$\alpha_{\mathbf{b}, \mathbf{a}}[\Phi_0](\tau) = -(1+w)^{-1} \tag{3.6c}$$

$$\alpha_{\mathbf{c}, \mathbf{a}}[\Phi_0](\tau) = -(1+w) \tag{3.6d}$$

$$\xi_{\mathbf{b}, \mathbf{a}}[\Phi_0, \mathbf{h}](\tau) = -1 - \mathbf{h} \log 2 - (1+w)^{-1} \tau \tag{3.6e}$$

$$\xi_{\mathbf{c}, \mathbf{a}}[\Phi_0, \mathbf{h}](\tau) = -(1+w)q - \mathbf{h} \log 2 - (1+w) \tau \tag{3.6f}$$

(see Definition 3.3) for all $\tau \in \mathbb{R}$.

Remark 3.4. The field Φ_0 is, up to renaming, given by equation (3.12) in [BKL1].

Lemma 3.3. *Let Φ_0 be as in Definition 3.9. Then $(\mathbf{a}, \mathbf{b}, c)[\Phi_0, \mathbf{h}, B_{\mathbf{a}}] = 0$ on $\mathbb{R}$.*

Proof. Let $\alpha = \alpha[\Phi_0]$, $\beta = \beta[\Phi_0]$, $\xi = \xi[\Phi_0, \mathbf{h}]$. We have $(\mathbf{a}_{\mathbf{a}}, \mathbf{b}_{\mathbf{a}})[\Phi_0, \mathbf{h}, B_{\mathbf{a}}] = 0$ by Lemma 3.1. For $\mathbf{p} \in \{\mathbf{b}, \mathbf{c}\}$, we have $\mathbf{a}_{\mathbf{a}}[\Phi_0, \mathbf{h}, B_{\mathbf{a}}] + \mathbf{a}_{\mathbf{p}}[\Phi_0, \mathbf{h}, B_{\mathbf{a}}] = -\mathbf{h} \frac{d}{d\tau} \alpha_{\mathbf{a}, \mathbf{p}} = 0$, that is $\mathbf{a}_{\mathbf{p}}[\Phi_0, \mathbf{h}, B_{\mathbf{a}}] = 0$. We also have $\beta_{\mathbf{a}}^{-1} \mathbf{b}_{\mathbf{a}}[\Phi_0, \mathbf{h}, B_{\mathbf{a}}] + \beta_{\mathbf{p}}^{-1} \mathbf{b}_{\mathbf{p}}[\Phi_0, \mathbf{h}, B_{\mathbf{a}}] = -\frac{d}{d\tau} \xi_{\mathbf{a}, \mathbf{p}} + \alpha_{\mathbf{a}, \mathbf{p}} = 0$, that is $\mathbf{b}_{\mathbf{p}}[\Phi_0, \mathbf{h}, B_{\mathbf{a}}] = 0$. Finally, $c[\Phi_0, \mathbf{h}, B_{\mathbf{a}}] = -\alpha_{\mathbf{a}}^2 - \beta_{\mathbf{a}}^2 + \alpha_{\mathbf{a}, \mathbf{b}} \alpha_{\mathbf{a}, \mathbf{c}} = -A_{\mathbf{a}}^2 + \alpha_{\mathbf{a}, \mathbf{b}} \alpha_{\mathbf{a}, \mathbf{c}} = 0$. Here, $A_{\mathbf{a}} = A_{\mathbf{a}}[\Phi_0]$. $\square$

Definition 3.10. *For all $\mathbf{f} = (\mathbf{h}, w, q) \in (0, \infty)^3$ set*

$$\begin{aligned} \tau_-(\mathbf{f}) &= -\left(1 - \frac{1}{2+w}\right) \min\{1, q\} < 0 \\ \tau_+(\mathbf{f}) &= 1 + \frac{1}{w} > 0 \end{aligned}$$

Lemma 3.4 (Technical Lemma 1). *Let $\pi = (\mathbf{a}, \mathbf{b}, \mathbf{c}) \in S_3$, $\mathbf{f} = (\mathbf{h}, w, q) \in (0, \infty)^3$, $\sigma_* \in \{-1, +1\}^3$ and fix $\delta > 0$, $\epsilon_- \in (0, -\tau_-)$, $\epsilon_+ \in (0, \tau_+)$ where $\tau_{\pm} = \tau_{\pm}(\mathbf{f})$. Set*

$$\tau_{0-} = \tau_- + \epsilon_- < 0 \quad \Phi_0 = \Phi_0(\pi, \mathbf{f}, \sigma_*)|_{[\tau_{0-}, \tau_{0+}]} \quad (3.7a)$$

$$\tau_{0+} = \tau_+ - \epsilon_+ > 0 \quad \mathcal{E} = \mathcal{E}(\sigma_*; \tau_{0-}, \tau_{0+}) \quad (3.7b)$$

Then $\Phi_0 \in \mathcal{E}$. Furthermore, if the inequality

$$\delta \leq 2^{-4} \min\left\{1, w, \epsilon_-, \frac{\epsilon_+}{\tau_+ \tau_{0+}}\right\} \quad (3.8)$$

holds, then for all $\Phi = \alpha \oplus \beta \in B_{\mathcal{E}, (\pi, \mathbf{h})}[\delta, \Phi_0]$ the estimates

$$\begin{aligned} \max\left\{|\beta_{\mathbf{b}}|^2, |\beta_{\mathbf{c}}|^2, |\beta_{\mathbf{b}}\beta_{\mathbf{a}}|, |\beta_{\mathbf{c}}\beta_{\mathbf{a}}|\right\} &\leq 2^4 \exp\left(-\frac{1}{4\mathbf{h}} \min\{1, \epsilon_-, \frac{\epsilon_+}{\tau_+}\}\right) \\ |A_{\mathbf{a}}[\Phi] - 1| &\leq 2^{-1} \\ |\varphi_{\mathbf{a}}[\Phi]| &\leq \frac{1}{\mathbf{h}} 2(1 + |\tau|) \\ |\beta_{\mathbf{a}}| &\leq 2 \end{aligned}$$

hold on $[\tau_{0-}, \tau_{0+}]$.

Proof. The following estimates hold for the components of Φ , for all $\tau \in [\tau_{0-}, \tau_{0+}]$:

$$\begin{aligned} |\beta_{\mathbf{b}}\beta_{\mathbf{a}}| &= 4 \exp\left(\frac{1}{\mathbf{h}} \xi_{\mathbf{b}, \mathbf{a}}\right) \\ &\leq 4 \exp\left(\frac{1}{\mathbf{h}} \xi_{\mathbf{b}, \mathbf{a}}[\Phi_0, \mathbf{h}] + \frac{1}{\mathbf{h}} \delta\right) \\ &\leq 2 \exp\left(-\frac{1}{\mathbf{h}} - \frac{1}{\mathbf{h}}(1+w)^{-1}\tau + \frac{1}{\mathbf{h}} \delta\right) \\ &\leq 2 \exp\left(-\frac{1}{\mathbf{h}} - \frac{1}{\mathbf{h}}(1+w)^{-1}\tau_- + \frac{1}{\mathbf{h}} \delta\right) \\ &\leq 2 \exp\left(-\frac{1}{\mathbf{h}} + \frac{1}{\mathbf{h}}(2+w)^{-1} + \frac{1}{\mathbf{h}} \delta\right) \\ &\leq 2 \exp\left(-\frac{1}{4\mathbf{h}}\right) \\ |\beta_{\mathbf{c}}\beta_{\mathbf{a}}| &= 4 \exp\left(\frac{1}{\mathbf{h}} \xi_{\mathbf{c}, \mathbf{a}}\right) \\ &\leq 4 \exp\left(\frac{1}{\mathbf{h}} \xi_{\mathbf{c}, \mathbf{a}}[\Phi_0, \mathbf{h}] + \frac{1}{\mathbf{h}} \delta\right) \\ &\leq 2 \exp\left(-\frac{1}{\mathbf{h}}(1+w)q - \frac{1}{\mathbf{h}}(1+w)\tau + \frac{1}{\mathbf{h}} \delta\right) \\ &\leq 2 \exp\left(-\frac{1}{\mathbf{h}}(1+w)q - \frac{1}{\mathbf{h}}(1+w)(\tau_- + \epsilon_-) + \frac{1}{\mathbf{h}} \delta\right) \\ &\leq 2 \exp\left(-\frac{1}{\mathbf{h}}(1+w)q + \frac{1}{\mathbf{h}} \frac{(1+w)^2}{2+w} q - \frac{1}{\mathbf{h}} \epsilon_- + \frac{1}{\mathbf{h}} \delta\right) \\ &\leq 2 \exp\left(-\frac{1}{2\mathbf{h}} \epsilon_-\right) \end{aligned}$$

$$\begin{aligned}
|\varphi_{\mathbf{a}}| &= \frac{1}{\mathbf{h}} A_{\mathbf{a}} |\tau - \theta_{\mathbf{a}}| \\
&\leq \frac{1}{\mathbf{h}} (1 + \delta) (|\tau| + \delta) \\
&\leq \frac{1}{\mathbf{h}} (|\tau| + \delta |\tau| + 2\delta) \\
&\leq \frac{1}{\mathbf{h}} 2(1 + |\tau|) \\
|\beta_{\mathbf{a}}|^{-1} &= |A_{\mathbf{a}}|^{-1} \cosh \varphi_{\mathbf{a}} \\
&\leq 2 \exp(|\varphi_{\mathbf{a}}|) \\
&\leq 2 \exp\left(\frac{1}{\mathbf{h}} |\tau| + \frac{1}{\mathbf{h}} \delta |\tau| + \frac{1}{\mathbf{h}} 2\delta\right) \\
|\beta_{\mathbf{b}}| &= |\beta_{\mathbf{b}} \beta_{\mathbf{a}}| \cdot |\beta_{\mathbf{a}}|^{-1} \\
&\leq 4 \exp\left(-\frac{1}{\mathbf{h}} - \frac{1}{\mathbf{h}} (1 + w)^{-1} \tau + \frac{1}{\mathbf{h}} |\tau| + \frac{1}{\mathbf{h}} \delta |\tau| + \frac{1}{\mathbf{h}} 3\delta\right) \\
&\leq 4 \exp\left(\frac{1}{\mathbf{h}} \max\left\{-1 - \frac{2+w}{1+w} \tau_{0-} - \delta \tau_{0-}, -1 + \frac{w}{1+w} \tau_{0+} + \delta \tau_{0+}\right\} + \frac{1}{\mathbf{h}} 3\delta\right) \\
&\leq 4 \exp\left(\frac{1}{\mathbf{h}} \max\{-\epsilon_- - \delta \tau_{0-}, -\frac{\epsilon_+}{\tau_+} + \delta \tau_{0+}\} + \frac{1}{\mathbf{h}} 3\delta\right) \\
&\leq 4 \exp\left(-\frac{1}{\mathbf{h}} \frac{15}{16} \min\{\epsilon_-, \frac{\epsilon_+}{\tau_+}\} + \frac{1}{\mathbf{h}} 3\delta\right) \\
&\leq 4 \exp\left(-\frac{1}{2\mathbf{h}} \min\{\epsilon_-, \frac{\epsilon_+}{\tau_+}\}\right)
\end{aligned}$$

The last step uses $\delta \leq 2^{-3} \frac{\epsilon_+}{\tau_+}$. In the case $\epsilon_+ \geq \frac{1}{2} \tau_+$, this follows from $\delta \leq 2^{-4}$. If $\epsilon_+ \leq \frac{1}{2} \tau_+$, then this follows from $\delta \leq 2^{-4} \frac{\epsilon_+}{\tau_+ \tau_{0+}}$, because $\tau_{0+} = \tau_+ - \epsilon_+ \geq \frac{1}{2} \tau_+ \geq \frac{1}{2}$.

$$\begin{aligned}
|\beta_{\mathbf{c}}| &= |\beta_{\mathbf{c}} \beta_{\mathbf{a}}| \cdot |\beta_{\mathbf{a}}|^{-1} \\
&\leq 4 \exp\left(-\frac{1}{\mathbf{h}} (1 + w) q - \frac{1}{\mathbf{h}} (1 + w) \tau + \frac{1}{\mathbf{h}} |\tau| + \frac{1}{\mathbf{h}} \delta |\tau| + \frac{1}{\mathbf{h}} 3\delta\right) \\
&\leq 4 \exp\left(-\frac{1}{\mathbf{h}} (1 + w) q \right. \\
&\quad \left. + \frac{1}{\mathbf{h}} \max\left\{-(2 + w + \delta) \tau_{0-}, -(w - \delta) \tau_{0+}\right\} + \frac{1}{\mathbf{h}} 3\delta\right) \\
&\leq 4 \exp\left(-\frac{1}{\mathbf{h}} (1 + w) q + \frac{1}{\mathbf{h}} (2 + w) |\tau_-| - \frac{1}{\mathbf{h}} (2 + w + \delta) \epsilon_- + \frac{1}{\mathbf{h}} 4\delta\right) \\
&\leq 4 \exp\left(-\frac{1}{\mathbf{h}} 2\epsilon_- + \frac{1}{\mathbf{h}} 4\delta\right) \\
&\leq 4 \exp\left(-\frac{1}{\mathbf{h}} \epsilon_-\right)
\end{aligned}$$

This concludes the proof. $\square$

Lemma 3.5. *Recall Definitions 3.1 and 3.2. We have*

$$\begin{aligned}
\mathbf{a}_{\mathbf{a}}[\Phi, \mathbf{h}, Z, B_{\mathbf{a}}] &= +\beta_{\mathbf{b}}^2 + \beta_{\mathbf{c}}^2 - 2\beta_{\mathbf{b}}\beta_{\mathbf{c}} \\
\mathbf{a}_{\mathbf{b}}[\Phi, \mathbf{h}, Z, B_{\mathbf{a}}] &= -\beta_{\mathbf{b}}^2 + \beta_{\mathbf{c}}^2 - 2\beta_{\mathbf{a}}\beta_{\mathbf{c}} \\
\mathbf{a}_{\mathbf{c}}[\Phi, \mathbf{h}, Z, B_{\mathbf{a}}] &= +\beta_{\mathbf{b}}^2 - \beta_{\mathbf{c}}^2 - 2\beta_{\mathbf{a}}\beta_{\mathbf{b}}
\end{aligned}$$

for all $(\mathbf{a}, \mathbf{b}, \mathbf{c}) \in S_3$.

Remark 3.5. Lemma 3.5 displays the differences between the equations $\mathbf{a}[\Phi, \mathbf{h}, Z] = 0$ and $\mathbf{a}[\Phi, \mathbf{h}, B_{\mathbf{a}}] = 0$. Lemma 3.4 gives bounds for the terms that appear in these differences. Informally, they tend exponentially to zero as $\mathbf{h} \downarrow 0$. This quantifies a basic guiding intuition of [BKL1].

Definition 3.11. For all vectors $\Phi = \alpha \oplus \beta \in \mathbb{R}^3 \oplus \mathbb{R}^3$ with $\beta_1, \beta_2, \beta_3 \neq 0$, all $\pi = (\mathbf{a}, \mathbf{b}, \mathbf{c}) \in S_3$ and all $\mathbf{h} > 0$, define four real numbers by

$$\mathbf{I}_1[\Phi, \mathbf{h}, \pi] = -\frac{1}{\mathbf{h}} \mathbf{a}_{\mathbf{a}}[\Phi, \mathbf{h}, Z, B_{\mathbf{a}}] \tanh \varphi_{\mathbf{a}}[\Phi]$$

$$\mathbf{I}_2[\Phi, \mathbf{h}, \pi] = (A_{\mathbf{a}}[\Phi])^{-2} \mathbf{a}_{\mathbf{a}}[\Phi, \mathbf{h}, Z, B_{\mathbf{a}}] \left(1 - \varphi_{\mathbf{a}}[\Phi] \tanh \varphi_{\mathbf{a}}[\Phi]\right)$$

$$\mathbf{I}_{(3, \mathbf{p})}[\Phi, \mathbf{h}, \pi] = \frac{1}{\mathbf{h}} \mathbf{a}_{\mathbf{p}}[\Phi, \mathbf{h}, Z, B_{\mathbf{a}}] + \frac{1}{\mathbf{h}} \mathbf{a}_{\mathbf{a}}[\Phi, \mathbf{h}, Z, B_{\mathbf{a}}]$$

where $\mathbf{p} \in \{\mathbf{b}, \mathbf{c}\}$. If Φ is not an element of $\mathbb{R}^3 \oplus \mathbb{R}^3$, but rather a function with values in $\mathbb{R}^3 \oplus \mathbb{R}^3$, with $\beta_1, \beta_2, \beta_3 \neq 0$ everywhere, then $\mathbf{I}_1, \mathbf{I}_2, \mathbf{I}_{(3, \mathbf{b})}, \mathbf{I}_{(3, \mathbf{c})}$ are functions, too.

Lemma 3.6 (Technical Lemma 2). In the context of Lemma 3.4, if $\delta > 0$ satisfies (3.8), then, for all $\Phi, \Psi \in B_{\mathcal{E}, (\pi, \mathbf{h})}[\delta, \Phi_0]$ and all $S \in \{1, 2, (3, \mathbf{b}), (3, \mathbf{c})\}$, the estimates

$$|\mathbf{I}_S[\Phi]| \leq 2^{11} \max\{1, \frac{1}{\mathbf{h}}, \frac{1}{\mathbf{h}}|\tau|\} \exp\left(-\frac{1}{4\mathbf{h}} \min\{1, \epsilon_-, \frac{\epsilon_+}{\tau_+}\}\right) \quad (3.9a)$$

$$|\mathbf{I}_S[\Phi] - \mathbf{I}_S[\Psi]| \leq 2^{17} \left(\max\{1, \frac{1}{\mathbf{h}}, \frac{1}{\mathbf{h}}|\tau|\}\right)^2 \exp\left(-\frac{1}{4\mathbf{h}} \min\{1, \epsilon_-, \frac{\epsilon_+}{\tau_+}\}\right) d_{\mathcal{E}}(\Phi, \Psi) \quad (3.9b)$$

hold on $[\tau_{0-}, \tau_{0+}]$. Here, $\mathbf{I}_S[\Phi] = \mathbf{I}_S[\Phi, \mathbf{h}, \pi]$, $\mathbf{I}_S[\Psi] = \mathbf{I}_S[\Psi, \mathbf{h}, \pi]$ and $d_{\mathcal{E}} = d_{\mathcal{E}, (\pi, \mathbf{h})}$.

Proof. In this proof, we simplify the notation by suppressing $\mathbf{h} > 0$ and abbreviating

$$M = \exp\left(-\frac{1}{4\mathbf{h}} \min\{1, \epsilon_-, \frac{\epsilon_+}{\tau_+}\}\right) \quad M_1 = \max\{1, \frac{1}{\mathbf{h}}, \frac{1}{\mathbf{h}}|\tau|\}$$

Lemmas 3.4, 3.5 imply $|\mathbf{a}_{\mathbf{i}}[\Phi, \mathbf{h}, Z, B_{\mathbf{a}}]| \leq 2^6 M$, $\mathbf{i} = 1, 2, 3$, and $|\varphi_{\mathbf{a}}[\Phi]| \leq 2^2 M_1$ and $(A_{\mathbf{a}}[\Phi])^{-2} \leq 2^2$. This implies (3.9a). To show (3.9b), observe that (here $\mathbf{p}, \mathbf{q} \in \{\mathbf{b}, \mathbf{c}\}$)

$$\begin{aligned} |\varphi_{\mathbf{a}}[\Phi] - \varphi_{\mathbf{a}}[\Psi]| &\leq \frac{1}{\mathbf{h}} |A_{\mathbf{a}}[\Phi] - A_{\mathbf{a}}[\Psi]| |\tau| + \frac{1}{\mathbf{h}} |A_{\mathbf{a}}[\Phi] \theta_{\mathbf{a}}[\Phi] - A_{\mathbf{a}}[\Psi] \theta_{\mathbf{a}}[\Psi]| \\ &\leq \frac{1}{\mathbf{h}} (1 + |\tau|) |A_{\mathbf{a}}[\Phi] - A_{\mathbf{a}}[\Psi]| + \frac{1}{\mathbf{h}} 2 |\theta_{\mathbf{a}}[\Phi] - \theta_{\mathbf{a}}[\Psi]| \\ &\leq 2^2 M_1 d_{\mathcal{E}}(\Phi, \Psi) \\ |\xi_{\mathbf{a}}[\Phi] - \xi_{\mathbf{a}}[\Psi]| &\leq \mathbf{h} |\log A_{\mathbf{a}}[\Phi] - \log A_{\mathbf{a}}[\Psi]| \\ &\quad + \mathbf{h} |\log \cosh \varphi_{\mathbf{a}}[\Phi] - \log \cosh \varphi_{\mathbf{a}}[\Psi]| \\ &\leq 2^3 \mathbf{h} M_1 d_{\mathcal{E}}(\Phi, \Psi) \\ |\beta_{\mathbf{p}}[\Phi] \beta_{\mathbf{a}}[\Phi] - \beta_{\mathbf{p}}[\Psi] \beta_{\mathbf{a}}[\Psi]| &\leq \frac{1}{\mathbf{h}} \max\{|\beta_{\mathbf{p}}[\Phi] \beta_{\mathbf{a}}[\Phi]|, |\beta_{\mathbf{p}}[\Psi] \beta_{\mathbf{a}}[\Psi]|\} \\ &\quad \times |\xi_{\mathbf{a}, \mathbf{p}}[\Phi] - \xi_{\mathbf{a}, \mathbf{p}}[\Psi]| \\ &\leq 2^4 M_1 M d_{\mathcal{E}}(\Phi, \Psi) \\ |\beta_{\mathbf{p}}[\Phi] - \beta_{\mathbf{p}}[\Psi]| &\leq \frac{1}{\mathbf{h}} \max\{|\beta_{\mathbf{p}}[\Phi]|, |\beta_{\mathbf{p}}[\Psi]|\} |\xi_{\mathbf{p}}[\Phi] - \xi_{\mathbf{p}}[\Psi]| \\ &\leq \frac{1}{\mathbf{h}} 2^2 M^{1/2} (|\xi_{\mathbf{a}, \mathbf{p}}[\Phi] - \xi_{\mathbf{a}, \mathbf{p}}[\Psi]| + |\xi_{\mathbf{a}}[\Phi] - \xi_{\mathbf{a}}[\Psi]|) \\ &\leq 2^6 M_1 M^{1/2} d_{\mathcal{E}}(\Phi, \Psi) \\ |\beta_{\mathbf{p}}[\Phi] \beta_{\mathbf{q}}[\Phi] - \beta_{\mathbf{p}}[\Psi] \beta_{\mathbf{q}}[\Psi]| &\leq 2^9 M_1 M d_{\mathcal{E}}(\Phi, \Psi) \end{aligned}$$

Consequently, for $\mathbf{i} = 1, 2, 3$,

$$|\mathbf{a}_{\mathbf{i}}[\Phi, \mathbf{h}, Z, B_{\mathbf{a}}] - \mathbf{a}_{\mathbf{i}}[\Psi, \mathbf{h}, Z, B_{\mathbf{a}}]| \leq 2^{11} M_1 M d_{\mathcal{E}}(\Phi, \Psi)$$

With these estimates, (3.9b) follows. Observe that $\mathbb{R} \rightarrow \mathbb{R}, x \mapsto x \tanh x$ is Lipschitz with Lipschitz-constant $L > 0$ determined by $L \tanh L = 1$, in particular $L < 2$. $\square$

Definition 3.12. For all $\pi = (\mathbf{a}, \mathbf{b}, \mathbf{c}) \in S_3$ and $\mathbf{f} = (\mathbf{h}, w, q) \in (0, \infty)^2 \times \mathbb{R}$ (we don't require $q > 0$ here) and $\sigma_* \in \{-1, +1\}^3$, let $\Phi_* = \Phi_*(\pi, \mathbf{f}, \sigma_*) \in \mathcal{D}(\sigma_*)$ be given by

$$\begin{aligned} \alpha_{\mathbf{a}}[\Phi_*] &= -1 & \xi_{\mathbf{a}}[\Phi_*, \mathbf{h}] &= -\frac{1+w}{1+2w}(1 + \mathbf{h} \log 2) \\ \alpha_{\mathbf{b}}[\Phi_*] &= \frac{w}{1+w} & \xi_{\mathbf{b}}[\Phi_*, \mathbf{h}] &= -\frac{1+w}{1+2w}(1 + \mathbf{h} \log 2) \\ \alpha_{\mathbf{c}}[\Phi_*] &= -w - \mu & \xi_{\mathbf{c}}[\Phi_*, \mathbf{h}] &= -(1+w)q - \frac{w(1+w)}{1+2w} - \frac{1+3w+w^2}{1+2w} \mathbf{h} \log 2 \end{aligned}$$

and

$$\mu = (1+w)(\beta_1^2 + \beta_2^2 + \beta_3^2 - 2\beta_2\beta_3 - 2\beta_3\beta_1 - 2\beta_1\beta_2)|_{\beta=\beta[\Phi_*]} \quad (3.10)$$

Definition 3.13. For all $\pi = (\mathbf{a}, \mathbf{b}, \mathbf{c}) \in S_3$, $\sigma_* \in \{-1, +1\}^3$ let $\mathcal{H}(\pi, \sigma_*) \subset \mathcal{D}(\sigma_*)$ be the set of all vectors $\Phi = \alpha \oplus \beta \in \mathcal{D}(\sigma_*)$ with

$$|\beta_{\mathbf{a}}| = |\beta_{\mathbf{b}}| \quad \sum_{(i,j,k) \in \mathcal{C}} (\alpha_j \alpha_k - (\beta_i)^2 + 2\beta_j \beta_k) = 0 \quad (3.11a)$$

$$0 < \alpha_{\mathbf{b}} < -\alpha_{\mathbf{a}} \quad (\alpha_{\mathbf{b}} + |\alpha_{\mathbf{a}}|) \log |\beta_{\mathbf{a}}/\alpha_{\mathbf{a}}| < \alpha_{\mathbf{b}} \log 2 \quad (3.11b)$$

Lemma 3.7. Let $\pi = (\mathbf{a}, \mathbf{b}, \mathbf{c}) \in S_3$ and $\sigma_* \in \{-1, +1\}^3$. The set $\mathcal{H}(\pi, \sigma_*) \subset \mathcal{D}(\sigma_*)$ is a smooth 4-dimensional submanifold. The map

$$\begin{aligned} (0, \infty)^3 \times \mathbb{R} &\rightarrow \mathcal{H}(\pi, \sigma_*) \\ (\lambda, \mathbf{h}, w, q) &\mapsto \lambda \Phi_*(\pi, (\mathbf{h}, w, q), \sigma_*) \end{aligned} \quad (3.12)$$

is a diffeomorphism. Its inverse is given by

$$w = -\alpha_{\mathbf{b}}/(\alpha_{\mathbf{a}} + \alpha_{\mathbf{b}}) \quad \frac{1}{\mathbf{h}} = -\frac{1+2w}{1+w} \log |\beta_{\mathbf{a}}/\alpha_{\mathbf{a}}| + \frac{w}{1+w} \log 2 \quad (3.13a)$$

$$\lambda = -\alpha_{\mathbf{a}} \quad q = -\frac{1}{1+w} \mathbf{h} \log |\beta_{\mathbf{c}}/\alpha_{\mathbf{a}}| - \frac{w}{1+2w} (1 + \mathbf{h} \log 2) \quad (3.13b)$$

Proof. $\mathcal{H}(\pi, \sigma_*)$ is the graph of a smooth map from an open subset of $\mathbb{R}^4$ to $\mathbb{R}^2$. Namely the map given by solving (3.11a) for $(\alpha_{\mathbf{c}}, \beta_{\mathbf{b}})$ in terms of $(\alpha_{\mathbf{a}}, \alpha_{\mathbf{b}}, \beta_{\mathbf{a}}, \beta_{\mathbf{c}})$, whose domain is given by (3.11b) and appropriate sign conditions inherited from $\mathcal{D}(\sigma_*)$. The map (3.12) is well-defined, i.e. $\lambda \Phi_*(\pi, (\mathbf{h}, w, q), \sigma_*) \in \mathcal{H}(\pi, \sigma_*)$. The map (3.13) is well-defined, because the two right hand sides in (3.13a) and the first right hand side in (3.13b) are positive, by (3.11b). By direct calculation, the two maps are inverses. $\square$

Definition 3.14. For all $\mathbf{f} = (\mathbf{h}, w, q) \in (0, \infty)^3$ set

$$\begin{aligned} \tau_{1-}(\mathbf{f}) &= \begin{cases} -\frac{1+w}{3+w} q - \frac{1}{3+w} \mathbf{h} \log 2 & \text{if } q \leq 1 \\ -\frac{1+w}{3+2w} - \frac{1+w}{3+2w} \mathbf{h} \log 2 & \text{if } q > 1 \end{cases} < 0 \\ \tau_{1+}(\mathbf{f}) &= (1 + \mathbf{h} \log 2) \frac{1+w}{1+2w} > 0 \end{aligned}$$

Definition 3.15. For all $\pi = (\mathbf{a}, \mathbf{b}, \mathbf{c}) \in S_3$ and $\mathbf{f} = (\mathbf{h}, w, q) \in (0, \infty)^3$ and $\sigma_* \in \{-1, +1\}^3$ let $\Phi_1 = \Phi_1(\pi, \mathbf{f}, \sigma_*) : \mathbb{R} \rightarrow \mathcal{D}(\sigma_*)$ be given by

$$\begin{aligned} A_{\mathbf{a}}[\Phi_1](\tau) &= A_{\mathbf{a}}[\Phi_*] & \alpha_{\mathbf{p}, \mathbf{a}}[\Phi_1](\tau) &= \alpha_{\mathbf{p}, \mathbf{a}}[\Phi_*] \\ \theta_{\mathbf{a}}[\Phi_1, \mathbf{h}](\tau) &= \theta_{\mathbf{a}}[\Phi_*, \mathbf{h}] & \xi_{\mathbf{p}, \mathbf{a}}[\Phi_1, \mathbf{h}](\tau) &= \xi_{\mathbf{p}, \mathbf{a}}[\Phi_*, \mathbf{h}] + (\tau - \tau_{1+}) \alpha_{\mathbf{p}, \mathbf{a}}[\Phi_*] \end{aligned}$$

for all $\tau \in \mathbb{R}$ and $\mathbf{p} \in \{\mathbf{b}, \mathbf{c}\}$. Here, $\tau_{1+} = \tau_{1+}(\mathbf{f})$ and $\Phi_* = \Phi_*(\pi, \mathbf{f}, \sigma_*)$.

Lemma 3.8. For all $\pi = (\mathbf{a}, \mathbf{b}, \mathbf{c}) \in S_3$, $\mathbf{f} = (\mathbf{h}, w, q) \in (0, \infty)^3$, $\sigma_* \in \{-1, +1\}^3$, set $\Phi_0 = \Phi_0(\pi, \mathbf{f}, \sigma_*)$ and $\Phi_1 = \Phi_1(\pi, \mathbf{f}, \sigma_*)$ and $\tau_{1+} = \tau_{1+}(\mathbf{f})$ and $d_{\mathcal{D}} = d_{\mathcal{D}(\sigma_*)}(\pi, \mathbf{h})$ and $\phi_{\mathcal{D}} = \phi_{\mathcal{D}(\sigma_*)}(\mathbf{h})$. Then

- (a) $|\beta_{\mathbf{a}}[\Phi_1](\tau_{1+})| = |\beta_{\mathbf{b}}[\Phi_1](\tau_{1+})|$
- (b) $c[\Phi_1, \mathbf{h}, Z](\tau_{1+}) = 0$, see Definitions 3.1 and 3.2 for c and Z , respectively
- (c) $\phi_{\mathcal{D}}(\Phi_0(\tau_{1+}), \Phi_1(\tau_{1+})) \leq 2^7 \max\{1 + w, \mathbf{h}\} \exp(-\frac{1}{2\mathbf{h}} \min\{1, w + q\})$
- (d) $d_{\mathcal{D}}(\Phi_0(\tau), \Phi_1(\tau)) \leq (1 + |\tau - \tau_{1+}|) d_{\mathcal{D}}(\Phi_0(\tau_{1+}), \Phi_1(\tau_{1+}))$ for all $\tau \in \mathbb{R}$

Proof. We discuss (c) only. By direct calculation,

$$\begin{aligned} \alpha_{\mathbf{a}}[\Phi_0](\tau_{1+}) - \alpha_{\mathbf{a}}[\Phi_1](\tau_{1+}) &= -X & \xi_{\mathbf{a}}[\Phi_0, \mathbf{h}](\tau_{1+}) - \xi_{\mathbf{a}}[\Phi_1, \mathbf{h}](\tau_{1+}) &= -Y \\ \alpha_{\mathbf{b}}[\Phi_0](\tau_{1+}) - \alpha_{\mathbf{b}}[\Phi_1](\tau_{1+}) &= +X & \xi_{\mathbf{b}}[\Phi_0, \mathbf{h}](\tau_{1+}) - \xi_{\mathbf{b}}[\Phi_1, \mathbf{h}](\tau_{1+}) &= +Y \\ \alpha_{\mathbf{c}}[\Phi_0](\tau_{1+}) - \alpha_{\mathbf{c}}[\Phi_1](\tau_{1+}) &= +X + \mu & \xi_{\mathbf{c}}[\Phi_0, \mathbf{h}](\tau_{1+}) - \xi_{\mathbf{c}}[\Phi_1, \mathbf{h}](\tau_{1+}) &= +Y \end{aligned}$$

with $X = -1 + \tanh(\frac{1}{\mathbf{h}}\tau_{1+})$ and $Y = \mathbf{h} \log(1 + \exp(-2\frac{1}{\mathbf{h}}\tau_{1+}))$. The estimates

$$\begin{aligned} |X| &\leq 2 \exp(-2\frac{1}{\mathbf{h}}\tau_{1+}) \leq 2 \exp(-\frac{1}{\mathbf{h}}) \\ |Y| &\leq \mathbf{h} \exp(-2\frac{1}{\mathbf{h}}\tau_{1+}) \leq \mathbf{h} \exp(-\frac{1}{\mathbf{h}}) \\ |\mu| &\leq (1 + w)2^6 \exp(-\frac{1}{2\mathbf{h}} \min\{1, w + q\}) \end{aligned}$$

imply (c). $\square$

Definition 3.16. This is, verbatim, Definition 1.2 in the Introduction.

Lemma 3.9. In the context of Definition 3.16, the identities

$$\lambda_L = 1 - \alpha_{\mathbf{a}, \mathbf{a}'}[\Phi_0](\tau_{1-}) = 1 - \alpha_{\mathbf{a}, \mathbf{a}'}[\Phi_0](\tau) \quad (3.14a)$$

$$w_L = -(\alpha_{\mathbf{a}, \mathbf{a}'}[\Phi_0](\tau_{1-}))^{-1} = -(\alpha_{\mathbf{a}, \mathbf{a}'}[\Phi_0](\tau))^{-1} \quad (3.14b)$$

$$\frac{\mathbf{h}}{\mathbf{h}_L} = \frac{1+2w_L}{1+w_L} (-\tau_{1-} + \mathbf{h} \log \lambda_L) - \mathbf{h} \log 2 \quad (3.14c)$$

$$\begin{aligned} q_L = \frac{1}{1+w_L} \left(\mathbf{h}_L \log \lambda_L - \frac{\mathbf{h}_L}{\mathbf{h}} \xi_{\mathbf{a}, \mathbf{c}'}[\Phi_0, \mathbf{h}](\tau_{1-}) + \frac{\mathbf{h}_L}{\mathbf{h}} \tau_{1-} - \frac{w_L(1+w_L)}{1+2w_L} \right. \\ \left. - \frac{1+3w_L+(w_L)^2}{1+2w_L} \mathbf{h}_L \log 2 \right) \end{aligned} \quad (3.14d)$$

hold, where $\Phi_0 = \Phi_0(\pi, \mathbf{f}, \sigma_*)$ and $\tau_{1-} = \tau_{1-}(\mathbf{f})$ and $\tau \in \mathbb{R}$. Furthermore,

$$(\xi_{\mathbf{a}}[\Phi_0, \mathbf{h}] - \xi_{\mathbf{a}'}[\Phi_0, \mathbf{h}]) F = \tau - \tau_{1-} - 2\mathbf{h} \log(1 + e^{2\tau/\mathbf{h}}) F \quad (3.15)$$

for all $\tau \in \mathbb{R}$, where

$$F = \begin{cases} \frac{1}{3+w} & q \leq 1 \\ \frac{1+w}{3+2w} & q > 1 \end{cases} \quad (3.16)$$

Proof. By direct calculation. In each case, distinguish $q \leq 1$ and $q > 1$. $\square$

Definition 3.17. For all $\mathbf{f} = (\mathbf{h}, w, q) \in (0, \infty)^3$ set

$$\tau_*(\mathbf{f}) = \begin{cases} \frac{q}{1+w} & \text{if } q \leq 1 \\ 1 & \text{if } q > 1 \end{cases}$$

Definition 3.18. For all $\mathbf{f} = (\mathbf{h}, w, q) \in (0, \infty)^3$ set

$$\mathbf{K}(\mathbf{f}) = 2^{40} \left(\frac{1}{\mathbf{h}}\right)^2 \max\left\{\left(\frac{1}{w}\right)^2, w^3\right\} \max\left\{\left(\frac{1}{q}\right)^2, q\right\} \exp\left(-\frac{1}{\mathbf{h}} 2^{-7} \tau_*(\mathbf{f})\right) \quad (3.17)$$

Definition 3.19. Let $\mathcal{F}$ be the open set of all $\mathbf{f} = (\mathbf{h}, w, q) \in (0, \infty)^3$ for which

$$q \neq 1 \quad \mathbf{K}(\mathbf{f}) < 1 \quad \mathbf{h} < 2^{-7} \tau_*(\mathbf{f}) \quad (3.18)$$

Proposition 3.3. For all $\pi = (\mathbf{a}, \mathbf{b}, \mathbf{c}) \in S_3$, $\sigma_* \in \{-1, +1\}^3$, there are **unique** maps

$$\begin{aligned} \Pi &= \Pi[\pi, \sigma_*] : \quad \mathcal{F} \rightarrow (0, \infty)^2 \times \mathbb{R} \\ \Lambda &= \Lambda[\pi, \sigma_*] : \quad \mathcal{F} \rightarrow [1, \infty) \\ \tau_{2-} &= \tau_{2-}[\pi, \sigma_*] : \quad \mathcal{F} \rightarrow (-\infty, 0) \end{aligned}$$

so that for all $\mathbf{f} = (\mathbf{h}, w, q) \in \mathcal{F}$ (see Definitions 3.5, 3.7, 3.9, 3.10, 3.12, 3.14, 3.16)

- (a) $\|\Pi(\mathbf{f}) - \mathcal{Q}_L(\mathbf{f})\|_{\mathbb{R}^3} \leq \mathbf{K}(\mathbf{f})$
- (b) $|\Lambda(\mathbf{f}) - \lambda_L(\mathbf{f})| \leq \mathbf{K}(\mathbf{f})$
- (c) $\tau_-(\mathbf{f}) < \tau_{2-}(\mathbf{f}) < \frac{1}{2}\tau_{1-}(\mathbf{f})$ and $|\tau_{2-}(\mathbf{f}) - \tau_{1-}(\mathbf{f})| \leq \mathbf{K}(\mathbf{f})$
- (d) Π , Λ and τ_{2-} are continuous
- (e) if we set $\tau_{2-} = \tau_{2-}(\mathbf{f})$, $\tau_{2+} = \tau_{1+}(\mathbf{f})$, $\pi' = (\mathbf{a}', \mathbf{b}', \mathbf{c}') = \mathcal{P}_L(\pi, \mathbf{f})$, $\lambda = \Lambda(\mathbf{f})$ and $\mathbf{f}' = (\mathbf{h}', w', q') = \Pi(\mathbf{f})$, then $\frac{1}{2} \leq \tau_{2+} - \tau_{2-} \leq 3$ and there is a smooth field

$$\Phi = \alpha \oplus \beta \in \mathcal{E} = \mathcal{E}(\sigma_*; \tau_{2-}, \tau_{2+})$$

that satisfies

(e.1) $(\mathbf{a}, \mathbf{b}, \mathbf{c})[\Phi, \mathbf{h}, Z] = 0$ on $[\tau_{2-}, \tau_{2+}]$

(e.2) $\Phi(\tau_{2+}) = \Phi_*(\pi, \mathbf{f}, \sigma_*)$ and $\Phi(\tau_{2-}) = \lambda \Phi_*(\pi', \mathbf{f}', \sigma_*)$, in particular

$$\Phi(\tau_{2+}) \in \mathcal{H}(\pi, \sigma_*) \quad \text{and} \quad \Phi(\tau_{2-}) \in \mathcal{H}(\pi', \sigma_*)$$

(e.3) $|\beta_{\mathbf{a}}[\Phi](\tau)| \geq |\beta_{\mathbf{a}'}[\Phi](\tau)|$ for all $\tau \in [\tau_{2-}, \frac{1}{2}\tau_{1-}(\mathbf{f})]$ with equality iff $\tau = \tau_{2-}$

(e.4) $d_{\mathcal{E}, (\pi, \mathbf{h})}(\Phi, \Phi_0) \leq \mathbf{K}(\mathbf{f})$, where $\Phi_0 = \Phi_0(\pi, \mathbf{f}, \sigma_*)|_{[\tau_{2-}, \tau_{2+}]}$

(e.5) $\sup_{\tau \in [\tau_{2-}, \tau_{2+}]} \max\{\alpha_{\mathbf{b}, \mathbf{c}}[\Phi], \alpha_{\mathbf{c}, \mathbf{a}}[\Phi], \alpha_{\mathbf{a}, \mathbf{b}}[\Phi]\}(\tau) \leq -2^{-2} \min\{w^2, w^{-1}\}$

Proof. The main part of this proof is the construction of the field Φ that appears in (e). To make the proof more transparent, we replace some numerical constants in (3.17) and (3.18) by the components of a parameter vector $\ell = (\ell_1, \dots, \ell_8) \in \mathbb{R}^8$. In the course of the construction of Φ , we require a finite number of inequalities of the form $\ell \geq \ell'$. Each inequality of this kind is marked by $(\bullet)$ and is *assumed to hold for the rest of the proof*, once it has been stated. At the end of the construction, we check that the particular parameters appearing in (3.17) and (3.18) satisfy all these inequalities.

Let $\pi = (\mathbf{a}, \mathbf{b}, \mathbf{c}) \in S_3$ and $\sigma_* \in \{-1, +1\}^3$. Fix any $\mathbf{f} = (\mathbf{h}, w, q) \in (0, \infty)^3$ with $q \neq 1$ and $\mathbf{h} \leq 1$. Set $\tau_* = \tau_*(\mathbf{f})$. For any $s = (s_1, \dots, s_7) \in \mathbb{R}^7$, set

$$\begin{aligned} \mathbf{X}(s) &= \mathbf{X}(s_1, \dots, s_7) = \\ &2^{s_1} \left(\frac{1}{\mathbf{h}}\right)^{s_2} \times \begin{cases} \left(\frac{1}{w}\right)^{s_3} & \text{if } w \leq 1 \\ w^{s_4} & \text{if } w > 1 \end{cases} \times \begin{cases} \left(\frac{1}{q}\right)^{s_5} & \text{if } q \leq 1 \\ q^{s_6} & \text{if } q > 1 \end{cases} \times \exp\left(\frac{1}{\mathbf{h}} s_7 \tau_*\right) \end{aligned}$$

Basic properties of $\mathbf{X}(s)$. The quantity $\mathbf{X}(s)$ is positive, non-decreasing in each of its seven arguments (recall $0 < \mathbf{h} \leq 1$), and $\mathbf{X}(s)\mathbf{X}(s') = \mathbf{X}(s + s')$ for all $s, s' \in \mathbb{R}^7$, and $\mathbf{X}(0, \dots, 0) = 1$. Also, we have $\tau_* \geq \mathbf{X}(-1, 0, 0, -1, -1, 0, 0)$.

Basic smallness assumptions. Introduce a parameter vector $\ell = (\ell_1, \dots, \ell_8) \in \mathbb{R}^8$ with

$$(\ell_1, \dots, \ell_7) \geq (0, 0, 0, 0, 0, 0, -\infty) \quad \text{and} \quad \ell_8 \geq 0 \quad (\bullet)_1$$

Our basic assumptions on the vector $\mathbf{f} = (\mathbf{h}, w, q)$ are:

$$q \neq 1 \quad \mathbf{K} \stackrel{\text{def}}{=} \mathbf{X}(\ell_1, \dots, \ell_7) < 1 \quad \mathbf{h} < 2^{-\ell_8} \tau_* \quad (3.19)$$

Observe that our previous assumptions $q \neq 1$ and $\mathbf{h} \leq 1$ are subsumed in (3.19).

Abbreviations. $\tau_{\pm} = \tau_{\pm}(\mathbf{f})$ and $\tau_{1\pm} = \tau_{1\pm}(\mathbf{f})$ and $\tau_{2+} = \tau_{1+}(\mathbf{f})$ and

$$\tau_{0-} = \frac{1}{2}\tau_{1-} + \frac{1}{2}\tau_- < 0 \quad \tau_{0+} = \tau_{1+} > 0$$

and $\mathcal{D} = \mathcal{D}(\sigma_*)$ and $\mathcal{E} = \mathcal{E}(\sigma_*; \tau_{0-}, \tau_{0+})$ and $\Phi_0 = \Phi_0(\pi, \mathbf{f}, \sigma_*)|_{[\tau_{0-}, \tau_{0+}]}$ and $\Phi_1 = \Phi_1(\pi, \mathbf{f}, \sigma_*)|_{[\tau_{0-}, \tau_{0+}]}$ and $\Phi_* = \Phi_*(\pi, \mathbf{f}, \sigma_*)$ and $d_{\mathcal{E}} = d_{\mathcal{E}, (\pi, \mathbf{h})}$ and $d_{\mathcal{D}} = d_{\mathcal{D}, (\pi, \mathbf{h})}$ and $\phi_{\mathcal{D}} = \phi_{\mathcal{D}, \mathbf{h}}$ and $B_{\mathcal{E}}[\cdot, \cdot] = B_{\mathcal{E}, (\pi, \mathbf{h})}[\cdot, \cdot]$ and $\pi' = (\mathbf{a}', \mathbf{b}', \mathbf{c}') = \mathcal{P}_L(\pi, \mathbf{f})$.

Preliminaries 1. Introduce ϵ_- and ϵ_+ by $\tau_{0-} = \tau_- + \epsilon_-$ and $\tau_{0+} = \tau_+ - \epsilon_+$, just as in Lemma 3.4. We have

$$\begin{aligned} \epsilon_+ &= \tau_+ - \tau_{1+} = \frac{1+w}{1+2w} \left(1 + \frac{1}{w} - \mathbf{h} \log 2\right) \\ \epsilon_- &= \frac{1}{2}(\tau_{1-} - \tau_-) = \begin{cases} \frac{1}{2(3+w)} \left(\frac{1+w}{2+w} q - \mathbf{h} \log 2\right) & \text{if } q < 1 \\ \frac{1+w}{2(3+2w)} \left(\frac{1+w}{2+w} - \mathbf{h} \log 2\right) & \text{if } q > 1 \end{cases} \end{aligned}$$

Require $\ell_8 \geq 2 (\bullet)_2$. Then $\mathbf{h} \log 2 \leq \mathbf{h} \leq 2^{-2} \min\{1, q\}$, and (recall that $\tau_+ = 1 + \frac{1}{w}$)

$$2^{-2} \leq \epsilon_+/\tau_+ \leq 1 \quad 2^{-5} \leq \epsilon_-/\tau_* \leq 2^{-1}$$

and $\epsilon_- \in (0, -\tau_-)$ and $\epsilon_+ \in (0, \tau_+)$, as required by Lemma 3.4. We have

$$-1 < \tau_- < \tau_{0-} < \tau_{1-} < 0 < \frac{1}{2} < \tau_{0+} = \tau_{1+} = \tau_{2+} < \min\{2, \tau_+\}$$

Set

$$\delta \stackrel{\text{def}}{=} 2^{-9} \min\{1, w\} \tau_* = \mathbf{X}(-9, 0, -1, 0, 0, 0, 0) \tau_* \geq \mathbf{X}(-10, 0, -1, -1, -1, 0, 0) \quad (3.20)$$

This implies $\delta \leq 2^{-4} \min\{1, w, \epsilon_-, \frac{\epsilon_+}{\tau_+ \tau_{0+}}\}$, the main hypothesis of Lemma 3.4. This lemma will be applied later.

Preliminaries 2. Require $\ell_8 \geq 7 (\bullet)_3$. Then

$$\begin{aligned} d_{\mathcal{E}}(\Phi_0, \Phi_1) &\leq 2^2 d_{\mathcal{D}}(\Phi_0(\tau_{1+}), \Phi_*) \leq 2^{11} \phi_{\mathcal{D}}(\Phi_0(\tau_{1+}), \Phi_*) \\ &\leq 2^{18} (1+w) \exp\left(-\frac{1}{2\mathbf{h}} \min\{1, q\}\right) \leq 2^{18} (1+w) \exp\left(-\frac{1}{2\mathbf{h}} \tau_*\right) \\ &\leq \mathbf{X}(19, 0, 0, 1, 0, 0, -2^{-1}) \leq 2^{-2} \delta \mathbf{X}(31, 0, 1, 2, 1, 0, -2^{-1}) \quad (3.21) \end{aligned}$$

The first and third inequality follow from (d) and (c) in Lemma 3.8, respectively, using $\sup_{\tau \in [\tau_{0-}, \tau_{0+}]} (1 + |\tau - \tau_{1+}|) \leq 2^2$. The second inequality follows from Lemma 3.2 (b), with $C = D = 2$. Its assumptions are satisfied, because $\mathbf{h} \leq 2^{-7}$ and $A_{\mathbf{a}}[\Phi_0](\tau_{1+}) = 1$ and $\mathbf{h} \varphi_{\mathbf{a}}[\Phi_0](\tau_{1+}) = \tau_{1+} \in [\frac{1}{2}, 2]$ and $\xi_{\mathbf{a}}[\Phi_*, \mathbf{h}] \in [-\frac{3}{2}, -\frac{1}{2}]$ and $|\beta_{\mathbf{a}}[\Phi_*]| \leq 2 \exp(-\frac{1}{2\mathbf{h}}) < 1$ and $A_{\mathbf{a}}[\Phi_*] \in [1, 2]$ and $0 \leq \mathbf{h} |\varphi_{\mathbf{a}}[\Phi_*]| + \xi_{\mathbf{a}}[\Phi_*, \mathbf{h}] \leq 2^{-3}$

(see Remark 3.2) and $\mathbf{h}|\varphi_{\mathbf{a}}[\Phi_*]| \in [\frac{1}{2}, 2]$ and $\text{sgn } \varphi_{\mathbf{a}}[\Phi_*] = -\text{sgn } \alpha_{\mathbf{a}}[\Phi_*] = +1$, and because $\ell_8 \geq 7$ implies $\mathbf{h} \leq 2^{-7}$ and therefore $\exp(-\frac{1}{\mathbf{h}}2^{-3}) \leq 2^{-12}$.

Require $(\ell_1, \dots, \ell_7) \geq (31, 0, 1, 2, 1, 0, -2^{-1}) (\bullet)_4$. Then, by (3.19),

$$\Phi_1 \in B_{\mathcal{E}}[2^{-2}\delta, \Phi_0] \quad (3.22)$$

Construction of Φ . Define a map $P : B_{\mathcal{E}}[\delta, \Phi_0] \rightarrow B_{\mathcal{E}}[\delta, \Phi_0], \Psi \mapsto P(\Psi)$ by

$$A_{\mathbf{a}}[P(\Psi)](\tau) - A_{\mathbf{a}}[\Phi_1](\tau) = \int_{\tau_{0+}}^{\tau} d\tau' \mathbf{I}_1[\Psi, \mathbf{h}, \pi](\tau') \quad (3.23a)$$

$$\theta_{\mathbf{a}}[P(\Psi), \mathbf{h}](\tau) - \theta_{\mathbf{a}}[\Phi_1, \mathbf{h}](\tau) = \int_{\tau_{0+}}^{\tau} d\tau' \mathbf{I}_2[\Psi, \mathbf{h}, \pi](\tau') \quad (3.23b)$$

$$\alpha_{\mathbf{p}, \mathbf{a}}[P(\Psi)](\tau) - \alpha_{\mathbf{p}, \mathbf{a}}[\Phi_1](\tau) = \int_{\tau_{0+}}^{\tau} d\tau' \mathbf{I}_{(3, \mathbf{p})}[\Psi, \mathbf{h}, \pi](\tau') \quad (3.23c)$$

$$\xi_{\mathbf{p}, \mathbf{a}}[P(\Psi), \mathbf{h}](\tau) - \xi_{\mathbf{p}, \mathbf{a}}[\Phi_1, \mathbf{h}](\tau) = \int_{\tau_{0+}}^{\tau} d\tau'' \int_{\tau_{0+}}^{\tau''} d\tau' \mathbf{I}_{(3, \mathbf{p})}[\Psi, \mathbf{h}, \pi](\tau') \quad (3.23d)$$

for all $\mathbf{p} \in \{\mathbf{b}, \mathbf{c}\}$ and $\tau \in [\tau_{0-}, \tau_{0+}]$. To make sure that P is well defined, we require $(\ell_1, \dots, \ell_7) \geq (28, 2, 1, 1, 1, 0, -2^{-7}) (\bullet)_5$, in which case Lemma 3.6 (see Preliminaries 1) implies the uniform estimates

$$|\mathbf{I}_S[\Psi, \mathbf{h}, \pi]| \leq \mathbf{X}(12, 1, 0, 0, 0, 0, -2^{-7}) \leq 2^{-6}\delta \mathbf{X}(28, 1, 1, 1, 1, 0, -2^{-7}) \leq 2^{-6}\delta \quad (3.24a)$$

$$|\mathbf{I}_S[\Psi, \mathbf{h}, \pi] - \mathbf{I}_S[\Psi', \mathbf{h}, \pi]| \leq 2^{-5}\mathbf{X}(24, 2, 0, 0, 0, 0, -2^{-7})d_{\mathcal{E}}(\Psi, \Psi') \leq 2^{-5}d_{\mathcal{E}}(\Psi, \Psi') \quad (3.24b)$$

on the interval $[\tau_{0-}, \tau_{0+}]$, for all $\Psi, \Psi' \in B_{\mathcal{E}}[\delta, \Phi_0]$ and all $S \in \{1, 2, (3, \mathbf{b}), (3, \mathbf{c})\}$. Since $\sup_{\tau \in [\tau_{0-}, \tau_{0+}]} |\tau - \tau_{0+}| \leq 4$, we have:

- $A_{\mathbf{a}}[P(\Psi)] > \frac{1}{2}$ on $[\tau_{0-}, \tau_{0+}]$, which makes $P(\Psi)$ a well defined element of $\mathcal{E}$.
- Each right hand side of (3.23) is $\leq 2^{-2}\delta$, hence $P(\Psi) \in B_{\mathcal{E}}[\frac{1}{2}\delta, \Phi_0]$.
- The map P is Lipschitz-continuous with constant $\leq \frac{1}{2}$.

The metric space $B_{\mathcal{E}}[\delta, \Phi_0]$ is nonempty and complete. By the Banach Fixed Point Theorem, the contraction P admits a unique fixed point

$$\Phi \in B_{\mathcal{E}}[\frac{1}{2}\delta, \Phi_0] \quad (3.25)$$

Proof that the fixed point satisfies $(\mathbf{a}, \mathbf{b}, c)[\Phi, \mathbf{h}, Z] = 0$. The fixed point Φ is smooth. We have $\Phi(\tau_{0+}) = \Phi_1(\tau_{0+}) = \Phi_*$ and $c[\Phi, \mathbf{h}, Z](\tau_{0+}) = 0$, by Lemma 3.8 (b), and because $\tau_{0+} = \tau_{1+}$. Set $\Psi = P(\Psi) = \Phi$ in (3.23) and differentiate with respect to τ . The result of differentiating (3.23a) and (3.23b) can be written as

$$\begin{aligned} \frac{d}{d\tau} \begin{pmatrix} A_{\mathbf{a}} \\ \theta_{\mathbf{a}} \end{pmatrix} &= \frac{1}{(A_{\mathbf{a}})^2} \begin{pmatrix} \frac{1}{\mathbf{h}}(A_{\mathbf{a}})^2 \tanh \varphi_{\mathbf{a}} & \frac{1}{\mathbf{h}}(A_{\mathbf{a}})^2 \text{sech } \varphi_{\mathbf{a}} \\ \varphi_{\mathbf{a}} \tanh \varphi_{\mathbf{a}} - 1 & \sinh \varphi_{\mathbf{a}} + \varphi_{\mathbf{a}} \text{sech } \varphi_{\mathbf{a}} \end{pmatrix} \\ &\quad \times \begin{pmatrix} \mathbf{a}_{\mathbf{a}}[\Phi, \mathbf{h}, B_{\mathbf{a}}] - \mathbf{a}_{\mathbf{a}}[\Phi, \mathbf{h}, Z] \\ -(\sigma_*)_{\mathbf{a}} \mathbf{b}_{\mathbf{a}}[\Phi, \mathbf{h}, B_{\mathbf{a}}] + (\sigma_*)_{\mathbf{a}} \mathbf{b}_{\mathbf{a}}[\Phi, \mathbf{h}, Z] \end{pmatrix} \end{aligned}$$

where $A_{\mathbf{a}} = A_{\mathbf{a}}[\Phi]$, $\theta_{\mathbf{a}} = \theta_{\mathbf{a}}[\Phi, \mathbf{h}]$, $\varphi_{\mathbf{a}} = \varphi_{\mathbf{a}}[\Phi]$, because $\mathbf{b}_{\mathbf{a}}[\Phi, \mathbf{h}, B_{\mathbf{a}}] = \mathbf{b}_{\mathbf{a}}[\Phi, \mathbf{h}, Z]$. Now, Lemma 3.1 implies $\mathbf{a}_{\mathbf{a}}[\Phi, \mathbf{h}, Z] = \mathbf{b}_{\mathbf{a}}[\Phi, \mathbf{h}, Z] = 0$. Differentiation of (3.23c) gives $\frac{d}{d\tau} \alpha_{\mathbf{p}, \mathbf{a}}[\Phi] = \frac{1}{\mathbf{h}} \mathbf{a}_{\mathbf{p}}[\Phi, \mathbf{h}, Z, B_{\mathbf{a}}] + \frac{1}{\mathbf{h}} \mathbf{a}_{\mathbf{a}}[\Phi, \mathbf{h}, Z, B_{\mathbf{a}}]$. Together with $\mathbf{a}_{\mathbf{a}}[\Phi, \mathbf{h}, Z] = 0$ and the general identity $\mathbf{a}_{\mathbf{p}}[\Phi, \mathbf{h}, B_{\mathbf{a}}] + \mathbf{a}_{\mathbf{a}}[\Phi, \mathbf{h}, B_{\mathbf{a}}] = -\mathbf{h} \frac{d}{d\tau} \alpha_{\mathbf{p}, \mathbf{a}}[\Phi]$, we obtain $\mathbf{a}_{\mathbf{p}}[\Phi, \mathbf{h}, Z] = 0$. Differentiating (3.23d) and simplifying the result with (3.23c) gives

$\frac{d}{d\tau}\xi_{\mathbf{p},\mathbf{a}}[\Phi, \mathbf{h}] = \alpha_{\mathbf{p},\mathbf{a}}[\Phi]$ which, by $\mathbf{b}_{\mathbf{a}}[\Phi, \mathbf{h}, Z] = 0$, implies $\mathbf{b}_{\mathbf{p}}[\Phi, \mathbf{h}, Z] = 0$. Now, Proposition 3.2 and the fact that $c[\Phi, \mathbf{h}, Z](\tau_{0+}) = 0$ imply that $c[\Phi, \mathbf{h}, Z] = 0$ identically on $[\tau_{0-}, \tau_{0+}]$.

Estimates on Φ . By the fixed point equation $P(\Phi) = \Phi$ and by (3.21) and (3.24a),

$$\begin{aligned} d_{\mathcal{E}}(\Phi_0, \Phi) &\leq d_{\mathcal{E}}(\Phi_0, \Phi_1) + d_{\mathcal{E}}(\Phi_1, P(\Phi)) \\ &\leq \mathbf{X}(19, 0, 0, 1, 0, 0, -2^{-1}) + \mathbf{X}(16, 1, 0, 0, 0, 0, -2^{-7}) \\ &\leq \mathbf{X}(20, 1, 0, 1, 0, 0, -2^{-7}) \end{aligned} \quad (3.26)$$

We require $(\ell_1, \dots, \ell_7) \geq (20, 1, 0, 1, 0, 0, -2^{-7})$ $(\bullet)_6$, which implies $d_{\mathcal{E}}(\Phi_0, \Phi) \leq \mathbf{K}$. To apply Lemma 3.2 (a), set $\mathcal{J} = [\tau_{0-}, \frac{1}{2}\tau_{1-}] \subset [\tau_{0-}, \tau_{0+}]$ and $C = 2$ and $D = 12 \max\{1, q^{-1}\}$. We check the assumptions of Lemma 3.2. The inequalities

$$C^{-1} \leq A_{\mathbf{a}}[X] \leq C \quad D^{-1} \leq \mathbf{h}|\varphi_{\mathbf{a}}[X]| \leq D$$

hold for both $X = \Phi_0(\tau)$ and $X = \Phi(\tau)$, for all $\tau \in \mathcal{J}$. The inequality for $A_{\mathbf{a}}$ follows from $A_{\mathbf{a}}[\Phi_0](\tau) = 1$ and the bound $d_{\mathcal{E}}(\Phi_0, \Phi) \leq \delta \leq 2^{-9}$. To check the inequality for $\varphi_{\mathbf{a}}$, observe that $\mathbf{h}\varphi_{\mathbf{a}}[\Phi_0](\tau) = \tau \in \mathcal{J} \subset [-(D/2), -(D/2)^{-1}]$, see the definitions of τ_{-} and τ_{1-} . Furthermore, for all $\tau \in \mathcal{J}$, we have

$$|\mathbf{h}\varphi_{\mathbf{a}}[\Phi] - \mathbf{h}\varphi_{\mathbf{a}}[\Phi_0]| \leq |\tau| |A_{\mathbf{a}}[\Phi] - 1| + A_{\mathbf{a}}[\Phi] |\theta_{\mathbf{a}}[\Phi, \mathbf{h}]| \leq 4\delta \leq (2D)^{-1}$$

This implies $\mathbf{h}\varphi_{\mathbf{a}}[\Phi](\tau) \in [-D, -D^{-1}]$ and $\text{sgn } \varphi_{\mathbf{a}}[\Phi_0](\tau) = \text{sgn } \varphi_{\mathbf{a}}[\Phi](\tau) = -1$ for all $\tau \in \mathcal{J}$. Now, Lemma 3.2 (a) and $2^3 C^2 D \leq \mathbf{X}(9, 0, 0, 0, 1, 0, 0)$ imply for all $\tau \in \mathcal{J}$

$$\phi_{\mathcal{D}}(\Phi_0(\tau), \Phi(\tau)) \leq 2^3 C^2 D d_{\mathcal{E}}(\Phi_0, \Phi) \leq \mathbf{X}(29, 1, 0, 1, 1, 0, -2^{-7}) \stackrel{\text{def}}{=} \mathbf{M} \quad (3.27)$$

Construction of τ_{2-} . Recall that $\mathbf{a}' = \mathbf{c}$ if $q < 1$ and $\mathbf{a}' = \mathbf{b}$ if $q > 1$. By (3.15),

$$(\xi_{\mathbf{a}}[\Phi_0, \mathbf{h}] - \xi_{\mathbf{a}'}[\Phi_0, \mathbf{h}]) F = \tau - \tau_{1-} - T_1 \quad (3.28a)$$

$$(\xi_{\mathbf{a}}[\Phi, \mathbf{h}] - \xi_{\mathbf{a}'}[\Phi, \mathbf{h}]) F = \tau - \tau_{1-} - T_2 \quad (3.28b)$$

for all $\tau \in \mathcal{J}$, where F is given by (3.16), and

$$T_1 = T_1(\tau) = 2\mathbf{h} \log(1 + e^{2\tau/\mathbf{h}}) F$$

$$T_2 = T_2(\tau) = T_1 - (\xi_{\mathbf{a}}[\Phi, \mathbf{h}] - \xi_{\mathbf{a}}[\Phi_0, \mathbf{h}]) F + (\xi_{\mathbf{a}'}[\Phi, \mathbf{h}] - \xi_{\mathbf{a}'}[\Phi_0, \mathbf{h}]) F$$

For all $\tau \in \mathcal{J}$ we have $0 < T_1 \leq 2\mathbf{h} e^{2\tau/\mathbf{h}} F \leq 2\mathbf{h} e^{\tau_{1-}/\mathbf{h}} F \leq \mathbf{M}F$ and therefore $|T_2| \leq 3\mathbf{M}F \leq \frac{3}{2}\mathbf{M}$. Estimate

$$\text{dist}_{\mathbb{R}}(\tau_{1-}, \mathbb{R} \setminus \mathcal{J}) = \min \left\{ \frac{1}{2}|\tau_{1-}|, \epsilon_- \right\} \geq 2^{-5}\tau_* \geq \mathbf{X}(-6, 0, 0, -1, -1, 0, 0)$$

Therefore, the condition $(\ell_1, \dots, \ell_7) \geq (37, 1, 0, 2, 2, 0, -2^{-7})$ $(\bullet)_7$ yields

$$|T_2| \leq \frac{1}{2}\text{dist}_{\mathbb{R}}(\tau_{1-}, \mathbb{R} \setminus \mathcal{J}) \quad (3.29)$$

for all $\tau \in \mathcal{J}$. Set

$$\tau_{2-} = \sup \left\{ \tau \in \mathcal{J} \mid \xi_{\mathbf{a}}[\Phi, \mathbf{h}](\tau) \leq \xi_{\mathbf{a}'}[\Phi, \mathbf{h}](\tau) \right\} \quad (3.30)$$

The set on the right is nonempty, by (3.28b) and (3.29), it contains τ_{0-} . We have $\tau_{2-} \in (\tau_{0-}, \frac{1}{2}\tau_{1-}) \subset \mathcal{J}$ and, by continuity, $\xi_{\mathbf{a}}[\Phi, \mathbf{h}](\tau_{2-}) = \xi_{\mathbf{a}'}[\Phi, \mathbf{h}](\tau_{2-})$, and $|\tau_{2-} -$

$\tau_{1-} \leq \frac{3}{2}\mathbf{M}$. For all $\tau \in [\tau_{2-}, \frac{1}{2}\tau_{1-}]$, we have $|\beta_{\mathbf{a}}[\Phi](\tau)| \geq |\beta_{\mathbf{a}'}[\Phi](\tau)|$ with equality iff $\tau = \tau_{2-}$. The condition $(\ell_1, \dots, \ell_7) \geq (31, 1, 0, 1, 1, 0, -2^{-7}) (\bullet)_8$ implies $|\tau_{2-} - \tau_{1-}| \leq \mathbf{K}$.

Estimates on Φ_0 . For all $\tau \in \mathcal{J}$, we have

$$\begin{aligned} |\alpha_{\mathbf{a}}[\Phi_0](\tau) - 1| &= |\tanh \frac{1}{\mathbf{h}}|\tau| - 1| \leq 2 \exp(-\frac{2}{\mathbf{h}}|\tau|) \\ |\xi_{\mathbf{a}}[\Phi_0, \mathbf{h}](\tau) - \tau| &= |\mathbf{h} \log(2 \cosh \frac{1}{\mathbf{h}}|\tau|) - |\tau|| \leq \mathbf{h} \exp(-\frac{2}{\mathbf{h}}|\tau|) \\ \exp(-\frac{2}{\mathbf{h}}|\tau|) &\leq \exp(-\frac{1}{\mathbf{h}}|\tau_{1-}|) \leq \exp(-\frac{1}{\mathbf{h}}2^{-2}\tau_*) \leq 2^{-29}\mathbf{M} \end{aligned}$$

These estimates will be used without further comment.

Construction of λ . Set $\lambda_L = \lambda_L(\mathbf{f})$ and recall (3.14a). Set

$$\lambda = -\alpha_{\mathbf{a}'}[\Phi](\tau_{2-}) \quad (3.31)$$

Then,

$$\begin{aligned} |\lambda - \lambda_L| &\leq |\alpha_{\mathbf{a}'}[\Phi](\tau_{2-}) - \alpha_{\mathbf{a}'}[\Phi_0](\tau_{2-})| + |\alpha_{\mathbf{a}'}[\Phi_0](\tau_{2-}) - \alpha_{\mathbf{a}'}[\Phi_0](\tau_{1-})| \\ &\quad + |\alpha_{\mathbf{a}'}[\Phi_0](\tau_{1-}) + (1 - \alpha_{\mathbf{a}, \mathbf{a}'}[\Phi_0](\tau_{1-}))| \\ &\leq |\alpha_{\mathbf{a}'}[\Phi](\tau_{2-}) - \alpha_{\mathbf{a}'}[\Phi_0](\tau_{2-})| \\ &\quad + (|\alpha_{\mathbf{a}}[\Phi_0](\tau_{2-}) - 1| + |\alpha_{\mathbf{a}}[\Phi_0](\tau_{1-}) - 1|) + |1 - \alpha_{\mathbf{a}}[\Phi_0](\tau_{1-})| \leq 2\mathbf{M} \end{aligned}$$

See (3.27). Require $(\ell_1, \dots, \ell_7) \geq (32, 1, 0, 2, 1, 0, -2^{-7}) (\bullet)_9$. Then $4\mathbf{M}(1+w) \leq \mathbf{K} \leq 1$ and $|\lambda - \lambda_L| \leq \frac{1}{2}(1+w)^{-1}\mathbf{K}$. In particular $\lambda \geq \lambda_L - (1+w)^{-1} \geq 1$.

We now construct the components of $\mathbf{f}' = (\mathbf{h}', w', q')$.

Construction of w' . Require $(\ell_1, \dots, \ell_7) \geq (32, 1, 0, 2, 1, 0, -2^{-7}) (\bullet)_{10}$ and set

$$w' = \frac{\alpha_{\mathbf{a}}[\Phi](\tau_{2-})}{-\alpha_{\mathbf{a}, \mathbf{a}'}[\Phi](\tau_{2-})} > 0 \quad (3.32)$$

To check that the denominator is nonzero and that $w' > 0$, note that for all $\tau \in \mathcal{J}$:

$$\begin{aligned} |\alpha_{\mathbf{a}, \mathbf{a}'}[\Phi](\tau) - \alpha_{\mathbf{a}, \mathbf{a}'}[\Phi_0](\tau)| &\leq 2\mathbf{M} \\ |\alpha_{\mathbf{a}}[\Phi](\tau) - 1| &\leq |\alpha_{\mathbf{a}}[\Phi](\tau) - \alpha_{\mathbf{a}}[\Phi_0](\tau)| + |\alpha_{\mathbf{a}}[\Phi_0](\tau) - 1| \leq 2\mathbf{M} \end{aligned}$$

and $4\mathbf{M} \leq \mathbf{X}(-1, 0, 0, -1, 0, 0, 0) \leq \frac{1}{1+w} \leq |\alpha_{\mathbf{a}, \mathbf{a}'}[\Phi_0](\tau)|$ and $4\mathbf{M} \leq 1$. Hence,

$$|\alpha_{\mathbf{a}, \mathbf{a}'}[\Phi](\tau) - \alpha_{\mathbf{a}, \mathbf{a}'}[\Phi_0](\tau)| \leq \frac{1}{2}|\alpha_{\mathbf{a}, \mathbf{a}'}[\Phi_0](\tau)| \quad |\alpha_{\mathbf{a}}[\Phi](\tau) - 1| \leq \frac{1}{2}$$

In particular, $\alpha_{\mathbf{a}, \mathbf{a}'}[\Phi](\tau_{2-}) \leq \frac{1}{2}\alpha_{\mathbf{a}, \mathbf{a}'}[\Phi_0](\tau_{2-}) < 0$ and $\alpha_{\mathbf{a}}[\Phi](\tau_{2-}) > 0$. Consequently, w' is well defined and positive. Recall (3.14b) and estimate

$$\begin{aligned} |w' - w_L| &\leq \left| \frac{\alpha_{\mathbf{a}}[\Phi](\tau_{2-})}{\alpha_{\mathbf{a}, \mathbf{a}'}[\Phi](\tau_{2-})} - \frac{\alpha_{\mathbf{a}}[\Phi_0](\tau_{2-})}{\alpha_{\mathbf{a}, \mathbf{a}'}[\Phi](\tau_{2-})} \right| + \left| \frac{\alpha_{\mathbf{a}}[\Phi_0](\tau_{2-})}{\alpha_{\mathbf{a}, \mathbf{a}'}[\Phi](\tau_{2-})} - \frac{\alpha_{\mathbf{a}}[\Phi_0](\tau_{2-})}{\alpha_{\mathbf{a}, \mathbf{a}'}[\Phi_0](\tau_{2-})} \right| \\ &\quad + \left| \frac{\alpha_{\mathbf{a}}[\Phi_0](\tau_{2-})}{\alpha_{\mathbf{a}, \mathbf{a}'}[\Phi_0](\tau_{2-})} - \frac{1}{\alpha_{\mathbf{a}, \mathbf{a}'}[\Phi_0](\tau_{2-})} \right| \\ &\leq 2w_L\mathbf{M} + 4w_L^2\mathbf{M} + w_L\mathbf{M} \leq 2^3(1+w)^2\mathbf{M} \leq \frac{1}{2}\mathbf{X}(6, 0, 0, 2, 0, 0, 0)\mathbf{M} \end{aligned}$$

We require $(\ell_1, \dots, \ell_7) \geq (35, 1, 0, 3, 1, 0, -2^{-7})$ $(\bullet)_{11}$. Hence $|w' - w_L| \leq \frac{1}{2}\mathbf{K} \leq \frac{1}{2}$.
Construction of $\mathbf{h}'$. Let λ and w' be given by (3.31) and (3.32). Set

$$\mu = \frac{1+2w'}{1+w'} \left(-\xi_{\mathbf{a}}[\Phi, \mathbf{h}](\tau_{2-}) + \mathbf{h} \log \lambda \right) - \mathbf{h} \log 2$$

Recall (3.14c) and estimate

$$\begin{aligned} \left| \mu - \frac{\mathbf{h}}{\mathbf{h}_L} \right| &\leq \left| \frac{1+2w'}{1+w'} \right| \left| -\xi_{\mathbf{a}}[\Phi, \mathbf{h}](\tau_{2-}) + \mathbf{h} \log \lambda + \tau_{1-} - \mathbf{h} \log \lambda_L \right| \\ &\quad + \left| \frac{1+2w'}{1+w'} - \frac{1+2w_L}{1+w_L} \right| \left| \tau_{1-} - \mathbf{h} \log \lambda_L \right| \\ &\leq 2 \left| \tau_{1-} - \tau_{2-} \right| + 2 \left| \tau_{2-} - \xi_{\mathbf{a}}[\Phi_0, \mathbf{h}](\tau_{2-}) \right| \\ &\quad + 2 \left| \xi_{\mathbf{a}}[\Phi_0, \mathbf{h}](\tau_{2-}) - \xi_{\mathbf{a}}[\Phi, \mathbf{h}](\tau_{2-}) \right| + 4\mathbf{h} |\lambda - \lambda_L| + 4 \frac{|w' - w_L|}{(1+w_L)^2} \\ &\leq 2^2 \mathbf{M} + \mathbf{M} + 2\mathbf{M} + 2^3 \mathbf{M} + 2^4 \mathbf{M} \leq 2^5 \mathbf{M} \end{aligned}$$

For the second inequality, use $(1+2w') \leq 2(1+w')$ and $\lambda, \lambda_L \geq \frac{1}{2}$ and $|\tau_{1-}| \leq 1$ and $|\mathbf{h} \log \lambda_L| \leq |\tau_* \log \lambda_L| \leq 1$, see (3.19), and $1+w' \geq \frac{1}{2}(1+w_L)$. By inspection,

$$\frac{\mathbf{h}}{\mathbf{h}_L} \geq \frac{1+w}{2+w} \min\{1, q\} \geq \mathbf{X}(-1, 0, 0, 0, -1, 0, 0)$$

To make sure that $\mu > 0$, we require $(\ell_1, \dots, \ell_7) \geq (36, 1, 0, 1, 2, 0, -2^{-7})$ $(\bullet)_{12}$, so that $2^5 \mathbf{M} \leq \frac{1}{2} \mathbf{X}(-1, 0, 0, 0, -1, 0, 0) \mathbf{K} \leq \frac{1}{2} \frac{\mathbf{h}}{\mathbf{h}_L}$, that is $|\mu - \frac{\mathbf{h}}{\mathbf{h}_L}| \leq \frac{1}{2} \frac{\mathbf{h}}{\mathbf{h}_L}$ and $\mu > 0$. Set

$$\mathbf{h}' = \mathbf{h}/\mu > 0 \quad (3.33)$$

Require $\ell_8 \geq 7$ $(\bullet)_{13}$, so that $\mathbf{h} \leq \mathbf{X}(-7, 0, 0, 0, -1, 0, 0)$ and

$$|\mathbf{h}' - \mathbf{h}_L| = \mathbf{h} \frac{\mathbf{h}/\mathbf{h}_L}{\mu} \left(\frac{\mathbf{h}_L}{\mathbf{h}} \right)^2 \left| \mu - \frac{\mathbf{h}}{\mathbf{h}_L} \right| \leq \frac{1}{2} \mathbf{X}(2, 0, 0, 0, 1, 0, 0) \mathbf{M}$$

We require $(\ell_1, \dots, \ell_7) \geq (31, 1, 0, 1, 2, 0, -2^{-7})$ $(\bullet)_{14}$. Then $|\mathbf{h}' - \mathbf{h}_L| \leq \frac{1}{2} \mathbf{K} \leq \frac{1}{2}$.
Construction of q' . Set

$$q' = \frac{1}{1+w'} \left(\mathbf{h}' \log \lambda - \frac{\mathbf{h}'}{\mathbf{h}} \xi_{\mathbf{c}'}[\Phi, \mathbf{h}](\tau_{2-}) - \frac{w'(1+w')}{1+2w'} - \frac{1+3w'+(w')^2}{1+2w'} \mathbf{h}' \log 2 \right) \quad (3.34)$$

Recall (3.14d) and estimate

$$\begin{aligned} |q' - q_L| &\leq \left| \frac{1}{1+w'} \mathbf{h}' \log \lambda - \frac{1}{1+w_L} \mathbf{h}_L \log \lambda_L \right| \\ &\quad + \left| \frac{1}{1+w'} \frac{\mathbf{h}'}{\mathbf{h}} \xi_{\mathbf{c}'}[\Phi, \mathbf{h}](\tau_{2-}) - \frac{1}{1+w_L} \frac{\mathbf{h}_L}{\mathbf{h}} \left(\xi_{\mathbf{a}, \mathbf{c}'}[\Phi_0, \mathbf{h}](\tau_{1-}) - \tau_{1-} \right) \right| \\ &\quad + \left| \frac{w'}{1+2w'} - \frac{w_L}{1+2w_L} \right| + \left| \frac{1+3w'+(w')^2}{(1+w')(1+2w')} \mathbf{h}' - \frac{1+3w_L+(w_L)^2}{(1+w_L)(1+2w_L)} \mathbf{h}_L \right| \\ &\leq \frac{1}{1+w'} \mathbf{h}' \left| \log \lambda - \log \lambda_L \right| + \frac{1}{1+w'} |\mathbf{h}' - \mathbf{h}_L| \log \lambda_L + \left| \frac{1}{1+w'} - \frac{1}{1+w_L} \right| \mathbf{h}_L \log \lambda_L \\ &\quad + \frac{1}{1+w'} \frac{\mathbf{h}'}{\mathbf{h}} \left| \xi_{\mathbf{c}'}[\Phi, \mathbf{h}](\tau_{2-}) - \xi_{\mathbf{c}'}[\Phi_0, \mathbf{h}](\tau_{2-}) \right| + \frac{1}{1+w'} \left| \frac{\mathbf{h}'}{\mathbf{h}} - \frac{\mathbf{h}_L}{\mathbf{h}} \right| \left| \xi_{\mathbf{c}'}[\Phi_0, \mathbf{h}](\tau_{2-}) \right| \\ &\quad + \left| \frac{1}{1+w'} - \frac{1}{1+w_L} \right| \frac{\mathbf{h}_L}{\mathbf{h}} \left| \xi_{\mathbf{c}'}[\Phi_0, \mathbf{h}](\tau_{2-}) \right| \\ &\quad + \frac{1}{1+w_L} \frac{\mathbf{h}_L}{\mathbf{h}} \left| \xi_{\mathbf{c}'}[\Phi_0, \mathbf{h}](\tau_{2-}) - \xi_{\mathbf{a}, \mathbf{c}'}[\Phi_0, \mathbf{h}](\tau_{1-}) + \tau_{1-} \right| \\ &\quad + \left| \frac{w'}{1+2w'} - \frac{w_L}{1+2w_L} \right| + \left| \frac{1+3w'+(w')^2}{(1+w')(1+2w')} - \frac{1+3w_L+(w_L)^2}{(1+w_L)(1+2w_L)} \right| \mathbf{h}' + |\mathbf{h}' - \mathbf{h}_L| \end{aligned}$$

$$\begin{aligned}
&\leq 2^3 |\lambda - \lambda_L| + |\mathbf{h}' - \mathbf{h}_L| (1+w) + 4 \frac{|w' - w_L|}{(1+w_L)^2} (1+w) \\
&\quad + 2 \frac{\mathbf{h}_L}{\mathbf{h}} |\xi_{\mathbf{c}'}[\Phi, \mathbf{h}](\tau_{2-}) - \xi_{\mathbf{c}'}[\Phi_0, \mathbf{h}](\tau_{2-})| + \left| \frac{\mathbf{h}'}{\mathbf{h}} - \frac{\mathbf{h}_L}{\mathbf{h}} \right| |\xi_{\mathbf{c}'}[\Phi_0, \mathbf{h}](\tau_{2-})| \\
&\quad + 2 \frac{|w' - w_L|}{(1+w_L)^2} \frac{\mathbf{h}_L}{\mathbf{h}} |\xi_{\mathbf{c}'}[\Phi_0, \mathbf{h}](\tau_{2-})| + \frac{\mathbf{h}_L}{\mathbf{h}} |\xi_{\mathbf{a}, \mathbf{c}'}[\Phi_0, \mathbf{h}](\tau_{2-}) - \xi_{\mathbf{a}, \mathbf{c}'}[\Phi_0, \mathbf{h}](\tau_{1-})| \\
&\quad + \frac{\mathbf{h}_L}{\mathbf{h}} |\xi_{\mathbf{a}}[\Phi_0, \mathbf{h}](\tau_{2-}) - \tau_{2-}| + \frac{\mathbf{h}_L}{\mathbf{h}} |\tau_{2-} - \tau_{1-}| \\
&\quad + 2 \frac{|w' - w_L|}{(1+w_L)^2} + 2^3 \frac{|w' - w_L|}{(1+w_L)^2} + |\mathbf{h}' - \mathbf{h}_L| \\
&\leq 2^4 \mathbf{M} + \mathbf{X}(2, 0, 0, 1, 1, 0, 0) \mathbf{M} + \mathbf{X}(5, 0, 0, 1, 0, 0, 0) \mathbf{M} \\
&\quad + \mathbf{X}(2, 0, 0, 0, 1, 0, 0) \mathbf{M} + \mathbf{X}(6, 1, 0, 1, 1, 1, 0) \mathbf{M} \\
&\quad + \mathbf{X}(9, 0, 0, 1, 1, 1, 0) \mathbf{M} + \mathbf{X}(3, 0, 0, 1, 1, 0, 0) \mathbf{M} \\
&\quad + \mathbf{X}(1, 0, 0, 0, 1, 0, 0) \mathbf{M} + \mathbf{X}(2, 0, 0, 0, 1, 0, 0) \mathbf{M} \\
&\quad + 2^3 \mathbf{M} + 2^5 \mathbf{M} + \mathbf{X}(1, 0, 0, 0, 1, 0, 0) \mathbf{M} \\
&\leq \frac{1}{2} \mathbf{X}(11, 1, 0, 1, 1, 1, 0) \mathbf{M}
\end{aligned}$$

For the third inequality, use $\mathbf{h}_L \leq \mathbf{X}(1, 0, 0, 0, 1, 0, 0) \mathbf{h} \leq 2$ and $\mathbf{h}' = \mathbf{h}/\mu \leq 2\mathbf{h}_L \leq 2^2$ and $\lambda, \lambda_L \geq \frac{1}{2}$ and $|\mathbf{h} \log \lambda_L| \leq 1$ and $\log \lambda_L \leq 1 + w$ and $(1 + w') \geq \frac{1}{2}(1 + w_L)$.

For the fourth inequality, use $(1 + w) \leq \mathbf{X}(1, 0, 0, 1, 0, 0, 0)$ and $\frac{|w' - w_L|}{(1+w_L)^2} \leq 2^2 \mathbf{M}$ and

$$\begin{aligned}
|\xi_{\mathbf{c}'}[\Phi_0, \mathbf{h}](\tau_{2-})| &\leq |\xi_{\mathbf{a}, \mathbf{c}'}[\Phi_0, \mathbf{h}](\tau_{2-})| + |\xi_{\mathbf{a}}[\Phi_0, \mathbf{h}](\tau_{2-}) - \tau_{2-}| + |\tau_{2-}| \\
&\leq \mathbf{X}(3, 0, 0, 1, 0, 1, 0) + 1 + 1 \leq \mathbf{X}(5, 0, 0, 1, 0, 1, 0)
\end{aligned}$$

We require $(\ell_1, \dots, \ell_7) \geq (40, 2, 0, 2, 2, 1, -2^{-7}) (\bullet)_{15}$, such that $|q' - q_L| \leq \frac{1}{2} \mathbf{K} \leq \frac{1}{2}$. The maximum of $\alpha_{\mathbf{b}, \mathbf{c}}, \alpha_{\mathbf{c}, \mathbf{a}}, \alpha_{\mathbf{a}, \mathbf{b}}$. Require $(\ell_1, \dots, \ell_7) \geq (24, 1, 2, 2, 0, 0, -2^{-7}) (\bullet)_{16}$. Then $\mathbf{X}(20, 1, 0, 1, 0, 0, -2^{-7}) \leq 2^{-3}(1+w)^{-1} \min\{w^2, 1\}$. By the inequality (3.26), we have $d_{\mathcal{D}}(\Phi_0(\tau), \Phi(\tau)) \leq 2^{-3}(1+w)^{-1} \min\{w^2, 1\}$, for all $\tau \in [\tau_{0-}, \tau_{0+}]$. Hence,

$$\begin{aligned}
\alpha_{\mathbf{a}, \mathbf{p}}[\Phi] &\leq \alpha_{\mathbf{a}, \mathbf{p}}[\Phi_0] + d_{\mathcal{D}}(\Phi_0, \Phi) \leq -(1+w)^{-1} + d_{\mathcal{D}}(\Phi_0, \Phi) \leq -2^{-1}(1+w)^{-1} \\
\alpha_{\mathbf{b}, \mathbf{c}}[\Phi] &\leq \alpha_{\mathbf{a}, \mathbf{b}}[\Phi] + \alpha_{\mathbf{a}, \mathbf{c}}[\Phi] - 2\alpha_{\mathbf{a}}[\Phi] \leq \alpha_{\mathbf{a}, \mathbf{b}}[\Phi] + \alpha_{\mathbf{a}, \mathbf{c}}[\Phi] + 2A_{\mathbf{a}}[\Phi] \\
&\leq 2^2 d_{\mathcal{D}}(\Phi_0, \Phi) + \alpha_{\mathbf{a}, \mathbf{b}}[\Phi_0] + \alpha_{\mathbf{a}, \mathbf{c}}[\Phi_0] + 2A_{\mathbf{a}}[\Phi_0] \leq -2^{-1}(1+w)^{-1} w^2
\end{aligned}$$

for all $\tau \in [\tau_{0-}, \tau_{0+}]$ and all $\mathbf{p} \in \{\mathbf{b}, \mathbf{c}\}$.

Definition of the maps Π , Λ and τ_{2-} . Set $(\ell_1, \dots, \ell_7) = (40, 2, 2, 3, 2, 1, -2^{-7})$ and $\ell_8 = 7$. With this choice, all inequalities $(\bullet)$ hold. The constant $\mathbf{K}$ defined by (3.19) coincides with $\mathbf{K}(\mathbf{f})$, defined by (3.17). Furthermore, a vector $\mathbf{f} = (\mathbf{h}, w, q) \in (0, \infty)^3$ satisfies our basic assumption (3.19) if and only if $\mathbf{f} \in \mathcal{F}$. Therefore, we can set

$$\begin{aligned}
\Pi[\pi, \sigma_*] : \mathcal{F} &\rightarrow (0, \infty)^2 \times \mathbb{R} & \mathbf{f} &\mapsto \text{right hand sides of } ((3.33), (3.32), (3.34)) \\
\Lambda[\pi, \sigma_*] : \mathcal{F} &\rightarrow [1, \infty) & \mathbf{f} &\mapsto \text{right hand side of } (3.31) \\
\tau_{2-}[\pi, \sigma_*] : \mathcal{F} &\rightarrow (-\infty, 0) & \mathbf{f} &\mapsto \text{right hand side of } (3.30)
\end{aligned}$$

Properties (a), (b), (c) and (e) in Proposition 3.3 are by construction, where it is understood that the fixed point Φ of the map P , whose domain of definition is $[\tau_{0-}, \tau_{0+}]$, has to be restricted to the subinterval $[\tau_{2-}, \tau_{2+}]$ to comply with the statement in Proposition 3.3 (e). The statements of (e.1), (e.3), (e.4), (e.5) have already been discussed in this proof. Equation $\Phi(\tau_{2+}) = \Phi_*$ in (e.2), with $\Phi_* = \Phi_*(\pi, \mathbf{f}, \sigma_*)$, follows from the fixed point equation $P(\Phi) = \Phi$, see (3.23), and from $\Phi_1(\tau_{1+}) = \Phi_*$ and $\tau_{0+} = \tau_{1+} = \tau_{2+}$.

Equation $\Phi(\tau_{2-}) = \lambda \Phi_*(\pi', \mathbf{f}', \sigma_*)$ in (e.2) with $\mathbf{f}' = (\mathbf{h}', w', q')$ is equivalent to (recall $\mathbf{b}' = \mathbf{a}$)

$$\alpha_{\mathbf{a}'}[\Phi](\tau_{2-}) = -\lambda \quad (3.35a)$$

$$\alpha_{\mathbf{a}}[\Phi](\tau_{2-}) = \lambda \frac{w'}{1+w'} \quad (3.35b)$$

$$\alpha_{\mathbf{c}'}[\Phi](\tau_{2-}) = \lambda(-w' - \mu') \quad (3.35c)$$

$$\frac{1}{\mathbf{h}} \xi_{\mathbf{a}'}[\Phi, \mathbf{h}](\tau_{2-}) = \log \lambda + \frac{1}{\mathbf{h}'} \left\{ -\frac{1+w'}{1+2w'}(1 + \mathbf{h}' \log 2) \right\} \quad (3.35d)$$

$$\frac{1}{\mathbf{h}} \xi_{\mathbf{a}}[\Phi, \mathbf{h}](\tau_{2-}) = \log \lambda + \frac{1}{\mathbf{h}'} \left\{ -\frac{1+w'}{1+2w'}(1 + \mathbf{h}' \log 2) \right\} \quad (3.35e)$$

$$\frac{1}{\mathbf{h}} \xi_{\mathbf{c}'}[\Phi, \mathbf{h}](\tau_{2-}) = \log \lambda + \frac{1}{\mathbf{h}'} \left\{ -(1 + w')q' - \frac{w'(1+w')}{1+2w'} - \frac{1+3w'+(w')^2}{1+2w'} \mathbf{h}' \log 2 \right\} \quad (3.35f)$$

with $\mu' = (1 + w')(\beta_1^2 + \beta_2^2 + \beta_3^2 - 2\beta_2\beta_3 - 2\beta_3\beta_1 - 2\beta_1\beta_2)|_{\beta=\beta[\Phi_*(\pi', \mathbf{f}', \sigma_*)]}$. By inspection: (3.35a) follows from (3.31); (3.35b) follows from (3.31) and (3.32); (3.35e) follows from (3.33); (3.35f) follows from (3.34); (3.35d) follows from (3.35e) and the discussion following (3.30). These five equations and $c[\Phi, \mathbf{h}, Z](\tau_{2-}) = 0$ imply (3.35c). We have now checked (e.2). We now discuss (d).

Continuity of the maps Π , Λ and τ_{2-} . Fix $\mathbf{f}^\Psi = (\mathbf{h}^\Psi, w^\Psi, q^\Psi) \in \mathcal{F}$. Let $r > 0$ and let $\mathbf{f}^\Upsilon = (\mathbf{h}^\Upsilon, w^\Upsilon, q^\Upsilon) \in \mathcal{F}$ with $\|\mathbf{f}^\Psi - \mathbf{f}^\Upsilon\|_{\mathbb{R}^3} \leq r$. All the objects and abbreviations that have been introduced for a single element of $\mathcal{F}$ before, now come in two versions, one associated to each of $\mathbf{f}^B \in \mathcal{F}$ with $B = \Psi, \Upsilon$. By convention, these two versions are distinguished by a superscript B . For instance, $\tau_{0+}^B = \tau_{+1}^B = \tau_{2+}^B = \tau_{+1}(\mathbf{f}^B)$ and $\Phi_0^B = \Phi_0(\pi, \mathbf{f}^B, \sigma_*)|_{[\tau_{0-}^B, \tau_{0+}^B]}$ and $\mathcal{E}^B = \mathcal{E}(\sigma_*; \tau_{0-}^B, \tau_{0+}^B)$ and so forth. Following this convention, the contraction mapping fixed points are denoted $\Phi^B \in \mathcal{E}^B$. However, we also write $\Phi^\Psi = \Psi$ and $\Phi^\Upsilon = \Upsilon$. Suppose $r \leq \frac{1}{2}|q^\Psi - 1|$. Then

$$0 \neq \text{sgn}(q^\Psi - 1) = \text{sgn}(q^\Upsilon - 1) \quad (3.36)$$

Define $\chi : \mathbb{R} \rightarrow \mathbb{R}$ by $\chi(\tau) = \frac{\mathbf{h}^\Upsilon}{\mathbf{h}^\Psi}(\tau - \tau_{0+}^\Psi) + \tau_{0+}^\Upsilon$. Introduce four closed intervals $\mathcal{I}^B = [\tau_{0-}^B, \tau_{0+}^B]$, $B = \Psi, \Upsilon$, and $\mathcal{I}^\Xi = [\chi^{-1}(\tau_{0-}^\Upsilon), \tau_{0+}^\Psi]$ and $\mathcal{I} = \mathcal{I}^\Psi \cap \mathcal{I}^\Xi$. Observe that $\chi(\mathcal{I}^\Xi) = \mathcal{I}^\Upsilon$. By Proposition 3.1, the field $\Xi = \Upsilon \circ (\chi|_{\mathcal{I}^\Xi})$ satisfies $(\mathbf{a}, \mathbf{b}, c)[\Xi, \mathbf{h}^\Psi, Z] = 0$ on $\mathcal{I}^\Xi$. Recall $\mathcal{J}^B = [\tau_{0-}^B, \frac{1}{2}\tau_{1-}^B] \subset \mathcal{I}^B$ and $|\tau_{2-}^B - \tau_{1-}^B| \leq \frac{1}{2}\text{dist}_{\mathbb{R}}(\tau_{1-}^B, \mathbb{R} \setminus \mathcal{J}^B)$, see (3.28b) and (3.29). Set $\mathcal{J} = \mathcal{J}^\Psi \cap \mathcal{J}^\Xi \subset \mathcal{I}$ with $\mathcal{J}^\Xi = \chi^{-1}(\mathcal{J}^\Upsilon)$. If $r > 0$ is sufficiently small, then

$$\tau_{2-}^\Psi \in \mathcal{J} \quad \text{and} \quad \chi^{-1}(\tau_{2-}^\Upsilon) \in \mathcal{J}$$

These inclusions have similar proofs. We only verify $\tau_{2-}^\Psi \in \mathcal{J}$. We have $\tau_{2-}^\Psi \in \mathcal{J}^\Psi$ and

$$\begin{aligned} |\chi(\tau_{2-}^\Psi) - \tau_{1-}^\Upsilon| &\leq |\chi(\tau_{2-}^\Psi) - \chi(\tau_{1-}^\Psi)| + |\chi(\tau_{1-}^\Psi) - \tau_{1-}^\Upsilon| \\ &\leq \frac{\mathbf{h}^\Upsilon}{\mathbf{h}^\Psi} \frac{1}{2} \text{dist}_{\mathbb{R}}(\tau_{1-}^\Psi, \mathbb{R} \setminus \mathcal{J}^\Psi) + |\chi(\tau_{1-}^\Psi) - \tau_{1-}^\Upsilon| \end{aligned} \quad (3.37)$$

The right hand side of (3.37) is a continuous function of $\mathbf{f}^\Upsilon \in \mathcal{F}$ (with $\mathbf{f}^\Psi$ fixed) and is equal to $\frac{1}{2}\text{dist}_{\mathbb{R}}(\tau_{1-}^\Upsilon, \mathbb{R} \setminus \mathcal{J}^\Upsilon) > 0$ when $\mathbf{f}^\Upsilon = \mathbf{f}^\Psi$. Therefore (3.37) is $< \text{dist}_{\mathbb{R}}(\tau_{1-}^\Upsilon, \mathbb{R} \setminus \mathcal{J}^\Upsilon)$ if $r > 0$ is small enough. Hence $\chi(\tau_{2-}^\Psi) \in \mathcal{J}^\Upsilon$, that is $\tau_{2-}^\Psi \in \mathcal{J}^\Xi$. Set $\mathcal{D} = \mathcal{D}^\Psi = \mathcal{D}(\sigma_*)$ and $\mathcal{E} = \mathcal{E}(\sigma_*; \mathcal{I})$ and $\Phi_0 = \Phi_0(\pi, \mathbf{f}^\Psi, \sigma_*)|_{\mathcal{I}}$. Equivalently, $\Phi_0 = \Phi_0^\Psi|_{\mathcal{I}}$. Abbreviate $d_{\mathcal{X}} = d_{\mathcal{X}, (\pi, \mathbf{h}^\Psi)}$ for $\mathcal{X} = \mathcal{E}, \mathcal{D}$ and $d_{\mathcal{X}^B} = d_{\mathcal{X}^B, (\pi, \mathbf{h}^B)}$

for $B = \Psi, \Upsilon$. By (3.25), we have $d_{\mathcal{E}^B}(B, \Phi_0^B) \leq \frac{1}{2}\delta^B$ for $B = \Psi, \Upsilon$. If $r > 0$ is sufficiently small, then

$$d_{\mathcal{E}}(\Psi|_{\mathcal{I}}, \Phi_0) \leq \delta^\Psi \quad \text{and} \quad d_{\mathcal{E}}(\Xi|_{\mathcal{I}}, \Phi_0) \leq \delta^\Psi \quad (3.38)$$

The first follows from $d_{\mathcal{E}}(\Psi|_{\mathcal{I}}, \Phi_0) \leq d_{\mathcal{E}^\Psi}(\Psi, \Phi_0^\Psi) \leq \frac{1}{2}\delta^\Psi$. The second follows from

$$\begin{aligned} d_{\mathcal{E}}(\Xi|_{\mathcal{I}}, \Phi_0) &\leq d_{\mathcal{E}}(\Upsilon \circ \chi|_{\mathcal{I}}, \Phi_0^\Upsilon \circ \chi|_{\mathcal{I}}) + d_{\mathcal{E}}(\Phi_0^\Upsilon \circ \chi|_{\mathcal{I}}, \Phi_0) \\ &\leq \max\{1, \frac{\mathbf{h}^\Psi}{\mathbf{h}^\Upsilon}\} d_{\mathcal{E}^\Upsilon}(\Upsilon, \Phi_0^\Upsilon) + d_{\mathcal{E}}(\Phi_0^\Upsilon \circ \chi|_{\mathcal{I}}, \Phi_0) \\ &\leq \max\{1, \frac{\mathbf{h}^\Psi}{\mathbf{h}^\Upsilon}\} \frac{1}{2}\delta^\Upsilon + d_{\mathcal{E}}(\Phi_0^\Upsilon \circ \chi|_{\mathcal{I}}, \Phi_0) \end{aligned} \quad (3.39)$$

and because the right hand side of (3.39) is a continuous function of $\mathbf{f}^\Upsilon \in \mathcal{F}$ (with $\mathbf{f}^\Psi$ fixed), see (3.20), that is equal to $\frac{1}{2}\delta^\Psi$ when $\mathbf{f}^\Upsilon = \mathbf{f}^\Psi$.

Both $X = \Psi|_{\mathcal{I}}$ and $X = \Xi|_{\mathcal{I}}$ satisfy $(\mathbf{a}, \mathbf{b}, c)[X, \mathbf{h}^\Psi, Z] = 0$ on $\mathcal{I}$,

$$A_{\mathbf{a}}[X](\tau) = A_{\mathbf{a}}[X](\tau_{0+}^\Psi) + \int_{\tau_{0+}^\Psi}^\tau d\tau' \mathbf{I}_1[X, \mathbf{h}^\Psi, \pi](\tau') \quad (3.40a)$$

$$\theta_{\mathbf{a}}[X, \mathbf{h}^\Psi](\tau) = \theta_{\mathbf{a}}[X, \mathbf{h}^\Psi](\tau_{0+}^\Psi) + \int_{\tau_{0+}^\Psi}^\tau d\tau' \mathbf{I}_2[X, \mathbf{h}^\Psi, \pi](\tau') \quad (3.40b)$$

$$\alpha_{\mathbf{p}, \mathbf{a}}[X](\tau) = \alpha_{\mathbf{p}, \mathbf{a}}[X](\tau_{0+}^\Psi) + \int_{\tau_{0+}^\Psi}^\tau d\tau' \mathbf{I}_{(3, \mathbf{p})}[X, \mathbf{h}^\Psi, \pi](\tau') \quad (3.40c)$$

$$\begin{aligned} \xi_{\mathbf{p}, \mathbf{a}}[X, \mathbf{h}^\Psi](\tau) &= \xi_{\mathbf{p}, \mathbf{a}}[X, \mathbf{h}^\Psi](\tau_{0+}^\Psi) + \alpha_{\mathbf{p}, \mathbf{a}}[X](\tau_{0+}^\Psi) (\tau - \tau_{0+}^\Psi) \\ &\quad + \int_{\tau_{0+}^\Psi}^\tau d\tau'' \int_{\tau_{0+}^\Psi}^{\tau''} d\tau' \mathbf{I}_{(3, \mathbf{p})}[X, \mathbf{h}^\Psi, \pi](\tau') \end{aligned} \quad (3.40d)$$

for all $\mathbf{p} \in \{\mathbf{b}, \mathbf{c}\}$ and $\tau \in \mathcal{I}$. By (3.24b), (3.38), (3.40) and by $\sup_{\tau \in \mathcal{I}} |\tau - \tau_{0+}^\Psi| \leq 4$, we have $d_{\mathcal{E}}(\Psi|_{\mathcal{I}}, \Xi|_{\mathcal{I}}) \leq 2^3 d_{\mathcal{D}}(\Psi(\tau_{0+}^\Psi), \Xi(\tau_{0+}^\Psi)) + 2^{-1} d_{\mathcal{E}}(\Psi|_{\mathcal{I}}, \Xi|_{\mathcal{I}})$, and consequently

$$d_{\mathcal{E}}(\Psi|_{\mathcal{I}}, \Xi|_{\mathcal{I}}) \leq 2^4 d_{\mathcal{D}}(\Psi(\tau_{0+}^\Psi), \Xi(\tau_{0+}^\Psi)) = 2^4 d_{\mathcal{D}}(\Phi_\star(\pi, \mathbf{f}^\Psi, \sigma_\star), \Phi_\star(\pi, \mathbf{f}^\Upsilon, \sigma_\star))$$

In particular, $d_{\mathcal{E}}(\Psi|_{\mathcal{I}}, \Xi|_{\mathcal{I}}) \rightarrow 0$ as $\mathbf{f}^\Upsilon \rightarrow \mathbf{f}^\Psi$. Furthermore,

$$\begin{aligned} &d_{\mathcal{D}}(\lambda^\Psi \Phi_\star(\pi', \mathbf{f}'^\Psi, \sigma_\star), \lambda^\Upsilon \Phi_\star(\pi', \mathbf{f}'^\Upsilon, \sigma_\star)) \\ &= d_{\mathcal{D}}(\Psi(\tau_{2-}^\Psi), \Xi(\chi^{-1}(\tau_{2-}^\Upsilon))) \\ &\leq d_{\mathcal{D}}(\Psi(\tau_{2-}^\Psi), \Psi(\chi^{-1}(\tau_{2-}^\Upsilon))) + d_{\mathcal{D}}(\Psi(\chi^{-1}(\tau_{2-}^\Upsilon)), \Xi(\chi^{-1}(\tau_{2-}^\Upsilon))) \\ &\leq d_{\mathcal{D}}(\Psi(\tau_{2-}^\Psi), \Psi(\chi^{-1}(\tau_{2-}^\Upsilon))) + 2^4 d_{\mathcal{D}}(\Phi_\star(\pi, \mathbf{f}^\Psi, \sigma_\star), \Phi_\star(\pi, \mathbf{f}^\Upsilon, \sigma_\star)) \end{aligned}$$

By the last inequality, if we can show that $\chi^{-1}(\tau_{2-}^\Upsilon) \rightarrow \tau_{2-}^\Psi$ as $\mathbf{f}^\Upsilon \rightarrow \mathbf{f}^\Psi$, then $\tau_{2-}^\Upsilon \rightarrow \tau_{2-}^\Psi$ and $\lambda^\Upsilon \rightarrow \lambda^\Psi$ and $\mathbf{f}'^\Upsilon \rightarrow \mathbf{f}'^\Psi$. In other words, to show that Π , Λ and τ_{2-} are continuous, it suffices to show that $\chi^{-1}(\tau_{2-}^\Upsilon) \rightarrow \tau_{2-}^\Psi$ as $\mathbf{f}^\Upsilon \rightarrow \mathbf{f}^\Psi$.

By the discussion after (3.32), we have $\alpha_{\mathbf{a}}[\Psi](\tau) \geq \frac{1}{2}$ and $\alpha_{\mathbf{a}, \mathbf{a}'}[\Psi](\tau) \leq 0$ for all $\tau \in \mathcal{J} \subset \mathcal{J}^\Psi$. Hence, for all $\tau \in \mathcal{J}$,

$$\frac{d}{d\tau}(\xi_{\mathbf{a}}[\Psi, \mathbf{h}^\Psi] - \xi_{\mathbf{a}'}[\Psi, \mathbf{h}^\Psi]) = \alpha_{\mathbf{a}}[\Psi] - \alpha_{\mathbf{a}'}[\Psi] = 2\alpha_{\mathbf{a}}[\Psi] - \alpha_{\mathbf{a}, \mathbf{a}'}[\Psi] \geq 1$$

Hence, $|\tau - \tau_{2-}^\Psi| \leq |\xi_{\mathbf{a}}[\Psi, \mathbf{h}^\Psi](\tau) - \xi_{\mathbf{a}'}[\Psi, \mathbf{h}^\Psi](\tau)|$ if $\tau \in \mathcal{J}$. Set $\tau = \chi^{-1}(\tau_{2-}^\Upsilon) \in \mathcal{J}$:

$$\begin{aligned} |\chi^{-1}(\tau_{2-}^\Upsilon) - \tau_{2-}^\Psi| &\leq |\xi_{\mathbf{a}}[\Psi, \mathbf{h}^\Psi](\chi^{-1}(\tau_{2-}^\Upsilon)) - \xi_{\mathbf{a}'}[\Psi, \mathbf{h}^\Psi](\chi^{-1}(\tau_{2-}^\Upsilon))| \\ &\leq 2 d_{\mathcal{D}, \mathbf{h}^\Psi}(\Psi(\chi^{-1}(\tau_{2-}^\Upsilon)), \Xi(\chi^{-1}(\tau_{2-}^\Upsilon))) \end{aligned} \quad (3.41)$$

The last inequality follows from $\xi_{\mathbf{a}}[\Xi, \mathbf{h}^\Psi](\chi^{-1}(\tau_{2-}^\gamma)) = \xi_{\mathbf{a}'}[\Xi, \mathbf{h}^\Psi](\chi^{-1}(\tau_{2-}^\gamma))$ and the triangle inequality. Since $\mathbf{f}^\gamma \rightarrow \mathbf{f}^\Psi$ implies $d_{\mathcal{E}}(\Psi|_{\mathcal{I}}, \Xi|_{\mathcal{I}}) \rightarrow 0$, also the right hand side of (3.41) goes to zero, that is, $|\chi^{-1}(\tau_{2-}^\gamma) - \tau_{2-}^\Psi| \rightarrow 0$, as required.

Uniqueness of Π , Λ and τ_{2-} . Suppose we have two triples $\Pi_i, \Lambda_i, \tau_{2-,i}$ with $i = 1, 2$. Let $\mathbf{f} \in \mathcal{F}$ and let Φ_i be the corresponding fields in (e). By (e.1) and (e.2) and the local uniqueness for solutions to ODE's, we have $\Phi_1 = \Phi_2$ on the intersection of their domains of definition $[\max\{\tau_{2-,1}(\mathbf{f}), \tau_{2-,2}(\mathbf{f})\}, \tau_{2+}]$. Observe that $\tau_{-}(\mathbf{f}) < \tau_{2-,1}(\mathbf{f}), \tau_{2-,2}(\mathbf{f}) < \frac{1}{2}\tau_{1-}(\mathbf{f})$, by (c). By (e.3), we have $\tau_{2-,1}(\mathbf{f}) = \tau_{2-,2}(\mathbf{f})$. By (e.2), we have $\Pi_1(\mathbf{f}) = \Pi_2(\mathbf{f})$, $\Lambda_1(\mathbf{f}) = \Lambda_2(\mathbf{f})$. $\square$

Remark 3.6. In Proposition 3.3, the signature vector σ_* appears to play a passive role. However, observe that $\Phi_* = \Phi_*(\pi, \mathbf{f}, \sigma_*)$ in (e.2) depends on it in a crucial way, see Definition 3.12. For instance, while $\alpha_{\mathbf{a}}[\Phi_*]$ and $\alpha_{\mathbf{b}}[\Phi_*]$ do not depend on σ_* at all, and $\beta_i[\Phi_*]$, $i = 1, 2, 3$ only in a trivial way through their signs, the component $\alpha_{\mathbf{c}}[\Phi_*]$ does depend on σ_* in a more important way, because the right hand side of (3.10) does. That σ_* plays a role is not surprising, after all it distinguishes Bianchi VIII and IX.

4. The approximate epoch-to-epoch and era-to-era maps

This section is logically self-contained, and the notation is introduced from scratch. Its goal is to study two maps, denoted $\mathcal{Q}_R$ and $\mathcal{E}_R$, that we informally refer to (following [BKL1]) as the *epoch-to-epoch* and *era-to-era* maps. The two maps are related, the second is some iterate of the first. The subscript R is for *right* (as opposed to *left*). For the moment, the definition of $\mathcal{Q}_R$ is taken for granted without motivation. To understand its role, see Part 3 of Proposition 4.4 and its proof.

Definition 4.1 (Epoch-to-epoch map). *Set*

$$\mathcal{Q}_R : (0, \infty) \setminus \mathbb{Q} \rightarrow (0, \infty) \setminus \mathbb{Q}$$

$$w \mapsto \mathcal{Q}_R(w) = \begin{cases} \frac{1}{w} - 1 & \text{if } w < 1 \\ w - 1 & \text{if } w > 1 \end{cases}$$

For every $w \in (0, \infty) \setminus \mathbb{Q}$, set

$$\mathcal{Q}_R\{w\}(q, \mathbf{h}) = \left(\frac{\text{num1}}{\text{den}}, \frac{\text{num2}}{\text{den}} \right)$$

where, if $w < 1$,

$$\text{num1} = 1 + w + \mathbf{h} \log 2 - \mathbf{h}(1 + 2w) \log(1 + \frac{1}{w}) \quad (4.1a)$$

$$\text{num2} = \mathbf{h} \quad (4.1b)$$

$$\text{den} = (1 + w)(1 + q + \mathbf{h} \log 2) - \mathbf{h}(2 + w) \log(1 + \frac{1}{w}) \quad (4.1c)$$

and, if $w > 1$,

$$\text{num1} = (1 + w)(1 + q + \mathbf{h} \log 2) - \mathbf{h}(2 + w) \log(1 + \frac{1}{w}) \quad (4.2a)$$

$$\text{num2} = \mathbf{h}w \quad (4.2b)$$

$$\text{den} = 1 + w + \mathbf{h} \log 2 - \mathbf{h}(1 + 2w) \log(1 + \frac{1}{w}) \quad (4.2c)$$

Here, we regard $\mathcal{Q}_R\{w\}$ as a pair of rational functions over $\mathbb{R}$ of degree one in the pair of abstract variables $(q, \mathbf{h})$. Finally, for all $w \in (0, \infty) \setminus \mathbb{Q}$ and all integers $n \geq 0$, set

$$\mathcal{Q}_R^n(w) = \left(\underbrace{\mathcal{Q}_R \circ \cdots \circ \mathcal{Q}_R}_n \right)(w)$$

$$\mathcal{Q}_R^n\{w\} = \mathcal{Q}_R\{\mathcal{Q}_R^{n-1}(w)\} \circ \cdots \circ \mathcal{Q}_R\{\mathcal{Q}_R^2(w)\} \circ \mathcal{Q}_R\{\mathcal{Q}_R(w)\} \circ \mathcal{Q}_R\{w\}$$

Warning: $\mathcal{Q}_R^n\{w\}$ is not the n -fold composition of $\mathcal{Q}_R\{w\}$ with itself.

The goal of this section is to understand the bulk behavior of $\mathcal{Q}_R^n\{w\}$ for large $n \geq 0$.

Definition 4.2. The floor function is $\mathbb{R} \ni x \mapsto \lfloor x \rfloor = \max\{n \in \mathbb{Z} \mid n \leq x\}$.

Definition 4.3 (Era-to-era map). Define $\mathcal{E}_R : (0, 1) \setminus \mathbb{Q} \rightarrow (0, 1) \setminus \mathbb{Q}$ by $\mathcal{E}_R(w) = \mathcal{Q}_R^{\lfloor 1/w \rfloor}(w)$. For every $w \in (0, 1) \setminus \mathbb{Q}$, denote by $\mathcal{E}_R\{w\}$ the pair of rational functions over $\mathbb{R}$ given by $\mathcal{E}_R\{w\} = \mathcal{Q}_R^{\lfloor 1/w \rfloor}\{w\}$. Finally, for all $w \in (0, 1) \setminus \mathbb{Q}$ and all integers $n \geq 0$, set

$$\mathcal{E}_R^n(w) = \left(\underbrace{\mathcal{E}_R \circ \cdots \circ \mathcal{E}_R}_n \right)(w)$$

$$\mathcal{E}_R^n\{w\} = \mathcal{E}_R\{\mathcal{E}_R^{n-1}(w)\} \circ \cdots \circ \mathcal{E}_R\{\mathcal{E}_R^2(w)\} \circ \mathcal{E}_R\{\mathcal{E}_R(w)\} \circ \mathcal{E}_R\{w\}$$

Lemma 4.1. For all integers $m, n \geq 0$,

- $\mathcal{Q}_R^{m+n}\{w\} = \mathcal{Q}_R^m\{\mathcal{Q}_R^n(w)\} \circ \mathcal{Q}_R^n\{w\}$ for $w \in (0, \infty) \setminus \mathbb{Q}$
- $\mathcal{E}_R^{m+n}\{w\} = \mathcal{E}_R^m\{\mathcal{E}_R^n(w)\} \circ \mathcal{E}_R^n\{w\}$ for $w \in (0, 1) \setminus \mathbb{Q}$

Proposition 4.1. Let $w \in (0, 1) \setminus \mathbb{Q}$. Then, for every integer $1 \leq r \leq \lfloor \frac{1}{w} \rfloor$,

$$\mathcal{Q}_R^r\{w\}(q, \mathbf{h}) = \left(\frac{\text{num}1_r}{\text{den}_r}, \frac{\text{num}2_r}{\text{den}_r} \right) \quad (4.3)$$

where

$$\text{num}1_r = (1 + w)(r + rq - q) + \mathbf{h} A_1(w, r) \quad (4.4a)$$

$$\text{num}2_r = \mathbf{h}(w + 1 - wr) \quad (4.4b)$$

$$\text{den}_r = (1 + w)(1 + q) + \mathbf{h} A_2(w, r) \quad (4.4c)$$

and where

$$A_1(w, r) = (2r - 1 + wr - w) \log 2 - (2r - 1 + wr + w) \log(1 + \frac{1}{w})$$

$$+ \sum_{k=1}^{r-1} (1 + 2k - 2k^2w - w) \log(1 + \frac{w}{1-kw})$$

$$+ r \sum_{k=1}^{r-1} ((2k - 1)w - 2) \log(1 + \frac{w}{1-kw}) \quad (4.5a)$$

$$A_2(w, r) = (1 + wr) \log 2 - (2 + w) \log(1 + \frac{1}{w})$$

$$+ \sum_{k=1}^{r-1} ((2k - 1)w - 2) \log(1 + \frac{w}{1-kw}) \quad (4.5b)$$

Furthermore, $\mathcal{E}_R(w) = \frac{1}{w} - \lfloor \frac{1}{w} \rfloor$, that is, $\mathcal{E}_R$ is the Gauss map, and

$$\mathcal{E}_R\{w\}(q, \mathbf{h}) = \left(\frac{\text{num}1_{\lfloor 1/w \rfloor}}{\text{den}_{\lfloor 1/w \rfloor}}, \frac{\text{num}2_{\lfloor 1/w \rfloor}}{\text{den}_{\lfloor 1/w \rfloor}} \right)$$

Remark 4.1. In equation (4.5), we have $0 \leq \frac{w}{1-kw} \leq 1$ for all $1 \leq k \leq r-1$.

Proof. Let $w \in (0, 1) \setminus \mathbb{Q}$. We show (4.3) by induction over $1 \leq r \leq \lfloor \frac{1}{w} \rfloor$. The $r = 1$ base case of the induction argument, $\mathcal{Q}_R\{w\}(q, \mathbf{h}) = (\text{num}1_1/\text{den}_1, \text{num}2_1/\text{den}_1)$, is by direct inspection, using (4.1). The induction step becomes the identity

$$\mathcal{Q}_R\{\mathcal{Q}_R^{r-1}(w)\}\left(\frac{\text{num}1_{r-1}}{\text{den}_{r-1}}, \frac{\text{num}2_{r-1}}{\text{den}_{r-1}}\right) = \left(\frac{\text{num}1_r}{\text{den}_r}, \frac{\text{num}2_r}{\text{den}_r}\right) \quad (4.6)$$

for all $2 \leq r \leq \lfloor \frac{1}{w} \rfloor$. To calculate $\mathcal{Q}_R\{\mathcal{Q}_R^{r-1}(w)\}(\cdot)$, use formulas (4.2), since $\mathcal{Q}_R^{r-1}(w) = \frac{1}{w} - r + 1 > 1$. Observe that (4.6) follows from the identities

$$\begin{aligned} \lambda \text{ num}1_r &= (1 + (\frac{1}{w} - r + 1))(\text{den}_{r-1} + \text{num}1_{r-1} + \text{num}2_{r-1} \log 2) \\ &\quad - \text{num}2_{r-1} (2 + (\frac{1}{w} - r + 1)) \log(1 + \frac{w}{1-(r-1)w}) \\ \lambda \text{ num}2_r &= \text{num}2_{r-1} (\frac{1}{w} - r + 1) \\ \lambda \text{ den}_r &= \text{den}_{r-1} (1 + (\frac{1}{w} - r + 1)) + \text{num}2_{r-1} \log 2 \\ &\quad - \text{num}2_{r-1} (1 + 2(\frac{1}{w} - r + 1)) \log(1 + \frac{w}{1-(r-1)w}) \end{aligned}$$

where $\lambda = 2 + \frac{1}{w} - r > 2$. To verify each of these identities, divide both sides by λ , and use $\text{num}2_{r-1} = \mathbf{h}w\lambda$, to obtain the equivalent identities

$$\begin{aligned} \text{num}1_r &= \text{den}_{r-1} + \text{num}1_{r-1} + \text{num}2_{r-1} \log 2 \\ &\quad - \mathbf{h}(3w + 1 - rw) \log(1 + \frac{w}{1-(r-1)w}) \\ \text{num}2_r &= \mathbf{h}(1 - rw + w) \\ \text{den}_r &= \text{den}_{r-1} + \mathbf{h}w \log 2 - \mathbf{h}(3w + 2 - 2rw) \log(1 + \frac{w}{1-(r-1)w}) \end{aligned}$$

The last three identities are verified directly. $\square$

The following lemma will be used later.

Lemma 4.2. For every $w \in (0, 1) \setminus \mathbb{Q}$ and every integer r with $1 \leq r \leq \lfloor \frac{1}{w} \rfloor$,

$$\begin{aligned} 0 &\leq A_1(w, r) - rA_2(w, r) + \log 2 \leq 6\frac{1}{w} \\ -8 \log(1 + \frac{1}{w}) &\leq A_2(w, r) \leq 0 \end{aligned}$$

Here, A_1 and A_2 are defined by (4.5).

Proof. Observe that $rw \leq 1$. Calculate

$$\begin{aligned} \{A_1(w, r) - rA_2(w, r) + \log 2\} w &= w^2(r-1) \log 2 + rw(1-rw) \log 2 \\ &\quad + (1-w) w \log(1 + \frac{1}{w}) + w \sum_{k=1}^{r-1} (2k(1-kw) + (1-w)) \log(1 + \frac{w}{1-kw}) \end{aligned}$$

By inspection, the right hand side is non-negative, and bounded by

$$\leq 3 + \frac{1}{r} \sum_{k=1}^{r-1} (2k(1-kw) + (1-w)) \frac{w}{1-kw} \leq 6$$

We have $A_2(w, r) \leq 0$, because $rw \leq 1$, the sum of the first two terms on the right hand side of (4.5b) is non-positive, and the third term is non-positive. Estimate

$$\begin{aligned} |A_2(w, r)| &\leq (2 + w) \log(1 + \frac{1}{w}) + 2 \sum_{k=1}^{r-1} \log(1 + \frac{w}{1-kw}) \\ &\leq 3 \log(1 + \frac{1}{w}) + 2 \sum_{k=1}^{r-1} \frac{w}{1-kw} \leq 3 \log(1 + \frac{1}{w}) + 2(1 + \int_0^{r-1} dk \frac{w}{1-kw}) \\ &\leq 3 \log(1 + \frac{1}{w}) + 2(1 - \log(1 - (r-1)w)) \leq 8 \log(1 + \frac{1}{w}) \end{aligned}$$

since $2 < 3 \log 2 \leq 3 \log(1 + \frac{1}{w})$ and $-\log(1 - rw + w) \leq -\log w = \log \frac{1}{w} \leq \log(1 + \frac{1}{w})$. $\square$

Proposition 4.2. *For every $w \in (0, 1) \setminus \mathbb{Q}$, every $p > 0$ and every integer $1 \leq r \leq \lfloor \frac{1}{w} \rfloor$, let (μ', ν') be the pair of rational functions over $\mathbb{R}$ in the pair of abstract variables (μ, ν) given implicitly by*

$$\left(p' + \frac{\mu'}{\nu'}, \frac{1 + w'}{\nu'}\right) = \mathcal{Q}_R^r\{w\} \left(p + \frac{\mu}{\nu}, \frac{1 + w}{\nu}\right)$$

where $w' = \mathcal{Q}_R^r(w) = \frac{1}{w} - r$ and $p' = r - p/(1 + p)$, that is $(p', 0) = \mathcal{Q}_R^r\{w\}(p, 0)$. Then μ' is actually a linear polynomial over $\mathbb{R}$ in μ , and ν' is actually a linear polynomial over $\mathbb{R}$ in the pair (μ, ν) . Explicitly

$$\begin{pmatrix} \mu' \\ \nu' \end{pmatrix} = \frac{1}{w} \begin{pmatrix} -\frac{1}{1+p} & 0 \\ 1 & 1+p \end{pmatrix} \begin{pmatrix} \mu \\ \nu \end{pmatrix} + \frac{1}{w} \begin{pmatrix} A_1(w, r) - p' A_2(w, r) \\ A_2(w, r) \end{pmatrix} \quad (4.7)$$

The first and second entries of the vector

$$\frac{1}{w} \begin{pmatrix} A_1(w, r) - p' A_2(w, r) \\ A_2(w, r) \end{pmatrix} \quad (4.8)$$

are bounded in absolute value by $\leq 2^4(\frac{1}{w})^2$ and $\leq 2^3 \frac{1}{w} \log(1 + \frac{1}{w})$, respectively.

Proof. Equation (4.7) follows from equation (4.3). To check the bounds, observe that

$$A_1(w, r) - p' A_2(w, r) = \left(A_1(w, r) - r A_2(w, r) + \log 2\right) - \log 2 + \frac{p}{1+p} A_2(w, r)$$

Now, use Lemma 4.2 and $\log 2 \leq \frac{1}{w}$ and $\log(1 + \frac{1}{w}) \leq \frac{1}{w}$. $\square$

Definition 4.4. *For every sequence of strictly positive integers $(k_n)_{n \geq 0}$, we denote the associated infinite continued fraction by*

$$\langle k_0, k_1, \dots \rangle = \frac{1}{k_0 + \frac{1}{k_1 + \dots}} \in \left(\frac{1}{k_0+1}, \frac{1}{k_0}\right) \setminus \mathbb{Q}$$

Every element of $(0, 1) \setminus \mathbb{Q}$ has a unique continued fraction expansion of this form.

We now show that when $\mathbf{h} = 0$, the era-to-era maps can be realized as a left-shift operator on two-sided sequences of positive integers.

Proposition 4.3. Fix any two-sided sequence $(k_n)_{n \in \mathbb{Z}}$ of strictly positive integers and define two-sided sequences $(p_n)_{n \in \mathbb{Z}}$ and $(w_n)_{n \in \mathbb{Z}}$ by

$$\frac{1}{1+p_n} = \langle k_n, k_{n-1}, k_{n-2}, \dots \rangle \quad w_n = \langle k_{n+1}, k_{n+2}, k_{n+3}, \dots \rangle \quad (4.9)$$

Then $w_{n+1} = \mathcal{E}_R(w_n)$ and $(p_{n+1}, 0) = \mathcal{E}_R\{w_n\}(p_n, 0)$ for all $n \in \mathbb{Z}$, and $\mathcal{E}_R^n(w_0) = w_n$ and $\mathcal{E}_R^n\{w_0\}(p_0, 0) = (p_n, 0)$ for all $n \geq 0$.

Proof. Use $\mathcal{E}_R(w) = \frac{1}{w} - \lfloor \frac{1}{w} \rfloor$ and $\mathcal{E}_R\{w\}(p, 0) = (\lfloor \frac{1}{w} \rfloor - 1 + \frac{1}{1+p}, 0)$. $\square$

Definition 4.5. Fix any two-sided sequence $(k_n)_{n \in \mathbb{Z}}$ of strictly positive integers and define $(p_n)_{n \in \mathbb{Z}}$ and $(w_n)_{n \in \mathbb{Z}}$ by (4.9). For every integer $n \geq 0$, let (μ_n, ν_n) be the pair of linear polynomials over $\mathbb{R}$ in the abstract variables (μ_0, ν_0) , with coefficients depending only on the fixed sequence $(k_n)_{n \in \mathbb{Z}}$, given implicitly by

$$\left(p_n + \frac{\mu_n}{\nu_n}, \frac{1 + w_n}{\nu_n}\right) = \mathcal{E}_R^n\{w_0\}\left(p_0 + \frac{\mu_0}{\nu_0}, \frac{1 + w_0}{\nu_0}\right) \quad (4.10a)$$

or by the equivalent recursive prescription

$$\left(p_{n+1} + \frac{\mu_{n+1}}{\nu_{n+1}}, \frac{1 + w_{n+1}}{\nu_{n+1}}\right) = \mathcal{E}_R\{w_n\}\left(p_n + \frac{\mu_n}{\nu_n}, \frac{1 + w_n}{\nu_n}\right) \quad (4.10b)$$

By Proposition 4.2, equation (4.10b) is $V_{n+1} = X_n V_n + Y_n$, where $V_n = (\mu_n, \nu_n)^T$ and

$$X_n = \frac{1}{w_n} \begin{pmatrix} -\frac{1}{1+p_n} & 0 \\ 1 & 1+p_n \end{pmatrix} \quad Y_n = \frac{1}{w_n} \begin{pmatrix} A_1(w_n) - p_{n+1}A_2(w_n) \\ A_2(w_n) \end{pmatrix}$$

Here, $A_1(w) = A_1(w, \lfloor \frac{1}{w} \rfloor)$ and $A_2(w) = A_2(w, \lfloor \frac{1}{w} \rfloor)$, see equations (4.5).

Example 4.1. We consider Definition 4.5 when $k_n = 1$ for all $n \in \mathbb{Z}$. Then $w_n = p_n = w$ for all $n \in \mathbb{Z}$, where $w = \frac{1}{2}(\sqrt{5} - 1) \in (0, 1) \setminus \mathbb{Q}$. We have $w^2 + w - 1 = 0$ and $\lfloor \frac{1}{w} \rfloor = 1$ and

$$X_n = \begin{pmatrix} -1 & 0 \\ 1+w & 2+w \end{pmatrix} \quad Y_n = \begin{pmatrix} -2\log(1+w) \\ (2+w)\log 2 - (6+4w)\log(1+w) \end{pmatrix}$$

for all $n \geq 0$. It follows that $\mu_{n+2} = \mu_n$ for all $n \geq 0$, that is, $\mu_{2n} = \mu_0$ and $\mu_{2n+1} = -\mu_0 - 2\log(1+w)$. There are unique $\lambda_1 = \lambda_1(\mu_0)$ and $\lambda_2 = \lambda_2(\mu_0)$, depending only on μ_0 , such that $\nu_{2n+2} - \lambda_1 = (2+w)^2(\nu_{2n} - \lambda_1)$ and $\nu_{2n+3} - \lambda_2 = (2+w)^2(\nu_{2n+1} - \lambda_2)$. That is, $\nu_{2n} = (2+w)^{2n}(\nu_0 - \lambda_1) + \lambda_1$ and $\nu_{2n+1} = (2+w)^{2n}(\nu_1 - \lambda_2) + \lambda_2$. Here, $\nu_1 = (2+w)\nu_0 + (1+w)\mu_0 + (2+w)\log 2 - (6+4w)\log(1+w)$.

Definition 4.6 (Propagator). Let $(p_n)_{n \in \mathbb{Z}}$, $(w_n)_{n \in \mathbb{Z}}$, $(X_n)_{n \geq 0}$ be as in Definition 4.5. Then for all integers $n \geq m \geq 0$, let $P_{n,m} = X_{n-1} \cdots X_m$. Explicitly,

$$P_{n,m} = \begin{pmatrix} a_{n-1} \cdots a_m & 0 \\ \sum_{\ell=m}^{n-1} x_\ell & c_{n-1} \cdots c_m \end{pmatrix}$$

where $x_\ell = c_{n-1} \cdots c_{\ell+1} b_\ell a_{\ell-1} \cdots a_m$ whenever $n-1 \geq \ell \geq m$, and for all $\ell \geq 0$,

$$X_\ell = \begin{pmatrix} a_\ell & 0 \\ b_\ell & c_\ell \end{pmatrix} \quad a_\ell = \frac{-1}{w_\ell(1+p_\ell)} \quad b_\ell = \frac{1}{w_\ell} \quad c_\ell = \frac{1+p_\ell}{w_\ell}$$

In this definition, a sequence of dots $\cdots$ indicates that indices increase towards the left, one by one. A product of the form $F_k \cdots F_j$ is equal to one if $k = j - 1$. In particular, $P_{n,n} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$.

Lemma 4.3. In the context of Definition 4.5, we have $V_n = P_{n,0}V_0 + \sum_{\ell=0}^{n-1} P_{n,\ell+1}Y_\ell$.

Lemma 4.4. Recall Definition 4.6. For all integers $n \geq m \geq 0$, we have

$$\frac{1}{2} \leq \frac{w_{n-1}}{w_{m-1}}(-1)^{m+n}a_{n-1} \cdots a_m \leq 2 \quad (4.11a)$$

$$(1 - \delta_{mn}) \frac{1}{4} \leq \frac{w_{n-1}}{w_{m-1}^2} (w_{n-2} \cdots w_{m-1})^2 \sum_{\ell=m}^{n-1} x_\ell \leq 2 \quad (4.11b)$$

$$\frac{1}{2} \leq \frac{w_{n-1}}{w_{m-1}} (w_{n-2} \cdots w_{m-1})^2 c_{n-1} \cdots c_m \leq 2 \quad (4.11c)$$

Moreover,

$$w_{n-2} \cdots w_{m-1} \leq |\rho_-|^{n-m-1} = \rho_+^{-n+m+1} \quad \text{when} \quad n \geq m \geq 0 \quad (4.12)$$

Here, $\rho_\pm = \frac{1}{2}(1 \pm \sqrt{5})$ are the roots of the polynomial $\rho^2 - \rho - 1$. Observe that $|\rho_-| < 1$. In this lemma, a sequence of dots $\cdots$ indicates that indices increase towards the left, one by one. A product of the form $F_k \cdots F_j$ is equal to one if $k = j - 1$.

Proof. In this proof, abbreviate $v_\ell = 1/(1 + p_{\ell+1}) = \langle k_{\ell+1}, k_\ell, k_{\ell-1} \dots \rangle$. We have

$$(-1)^{m+n}a_{n-1} \cdots a_m = \frac{v_{n-2} \cdots v_{m-1}}{w_{n-2} \cdots w_{m-1}} \cdot \frac{w_{m-1}}{w_{n-1}} \quad (4.13a)$$

$$x_m = c_{n-1} \cdots c_{m+1}b_m = \frac{w_{n-2} \cdots w_m}{v_{n-2} \cdots v_m} \cdot \frac{w_{m-1}^2}{w_{n-1}} \cdot \left(\frac{1}{w_{n-2} \cdots w_{m-1}} \right)^2 \quad (4.13b)$$

$$c_{n-1} \cdots c_m = \frac{w_{n-2} \cdots w_{m-1}}{v_{n-2} \cdots v_{m-1}} \cdot \frac{w_{m-1}}{w_{n-1}} \cdot \left(\frac{1}{w_{n-2} \cdots w_{m-1}} \right)^2 \quad (4.13c)$$

where $n \geq m$ in (4.13a) and (4.13c) and $n > m$ in (4.13b). Each right hand side is written as a product of positive quotients, whose first factor is contained in the closed interval $[\frac{1}{2}, 2]$, see Proposition A.1 (a) of Appendix A. This implies (4.11a) and (4.11c). If $n = m$, the sum in (4.11b) vanishes and the estimate is trivial. Suppose $n > m$. We have $\text{sgn } x_\ell = (-1)^{\ell+m}$. If, in addition, ℓ satisfies $n - 2 \geq \ell \geq m$, we have $|x_{\ell+1}|/|x_\ell| = v_\ell v_{\ell-1} = v_\ell(v_\ell^{-1} - \lfloor v_\ell^{-1} \rfloor) \leq \frac{1}{2}$, that is $\frac{1}{2}|x_\ell| \geq |x_{\ell+1}|$. Therefore, the alternating sum in (4.11b) is non-negative and bounded from above by its first summand $x_m > 0$ and from below by $x_m + x_{m+1} \geq x_m - |x_{m+1}| \geq \frac{1}{2}x_m$. Actually, x_{m+1} is only defined when $n \geq m + 2$, but $\frac{1}{2}x_m$ is a lower bound for all $n \geq m + 1$. Now, estimate x_m , which is the left hand side of (4.13b).

Inequality (4.12) is a consequence of Proposition A.1 (b). $\square$

Warning: In the next proposition, the sequences $(w_j)_{j \in \mathbb{Z}}$ and $(p_j)_{j \in \mathbb{Z}}$ do not have the property that w_j and $(1 + p_j)^{-1}$ always lie in $(0, 1) \setminus \mathbb{Q}$. Rather, they lie in $(0, \infty) \setminus \mathbb{Q}$. However, in the proof of Proposition 4.4, the auxiliary sequences $(w_n^*)_{n \in \mathbb{Z}}$ and $(p_n^*)_{n \in \mathbb{Z}}$ do have the property that w_n^* and $(1 + p_n^*)^{-1}$ always lie in $(0, 1) \setminus \mathbb{Q}$. The discussion beginning with Proposition 4.3 and ending just above will be applied to the auxiliary sequences.

Proposition 4.4. For all $w_0 \in (0, 1) \setminus \mathbb{Q}$ and $q_0 \in (0, \infty) \setminus \mathbb{Q}$, introduce

- a two sided sequence of strictly positive integers $(k_n)_{n \in \mathbb{Z}}$ by

$$(1 + q_0)^{-1} = \langle k_0, k_{-1}, k_{-2}, \dots \rangle \quad w_0 = \langle k_1, k_2, k_3, \dots \rangle$$

- (Era Pointer) $J : \mathbb{Z} \rightarrow \mathbb{Z}$ by $J(0) = 0$ and $J(n+1) = J(n) + k_{n+1}$
- (Era Counter) $N : \mathbb{Z} \rightarrow \mathbb{Z}$ by $N(0) = 0$ and $N(j+1) = N(j) + \chi_{J(\mathbb{Z})}(j)$, where $\chi_{J(\mathbb{Z})}$ is the characteristic function of the image $J(\mathbb{Z}) \subset \mathbb{Z}$; equivalently

$$N(j) = \min\{n \in \mathbb{Z} \mid J(n) \geq j\} \quad (4.14)$$

- sequences $(w_j)_{j \in \mathbb{Z}}$ and $(p_j)_{j \in \mathbb{Z}}$ by (observe that w_0 is defined consistently)

$$w_j = \langle k_{N(j)+1}, k_{N(j)+2}, \dots \rangle + J(N(j)) - j \quad (4.15a)$$

$$p_j = \langle k_{N(j)-1}, k_{N(j)-2}, \dots \rangle + k_{N(j)} + j - J(N(j)) - 1 \quad (4.15b)$$

Part 1. Then $p_0 = q_0$ and $w_j, p_j > 0$ and $\mathcal{Q}_R(w_j) = w_{j+1}$ and $\mathcal{Q}_R\{w_j\}(p_j, 0) = (p_{j+1}, 0)$ for all $j \in \mathbb{Z}$, and $\mathcal{Q}_R^j(w_0) = w_j$ and $\mathcal{Q}_R^j\{w_0\}(q_0, 0) = (p_j, 0)$ for all $j \geq 0$.

Part 2. Let $\rho_+ = \frac{1}{2}(1 + \sqrt{5})$ and set

$$\mathbf{C}(w_0, q_0) = \sup_{n \geq 0} (n+1) \rho_+^{-2n} k_n \max\{k_{n-1}, k_{n-2}\} \in [1, \infty]$$

Suppose $\mathbf{C}(w_0, q_0) < \infty$. Fix any $0 < \mathbf{h}_0 \leq 2^{-14}(\mathbf{C}(w_0, q_0))^{-1}$. Then, there are sequences $(q_j)_{j \geq 0}$, $(\mathbf{h}_j)_{j \geq 0}$ of real numbers such that for every $j \geq 0$, the denominator appearing in the pair of rational functions $\mathcal{Q}_R\{w_j\}$, given by (4.1c) or (4.2c), is strictly positive at $(q_j, \mathbf{h}_j)$, and

$$(q_{j+1}, \mathbf{h}_{j+1}) = \mathcal{Q}_R\{w_j\}(q_j, \mathbf{h}_j)$$

or $(q_j, \mathbf{h}_j) = \mathcal{Q}_R^j\{w_0\}(q_0, \mathbf{h}_0)$. For all $j \geq 0$,

- $0 < \mathbf{h}_j \leq 2^6 \mathbf{h}_0 \rho_+^{-2N(j)}$ and

$$\frac{1}{4} \leq \frac{\mathbf{h}_j}{\mathbf{h}_0} \frac{1+w_0}{1+w_j} \prod_{\ell=0}^{N(j)-1} \frac{1}{w_{J(\ell)} w_{J(\ell-1)}} \leq 4$$

- $q_j \in (0, \infty) \setminus \mathbb{Z}$ and $|q_j - p_j| \leq 2^{12} \mathbf{h}_0 N(j) \rho_+^{-2N(j)} k_{N(j)}$
- $q_j \in (0, 1)$ if and only if $p_j \in (0, 1)$ if and only if $j-1 \in J(\mathbb{Z})$
- $\max\{\frac{1}{w_j}, w_j, \frac{1}{q_j}, \frac{1}{|q_j-1|}, q_j\} \leq 2^4 \max\{k_{N(j)-2}, k_{N(j)-1}, k_{N(j)}, k_{N(j)+1}\}$

Part 3. Let the map $\mathcal{Q}_L : (0, \infty)^3 \rightarrow (0, \infty)^2 \times \mathbb{R}$ be given as in Definition 3.16. Then the sequences $(\mathbf{h}_j)_{j \geq 0}$, $(w_j)_{j \geq 0}$, $(q_j)_{j \geq 0}$ in Part 2 satisfy for all $j \geq 0$:

$$(\mathbf{h}_j, w_j, q_j) = \mathcal{Q}_L(\mathbf{h}_{j+1}, w_{j+1}, q_{j+1})$$

Example 4.2. In Part 1 of Proposition 4.4, suppose the continued fraction expansions begin as follows: $(1 + q_0)^{-1} = \langle 1, 2, \dots \rangle$ and $w_0 = \langle 3, 1, 2, 4, \dots \rangle$. Then,

j	-3	-2	-1	0	1	2	3	4	5	6	7	8	9	10
$\chi_{J(\mathbb{Z})}(j)$	1	0	1	1	0	0	1	1	0	1	0	0	0	1
$N(j)$	-2	-1	-1	0	1	1	1	2	3	3	4	4	4	4
$J(N(j))$	-3	-1	-1	0	3	3	3	4	6	6	10	10	10	10

Proof (of Proposition 4.4). Two basic properties of J and N are, for all $j \in \mathbb{Z}$:

- $N \circ J$ is the identity; consequently $J(N(j)) = j$ if and only if $j \in J(\mathbb{Z})$
- $J(N(j)) \geq j$ and $J(N(j) - 1) \leq j - 1$ by (4.14); consequently

$$j \leq J(N(j)) \leq k_{N(j)} + j - 1 \quad (4.16)$$

The second bullet implies $w_j > 0$ and $p_j > 0$, for all $j \in \mathbb{Z}$. The first bullet implies that $w_j \in (0, 1)$ if and only if $j \in J(\mathbb{Z})$. Therefore, we have

$$\mathcal{Q}_R(w_j) = \begin{cases} \frac{1}{w_j} - 1 & \text{if } j \in J(\mathbb{Z}) \\ w_j - 1 & \text{if } j \notin J(\mathbb{Z}) \end{cases} \quad \mathcal{Q}_R\{w_j\}(p_j, 0) = \begin{cases} (\frac{1}{1+p_j}, 0) & \text{if } j \in J(\mathbb{Z}) \\ (1 + p_j, 0) & \text{if } j \notin J(\mathbb{Z}) \end{cases}$$

In the case $j \notin J(\mathbb{Z})$, we have $N(j+1) = N(j)$, and therefore $w_{j+1} = w_j - 1$ and $p_{j+1} = p_j + 1$, as required. In the case $j \in J(\mathbb{Z})$, we have $N(j+1) = N(j) + 1$ and $J(N(j+1)) = J(N(j) + 1) = J(N(j)) + k_{N(j)+1} = j + k_{N(j)+1}$, which implies

$$w_{j+1} = \langle k_{N(j)+2}, k_{N(j)+3}, \dots \rangle + k_{N(j)+1} - 1 = \frac{1}{w_j} - 1$$

$$p_{j+1} = \langle k_{N(j)}, k_{N(j)-1}, \dots \rangle = \frac{1}{1+p_j}$$

as required. **Part 1** is checked.

To prove **Part 2**, we first construct two sequences $(q_j)_{j \geq 0}$ and $(h_j)_{j \geq 0}$. Then we verify that they have the desired properties. Below, a sequence of dots $\dots$ in any product of the form $F_m \cdots F_n$ indicates that indices increase towards the left, one by one. The product is equal to one if $m = n - 1$. Define sequences $(w_n^*)_{n \in \mathbb{Z}}$ and $(p_n^*)_{n \in \mathbb{Z}}$ by $w_n^* = w_{J(n)} \in (0, 1) \setminus \mathbb{Q}$ and $p_n^* = p_{J(n)} \in (0, \infty) \setminus \mathbb{Q}$. Equivalently,

$$\frac{1}{1+p_n^*} = \langle k_n, k_{n-1}, k_{n-2}, \dots \rangle \quad w_n^* = \langle k_{n+1}, k_{n+2}, k_{n+3}, \dots \rangle$$

so that $w_{n+1}^* = \mathcal{E}_R(w_n^*)$ and $(p_{n+1}^*, 0) = \mathcal{E}_R\{w_n^*\}(p_n^*, 0)$, by Proposition 4.3. Let $(V_n^*)_{n \geq 0}$, with $V_n^* = (\mu_n^*, \nu_n^*)^T$, as in Definition 4.5, be the solution to $V_{n+1}^* = X_n^* V_n^* + Y_n^*$ for all $n \geq 0$ with $\mu_0^* = 0$ and $\nu_0^* = (1 + w_0^*)/\mathbf{h}_0 > 0$, where

$$X_n^* = \frac{1}{w_n^*} \begin{pmatrix} -\frac{1}{1+p_n^*} & 0 \\ 1 & 1+p_n^* \end{pmatrix} \quad Y_n^* = \frac{1}{w_n^*} \begin{pmatrix} A_1(w_n^*) - p_{n+1}^* A_2(w_n^*) \\ A_2(w_n^*) \end{pmatrix}$$

Let $(V_j)_{j \geq 0}$, with $V_j = (\mu_j, \nu_j)^T$, be given by $V_0 = V_0^*$ and for all $j \geq 1$ by $V_j = X_{N(j)-1}^* V_{N(j)-1}^* + Y_j$, where

$$Y_j = \frac{1}{w_s^*} \begin{pmatrix} A_1(w_s^*, j - J(s)) - p_j A_2(w_s^*, j - J(s)) \\ A_2(w_s^*, j - J(s)) \end{pmatrix} \Big|_{s=N(j)-1}$$

The functions A_1 and A_2 , on the right hand side, are well defined at $(w_s, j - J(s))$, where $s = N(j) - 1$, because $1 \leq j - J(s) \leq k_{N(j)} = \lfloor 1/w_s^* \rfloor$. The following two observations will be used later on:

- Recall (4.15b). For all $j \geq 1$, $s = N(j) - 1$, we have $p_j = (j - J(s)) - p_s^*/(1 + p_s^*)$, and consequently the estimates after (4.8) apply to Y_j , $j \geq 1$. They also apply to Y_n^* , $n \geq 0$, because $p_{n+1}^* = \lfloor 1/w_n^* \rfloor - p_n^*/(1 + p_n^*)$.
- $Y_{J(n)} = Y_{n-1}^*$ for all $n \geq 1$, and consequently $V_{J(n)} = V_n^*$. The last identity is also true when $n = 0$, because $J(0) = 0$.

As in Definition 4.6, set $P_{n,m}^* = X_{n-1}^* \cdots X_m^*$ for all $n \geq m \geq 0$. For all $j \geq 1$, $s = N(j) - 1$, Lemma 4.3 implies

$$V_j = X_s^* (P_{s,0}^* V_0^* + \sum_{\ell=0}^{s-1} P_{s,\ell+1}^* Y_\ell^*) + Y_j = P_{s+1,0}^* V_0^* + \sum_{\ell=0}^{s-1} P_{s+1,\ell+1}^* Y_\ell^* + Y_j$$

The last equation, the estimates after (4.8), and the estimates in Lemma 4.4 imply

$$|\mu_j| \leq \frac{2^5}{w_s^*} \sum_{\ell=0}^s \frac{1}{w_\ell^*} \quad (4.17a)$$

$$\nu_j \geq \frac{1}{2w_s^*} \left(\frac{1}{w_{s-1}^* \cdots w_{-1}^*} \right)^2 \left(w_{-1}^* \nu_0 - 2^8 \sum_{\ell=0}^s (w_{\ell-1}^* \cdots w_{-1}^*)^2 \log \left(1 + \frac{1}{w_\ell^*} \right) \right) \quad (4.17b)$$

$$\nu_j \leq \frac{2}{w_s^*} \left(\frac{1}{w_{s-1}^* \cdots w_{-1}^*} \right)^2 \left(w_{-1}^* \nu_0 + 2^6 \sum_{\ell=0}^s (w_{\ell-1}^* \cdots w_{-1}^*)^2 \log \left(1 + \frac{1}{w_\ell^*} \right) \right) \quad (4.17c)$$

for all $j \geq 1$ and $s = N(j) - 1$. All three estimates are also true when $j = 0$, $s = -1$. Abbreviate $\mathbf{C} = \mathbf{C}(w_0, q_0) \geq 1$. We have $k_n \leq \mathbf{C} \rho_+^{2n}$ for all $n \geq 0$. Estimate

$$\begin{aligned} & 2^8 \sum_{\ell=0}^s (w_{\ell-1}^* \cdots w_{-1}^*)^2 \log(1 + 1/w_\ell^*) \\ & \leq 2^8 w_{-1}^* \sum_{\ell=0}^\infty (\rho_+)^{-2\ell+2} \log(2 + k_{\ell+1}) \quad \text{see inequality (4.12)} \\ & \leq 2^8 w_{-1}^* \sum_{\ell=0}^\infty (\rho_+)^{-2\ell+2} (2 + \log k_{\ell+1}) \\ & \leq 2^{13} w_{-1}^* (1 + \log \mathbf{C}) \leq 2^{13} w_{-1}^* \mathbf{C} \leq 2^{-1} w_{-1}^* \frac{1}{\mathbf{h}_0} \leq 2^{-1} w_{-1}^* \nu_0 \end{aligned}$$

Hence, for all $j \geq 0$,

$$\frac{1}{4} \leq \frac{w_{N(j)-1}^*}{w_{-1}^*} (w_{N(j)-2}^* \cdots w_{-1}^*)^2 \frac{\mathbf{h}_0}{1 + w_0^*} \nu_j \leq 4 \quad (4.18)$$

Define sequences $(\mathbf{h}_j)_{j \geq 0}$ and $(q_j)_{j \geq 0}$ by $\mathbf{h}_j = (1 + w_j)/\nu_j > 0$ and $q_j = p_j + \mu_j/\nu_j$. These definitions are consistent when $j = 0$. Observe that $1 + w_j \leq 2 + J(N(j)) - j \leq 1 + k_{N(j)} \leq 2/w_{N(j)-1}^*$. Therefore, the estimates (4.12), (4.17), (4.18) imply for $j \geq 1$:

$$\frac{1}{4} \leq \frac{\mathbf{h}_j}{\mathbf{H}_j} \leq 4 \quad \text{where} \quad \mathbf{H}_j = \mathbf{h}_0 \frac{1 + w_j}{1 + w_0^*} \prod_{\ell=0}^{N(j)-1} (w_\ell^* w_{\ell-1}^*) \quad (4.19a)$$

$$\mathbf{H}_j \leq 2 \mathbf{h}_0 \left(\prod_{\ell=0}^{N(j)-2} w_\ell^* \right) \left(\prod_{\ell=-1}^{N(j)-2} w_\ell^* \right) \leq 2^4 \mathbf{h}_0 \rho_+^{-2N(j)} \quad (4.19b)$$

$$|q_j - p_j| \leq \frac{2^7 \mathbf{h}_0}{w_{N(j)-1}^*} \left(\sum_{\ell=0}^{N(j)-1} \frac{1}{w_\ell^*} \right) \prod_{\ell=0}^{N(j)-1} (w_\ell^* w_{\ell-1}^*) \leq 2^{12} \mathbf{h}_0 N(j) \rho_+^{-2N(j)} k_{N(j)}$$

The left hand sides are also less than or equal to the right hand sides when $j = 0$. Using (4.15b), one estimates

$$\begin{aligned} \text{dist}_{\mathbb{R}}(p_j, \mathbb{Z}) &= \text{dist}_{\mathbb{R}}(\langle k_{N(j)-1}, k_{N(j)-2}, \dots \rangle, \{0, 1\}) \\ &\geq \min \left\{ \frac{1}{k_{N(j)-1} + 1}, \frac{1}{k_{N(j)-2} + 2} \right\} \geq \frac{1}{3 \max\{k_{N(j)-1}, k_{N(j)-2}\}} \end{aligned}$$

By the definition of $\mathbf{C}$ and by the assumption $\mathbf{h}_0 \leq 2^{-14}\mathbf{C}^{-1}$, we have $|q_j - p_j| \leq \frac{3}{4}\text{dist}_{\mathbb{R}}(p_j, \mathbb{Z}) < \text{dist}_{\mathbb{R}}(p_j, \mathbb{Z})$ for all $j \geq 0$. Therefore, $q_j \in (0, \infty) \setminus \mathbb{Z}$. Moreover, $q_j \in (0, 1)$ iff $p_j \in (0, 1)$ iff $k_{N(j)} + j - J(N(j)) - 1 = 0$ iff $J(N(j) - 1) = j - 1$ iff $N(j) - 1 = N(j - 1)$ iff $j - 1 \in J(\mathbb{Z})$. For every $j \geq 0$,

$$\begin{aligned} w_j &\leq J(N(j)) - j + 1 \leq k_{N(j)} \\ 1/w_j &\leq k_{N(j)+1} + 1 \\ q_j &\leq p_j + 1 \leq k_{N(j)} + j - J(N(j)) + 1 \leq k_{N(j)} + 1 \\ (\text{dist}_{\mathbb{R}}(q_j, \mathbb{Z}))^{-1} &\leq 4 (\text{dist}_{\mathbb{R}}(p_j, \mathbb{Z}))^{-1} \leq 12 \max\{k_{N(j)-1}, k_{N(j)-2}\} \end{aligned}$$

Finally, we show that for all $j \geq 0$,

- (a) the denominator of $\mathcal{Q}_R\{w_j\}$, given by (4.1c) or (4.2c), is strictly positive at $(q_j, \mathbf{h}_j)$
- (b) $(q_{j+1}, \mathbf{h}_{j+1}) = \mathcal{Q}_R\{w_j\}(q_j, \mathbf{h}_j)$

For all $j \geq 0$, we have

$$2\mathbf{h}_j k_{N(j)+1} \leq 2(2^6 \mathbf{h}_0 \rho_+^{-2N(j)}) (\mathbf{C} \rho_+^{2N(j)+2}) \leq 2^9 \mathbf{h}_0 \mathbf{C} \leq 2^{-5}$$

This implies $\mathbf{h}_j \log(1 + 1/w_j) \leq 2\mathbf{h}_j k_{N(j)+1} < 2^{-1}$, which by inspection of (4.1c) and (4.2c) implies (a). To show (b), observe that by construction of $(V_n^*)_{n \geq 0}$,

$$\left(p_{n+1}^* + \frac{\mu_{n+1}^*}{\nu_{n+1}^*}, \frac{1 + w_{n+1}^*}{\nu_{n+1}^*}\right) = \mathcal{E}_R\{w_n^*\}\left(p_n^* + \frac{\mu_n^*}{\nu_n^*}, \frac{1 + w_n^*}{\nu_n^*}\right)$$

for all $n \geq 0$, see Definition 4.5 and Proposition 4.2. Since $V_{J(n)} = V_n^*$ for all $n \geq 0$ and since $\lfloor 1/w_n^* \rfloor = k_{n+1} = J(n+1) - J(n)$, the last equation is equivalent to

$$(q_{J(n+1)}, \mathbf{h}_{J(n+1)}) = \mathcal{Q}_R^{J(n+1)-J(n)}\{w_{J(n)}\}(q_{J(n)}, \mathbf{h}_{J(n)}) \quad (4.20)$$

By Proposition 4.2 and by the construction of $(V_j)_{j \geq 0}$, for all $j \geq 1$, $s = N(j) - 1$,

$$\left(p_j + \frac{\mu_j}{\nu_j}, \frac{1 + 1/w_s^* - j + J(s)}{\nu_j}\right) = \mathcal{Q}_R^{j-J(s)}\{w_s^*\}\left(p_s^* + \frac{\mu_s^*}{\nu_s^*}, \frac{1 + w_s^*}{\nu_s^*}\right)$$

Since $1/w_s^* - j + J(s) = w_j$, this implies $(q_j, \mathbf{h}_j) = \mathcal{Q}_R^{j-J(s)}\{w_{J(s)}\}(q_{J(s)}, \mathbf{h}_{J(s)})$, for all $j \geq 1$, $s = N(j) - 1$. The last identity and (4.20) imply $(q_j, \mathbf{h}_j) = \mathcal{Q}_R^j\{w_0\}(q_0, \mathbf{h}_0)$ for all $j \geq 0$, which is equivalent to (b).

To prove **Part 3**, check that for all $j \geq 0$ the following implication holds:

$$\left. \begin{aligned} w_{j+1} &= \mathcal{Q}_R(w_j) \\ (q_{j+1}, \mathbf{h}_{j+1}) &= \mathcal{Q}_R\{w_j\}(q_j, \mathbf{h}_j) \end{aligned} \right\} \implies \mathcal{Q}_L(\mathbf{h}_{j+1}, w_{j+1}, q_{j+1}) = (\mathbf{h}_j, w_j, q_j)$$

To make this calculation, distinguish the cases $j \in J(\mathbb{Z})$ and $j \notin J(\mathbb{Z})$, and recall $w_j, q_j \in (0, \infty) \setminus \mathbb{Z}$ and that $w_j \in (0, 1)$ iff $j \in J(\mathbb{Z})$ iff $q_{j+1} \in (0, 1)$. $\square$

5. An abstract semi-global existence theorem

This section is logically self-contained, and the notation is introduced from scratch. The objects in this section are abstractions of concrete objects that appear in other sections of this paper. This relationship is reflected in the choice of notation: abstract objects are named after their concrete counterparts, whenever possible. *This section is an independent unit. Definitions in other sections are irrelevant here and must be ignored.*

Definition 5.1. For every integer $d \geq 1$, denote by $\|\cdot\|$ the Euclidean distance in $\mathbb{R}^d$. Set $B[\delta, \mathbf{f}] = \{\mathbf{g} \in \mathbb{R}^d \mid \|\mathbf{g} - \mathbf{f}\| \leq \delta\}$ for every $\delta \geq 0$ and every $\mathbf{f} \in \mathbb{R}^d$.

Proposition 5.1. Fix an integer $d \geq 1$. Suppose:

- (a) $\mathcal{F} \subset \mathbb{R}^d$ is a nonempty open subset and $B\mathcal{F} = \{(\delta, \mathbf{f}) \in [0, \infty) \times \mathcal{F} \mid B[\delta, \mathbf{f}] \subset \mathcal{F}\}$.
- (b) $\Pi_j : \mathcal{F} \rightarrow \mathbb{R}^d$ is a continuous map, for every integer $j \geq 1$.
- (c) $\mathcal{Q}_L : \mathcal{F} \rightarrow \mathbb{R}^d$ and $\text{Err} : B\mathcal{F} \rightarrow [0, \infty)$ and $\text{Lip} : B\mathcal{F} \rightarrow [0, \infty)$ are maps such that for all $(\delta, \mathbf{f}) \in B\mathcal{F}$:

$$\sup_{j \geq 1} \sup_{\mathbf{g} \in B[\delta, \mathbf{f}]} \|\Pi_j(\mathbf{g}) - \mathcal{Q}_L(\mathbf{g})\| \leq \text{Err}(\delta, \mathbf{f}) \quad (5.1a)$$

$$\sup_{\mathbf{g}, \mathbf{g}' \in B[\delta, \mathbf{f}], \mathbf{g} \neq \mathbf{g}'} \frac{\|\mathcal{Q}_L(\mathbf{g}) - \mathcal{Q}_L(\mathbf{g}')\|}{\|\mathbf{g} - \mathbf{g}'\|} \leq \text{Lip}(\delta, \mathbf{f}) \quad (5.1b)$$

- (d) $(\delta_j, \mathbf{f}_j)_{j \geq 0}$ is a sequence in $B\mathcal{F}$ so that $\mathbf{f}_{j-1} = \mathcal{Q}_L(\mathbf{f}_j)$ for all $j \geq 1$, and so that

$$\sum_{n=j+1}^{\infty} \left\{ \prod_{k=j+1}^{n-1} \text{Lip}(\delta_k, \mathbf{f}_k) \right\} \text{Err}(\delta_n, \mathbf{f}_n) \leq \delta_j \quad (5.2)$$

for all $j \geq 0$.

Then, there exists a sequence $(\mathbf{g}_j)_{j \geq 0}$ with $\mathbf{g}_j \in B[\delta_j, \mathbf{f}_j] \subset \mathcal{F}$ such that for all $j \geq 1$:

$$\mathbf{g}_{j-1} = \Pi_j(\mathbf{g}_j)$$

Proof. For all integers $0 \leq j \leq \ell$, set

$$E_j^\ell = \sum_{n=j+1}^{\ell} \left\{ \prod_{k=j+1}^{n-1} \text{Lip}(\delta_k, \mathbf{f}_k) \right\} \text{Err}(\delta_n, \mathbf{f}_n) \in [0, \infty)$$

Then $E_j = \lim_{\ell \rightarrow \infty} E_j^\ell$ is the left hand side of (5.2). Observe that $E_j^j = 0$ and $E_j^\ell \leq E_j \leq \delta_j$ by (d). Moreover, $E_{j-1}^\ell = \text{Lip}(\delta_j, \mathbf{f}_j)E_j^\ell + \text{Err}(\delta_j, \mathbf{f}_j)$ when $1 \leq j \leq \ell$.

For all integers $0 \leq m \leq \ell$, let $(A)^{m, \ell}$ be the statement: *There is a finite sequence $\mathbf{g}^{m, \ell} = (\mathbf{g}_j^{m, \ell})_{m \leq j \leq \ell}$ with $\mathbf{g}_j^{m, \ell} \in B[E_j^\ell, \mathbf{f}_j] \subset B[\delta_j, \mathbf{f}_j] \subset \mathcal{F}$ for all $m \leq j \leq \ell$, such that $\mathbf{g}_\ell^{m, \ell} = \mathbf{f}_\ell$ and $\mathbf{g}_{j-1}^{m, \ell} = \Pi_j(\mathbf{g}_j^{m, \ell})$ when $m+1 \leq j \leq \ell$.* Observe that if $(A)^{m, \ell}$ is true, then the sequence $\mathbf{g}^{m, \ell}$ is unique.

For every fixed $\ell \geq 0$, we show by induction over m , one-by-one from $m = \ell$ down to $m = 0$, that $(A)^{m, \ell}$ is true. The base case $(A)^{\ell, \ell}$ is trivial. For the induction step, let $1 \leq m \leq \ell$ and suppose $(A)^{m, \ell}$ is true. Define $\mathbf{g}^{m-1, \ell}$ by $\mathbf{g}_j^{m-1, \ell} = \mathbf{g}_j^{m, \ell} \in B[E_j^\ell, \mathbf{f}_j] \subset B[\delta_j, \mathbf{f}_j] \subset \mathcal{F}$ when $m \leq j \leq \ell$, and set $\mathbf{g}_{m-1}^{m-1, \ell} = \Pi_m(\mathbf{g}_m^{m-1, \ell}) = \Pi_m(\mathbf{g}_m^{m, \ell}) \in \mathbb{R}^d$. The statement $(A)^{m-1, \ell}$ is true, if $\mathbf{g}_{m-1}^{m-1, \ell} \in B[E_{m-1}^\ell, \mathbf{f}_{m-1}]$, which follows from

$$\begin{aligned} \|\mathbf{g}_{m-1}^{m-1, \ell} - \mathbf{f}_{m-1}\| &= \|\Pi_m(\mathbf{g}_m^{m, \ell}) - \mathcal{Q}_L(\mathbf{f}_m)\| \\ &\leq \|\Pi_m(\mathbf{g}_m^{m, \ell}) - \mathcal{Q}_L(\mathbf{g}_m^{m, \ell})\| + \|\mathcal{Q}_L(\mathbf{g}_m^{m, \ell}) - \mathcal{Q}_L(\mathbf{f}_m)\| \\ &\leq \text{Err}(\delta_m, \mathbf{f}_m) + \text{Lip}(\delta_m, \mathbf{f}_m)E_m^\ell = E_{m-1}^\ell \end{aligned}$$

We have shown that $(A)^{m,\ell}$ is true for all $0 \leq m \leq \ell$. For all integers $0 \leq j \leq \ell$, set $\mathbf{g}_j^\ell = \mathbf{g}_j^{0,\ell} \in B[\delta_j, \mathbf{f}_j]$, where $\mathbf{g}^{0,\ell} = (\mathbf{g}_j^{0,\ell})_{0 \leq j \leq \ell}$ is the sequence in $(A)^{0,\ell}$. For every fixed $j \geq 0$, the sequence $\mathbf{g}_j = (\mathbf{g}_j^\ell)_{\ell \geq j}$ in the compact $B[\delta_j, \mathbf{f}_j]$ has a convergent subsequence $(\mathbf{g}_j^\ell)_{\ell \in \mathcal{L}_j}$, where $\mathcal{L}_j \subset [j, \infty) \cap \mathbb{Z}$ is infinite. One may choose $\mathcal{L}_0 \supset \mathcal{L}_1 \supset \dots$, that is $\mathcal{L}_{j-1} \supset \mathcal{L}_j$ for all $j \geq 1$. Pick a sequence $(\ell_j)_{j \geq 0}$ with $\ell_j \in \mathcal{L}_j$ for all $j \geq 0$, such that $\ell_{j-1} < \ell_j$ for all $j \geq 1$. Set $\mathcal{L} = \{\ell_j \mid j \geq 0\}$. By construction, all but a finite number of elements of $\mathcal{L}$ are in $\mathcal{L}_j$, for every $j \geq 0$. That is, $(\mathbf{g}_j^\ell)_{\ell \in \mathcal{L} \cap [j, \infty)}$ converges. Set $\mathbf{g}_j = \lim_{\ell \rightarrow \infty, \ell \in \mathcal{L} \cap [j, \infty)} \mathbf{g}_j^\ell \in B[\delta_j, \mathbf{f}_j]$. For all $j \geq 1$,

$$\begin{aligned} \Pi_j(\mathbf{g}_j) &= \lim_{\ell \rightarrow \infty, \ell \in \mathcal{L} \cap [j, \infty)} \Pi_j(\mathbf{g}_j^\ell) \\ &= \lim_{\ell \rightarrow \infty, \ell \in \mathcal{L} \cap [j, \infty)} \mathbf{g}_{j-1}^\ell = \mathbf{g}_{j-1} \end{aligned}$$

because Π_j is continuous by (b). $\square$

6. Main Theorems

In this section, τ_* , $\mathbf{K}$, $\mathcal{F}$ are given just as in Definitions 3.17, 3.18, 3.19, and $\mathcal{Q}_L$ is the map in Definition 3.16.

Definition 6.1. Let $\|\cdot\|$ be the Euclidean distance in $\mathbb{R}^3$. For every $\delta \geq 0$ and every $\mathbf{f} \in \mathbb{R}^3$, set $B[\delta, \mathbf{f}] = \{\mathbf{g} \in \mathbb{R}^3 \mid \|\mathbf{g} - \mathbf{f}\| \leq \delta\}$.

Definition 6.2. Let $\mathcal{F} \subset (0, \infty)^3$ be as in Definition 3.19. For all $\zeta \geq 1$ set

$$B_\zeta \mathcal{F} = \{(\delta, \mathbf{f}) \in [0, \infty) \times \mathcal{F} \mid B[\zeta\delta, \mathbf{f}] \subset \mathcal{F}\} \quad \text{and} \quad B\mathcal{F} = B_1 \mathcal{F}$$

Lemma 6.1. For all $(\delta, \mathbf{f}) \in B\mathcal{F}$ set

$$\begin{aligned} W(\delta, \mathbf{f}) &= \max\left\{\frac{1}{w-\delta}, w + \delta, \frac{1}{q-\delta}, \frac{1}{|q-1|-\delta}, q + \delta\right\} && \in [1, \infty) \\ W(\mathbf{f}) &= W(0, \mathbf{f}) = \max\left\{\frac{1}{w}, w, \frac{1}{q}, \frac{1}{|q-1|}, q\right\} && \in [1, \infty) \end{aligned}$$

where $\mathbf{f} = (\mathbf{h}, w, q)$. Then:

- (a) $W(\mathbf{g}) \leq W(\delta, \mathbf{f})$ for all $\mathbf{g} \in B[\delta, \mathbf{f}]$.
- (b) If $(\delta, \mathbf{f}) \in B_2 \mathcal{F} \subset B\mathcal{F}$ then $W(\delta, \mathbf{f}) \leq 2W(\mathbf{f})$.

Lemma 6.2. Let $\text{Err} : B\mathcal{F} \rightarrow [0, \infty)$ be given by

$$\text{Err}(\delta, \mathbf{f}) = 2^{40} \left(\frac{1}{\mathbf{h}-\delta}\right)^2 W(\delta, \mathbf{f})^5 \exp\left(-\frac{1}{\mathbf{h}} 2^{-9} W(\delta, \mathbf{f})^{-2}\right)$$

where $\mathbf{f} = (\mathbf{h}, w, q)$. Then for all $(\delta, \mathbf{f}) \in B\mathcal{F}$, we have $\mathbf{K}(\mathbf{g}) \leq \text{Err}(\delta, \mathbf{f})$ for all $\mathbf{g} \in B[\delta, \mathbf{f}] \subset \mathcal{F}$ (see Definition 3.18).

Proof. Let $\mathbf{g} = (\mathbf{h}', w', q') \in B[\delta, \mathbf{f}]$. Then $\tau_*(\mathbf{g}) \geq \frac{1}{2}W(\mathbf{g})^{-2}$ and $0 < \mathbf{h} - \delta \leq \mathbf{h}' \leq \mathbf{h} + \delta \leq 2\mathbf{h}$ and $\frac{1}{\mathbf{h}'} \geq \frac{1}{2\mathbf{h}}$. Hence, $\mathbf{K}(\mathbf{g}) \leq 2^{40} \left(\frac{1}{\mathbf{h}-\delta}\right)^2 W(\mathbf{g})^5 \exp\left(-\frac{1}{\mathbf{h}} 2^{-9} W(\mathbf{g})^{-2}\right)$. Now use Lemma 6.1 (a). $\square$

Lemma 6.3. Let $\mathcal{Q}_L$ be as in Definition 3.16. Set $\text{Lip} : B\mathcal{F} \rightarrow [0, \infty)$, $\text{Lip}(\delta, \mathbf{f}) = 2^{13}W(\delta, \mathbf{f})^3$. Then $\|\mathcal{Q}_L(\mathbf{g}) - \mathcal{Q}_L(\mathbf{g}')\| \leq \text{Lip}(\delta, \mathbf{f}) \|\mathbf{g} - \mathbf{g}'\|$ for all $\mathbf{g}, \mathbf{g}' \in B[\delta, \mathbf{f}]$.

Proof. Let $\mathbf{f} = (\mathbf{h}, w, q)$. If $\mathbf{g} = \mathbf{g}'$, there is nothing to prove. Suppose $\mathbf{g} \neq \mathbf{g}'$. In Lemma B.1 of Appendix B, set $\mathbf{f}_1 = (\mathbf{h}_1, w_1, q_1) = \mathbf{g}$ and $\mathbf{f}_2 = (\mathbf{h}_2, w_2, q_2) = \mathbf{g}'$. Observe that $0 < \mathbf{h}_i \leq 1$ by $\mathbf{g}, \mathbf{g}' \in B[\delta, \mathbf{f}] \subset \mathcal{F}$. Since $\delta < |q - 1|$, either $q, q_1, q_2 < 1$ or $q, q_1, q_2 > 1$. We have $w_{\max} \leq \max\{W(\mathbf{g}), W(\mathbf{g}')\}$ and $q_{\max} \leq \max\{W(\mathbf{g}), W(\mathbf{g}')\}$ and $q_{\min}^{-1} = \max\{q_1^{-1}, q_2^{-1}\} \leq \max\{W(\mathbf{g}), W(\mathbf{g}')\}$. Now use $\log(2 + w_{\max}) \leq 1 + w_{\max}$ and Lemma 6.1 (a). $\square$

Theorem 6.1 (Main Theorem 1). Recall the definitions of $\mathcal{P}_L$ and $\mathcal{Q}_L$ (Definition 3.16), $\mathcal{F}$ (Definition 3.19), Π (Proposition 3.3), $B_\zeta \mathcal{F}$ (Definition 6.2), W (Lemma 6.1), Err (Lemma 6.2), Lip (Lemma 6.3). Suppose:

- (a) $(\mathbf{f}_j)_{j \geq 0}$, with $\mathbf{f}_j = (\mathbf{h}_j, w_j, q_j) \in \mathcal{F}$, satisfies $\mathbf{f}_{j-1} = \mathcal{Q}_L(\mathbf{f}_j)$ for all $j \geq 1$.
- (b) The sequence $(\delta_j)_{j \geq 0}$ given by

$$\delta_j = \sum_{\ell=j+1}^{\infty} \left\{ \prod_{k=j+1}^{\ell-1} 2^{16} W(\mathbf{f}_k)^3 \right\} 2^{47} \left(\frac{1}{\mathbf{h}_\ell} \right)^2 W(\mathbf{f}_\ell)^5 \exp \left(- \frac{1}{\mathbf{h}_\ell} 2^{-11} W(\mathbf{f}_\ell)^{-2} \right)$$

satisfies $\delta_j < \infty$ and $(\delta_j, \mathbf{f}_j) \in B_2 \mathcal{F}$ for all $j \geq 0$.

- (c) $\pi_0 \in S_3$ and $(\pi_j)_{j \geq 0}$ is the unique sequence in S_3 that satisfies $\pi_{j-1} = \mathcal{P}_L(\pi_j, \mathbf{f}_j)$ for all $j \geq 1$.
- (d) $\sigma_* \in \{-1, +1\}^3$.

Then, there exists a sequence $(\mathbf{g}_j)_{j \geq 0}$ with $\mathbf{g}_j \in B[\delta_j, \mathbf{f}_j] \subset \mathcal{F}$ such that for all $j \geq 1$:

$$\mathbf{g}_{j-1} = \Pi[\pi_j, \sigma_*](\mathbf{g}_j) \quad \text{and} \quad \pi_{j-1} = \mathcal{P}_L(\pi_j, \mathbf{g}_j)$$

Proof. We use Proposition 5.1, with the understanding that the abstract objects of Proposition 5.1 in the left column are given by the special objects in the right column:

d	3
$\mathcal{F}$	$\mathcal{F}$ as in Definition 3.19
Π_j	$\Pi[\pi_j, \sigma_*]$, see Proposition 3.3 and the hypotheses Theorem 6.1 (c), (d)
$\mathcal{Q}_L$	$\mathcal{Q}_L _{\mathcal{F}}$, with $\mathcal{Q}_L$ as in Definition 3.16
Err	Err as in Lemma 6.2
Lip	Lip as in Lemma 6.3
$(\delta_j, \mathbf{f}_j)$	$(\delta_j, \mathbf{f}_j)$ as in hypotheses Theorem 6.1 (a) and (b)

We check that the assumptions (a), (b), (c), (d) of Proposition 5.1 are satisfied:

- (a) The definitions of $B\mathcal{F}$ in Proposition 5.1 and in Definition 6.2 are consistent.
- (b) $\Pi[\pi_j, \sigma_*] : \mathcal{F} \rightarrow (0, \infty)^2 \times \mathbb{R} \subset \mathbb{R}^3$ is continuous, by Proposition 3.3.
- (c) The domains of definition of $\mathcal{Q}_L|_{\mathcal{F}}$ and Err and Lip are just as required by Proposition 5.1 (c). For all $(\delta, \mathbf{f}) \in B\mathcal{F}$ and $\mathbf{g}, \mathbf{g}' \in B[\delta, \mathbf{f}] \subset \mathcal{F}$ and $j \geq 1$,

$$\begin{aligned} \|\Pi[\pi_j, \sigma_*](\mathbf{g}) - \mathcal{Q}_L|_{\mathcal{F}}(\mathbf{g})\| &\leq \mathbf{K}(\mathbf{g}) \leq \text{Err}(\delta, \mathbf{f}) \\ \|\mathcal{Q}_L|_{\mathcal{F}}(\mathbf{g}) - \mathcal{Q}_L|_{\mathcal{F}}(\mathbf{g}')\| &\leq \text{Lip}(\delta, \mathbf{f}) \|\mathbf{g} - \mathbf{g}'\| \end{aligned}$$

by Proposition 3.3 (a) and by Lemmas 6.2 and 6.3. That is, (5.1a) and (5.1b) hold.

- (d) By assumption, $(\delta_j, \mathbf{f}_j) \in B_2\mathcal{F} \subset B\mathcal{F}$ for all $j \geq 0$. Hence $\frac{1}{2}\mathbf{h}_j \leq \mathbf{h}_j - \delta_j$ and, by Lemma 6.1 (b), we have $W(\delta_j, \mathbf{f}_j) \leq 2W(\mathbf{f}_j)$. Consequently, for all $j \geq 0$,

$$\begin{aligned} & \sum_{\ell=j+1}^{\infty} \left\{ \prod_{k=j+1}^{\ell-1} \text{Lip}(\delta_k, \mathbf{f}_k) \right\} \text{Err}(\delta_\ell, \mathbf{f}_\ell) \\ &= \sum_{\ell=j+1}^{\infty} \left\{ \prod_{k=j+1}^{\ell-1} 2^{13} W(\delta_k, \mathbf{f}_k)^3 \right\} 2^{40} \left(\frac{1}{\mathbf{h}_\ell - \delta_\ell} \right)^2 W(\delta_\ell, \mathbf{f}_\ell)^5 \\ & \quad \times \exp \left(- \frac{1}{\mathbf{h}_\ell} 2^{-9} W(\delta_\ell, \mathbf{f}_\ell)^{-2} \right) \\ &\leq \sum_{\ell=j+1}^{\infty} \left\{ \prod_{k=j+1}^{\ell-1} 2^{16} W(\mathbf{f}_k)^3 \right\} 2^{47} \left(\frac{1}{\mathbf{h}_\ell} \right)^2 W(\mathbf{f}_\ell)^5 \exp \left(- \frac{1}{\mathbf{h}_\ell} 2^{-11} W(\mathbf{f}_\ell)^{-2} \right) \end{aligned}$$

The last expression is equal to δ_j , and (5.2) is checked.

Now, Theorem 6.1 follows from Proposition 5.1. $\square$

Theorem 6.2 (Main Theorem 2). *Suppose the vector $\mathbf{f}_0 = (\mathbf{h}_0, w_0, q_0)$ satisfies the assumptions of Proposition 4.4, that is*

$$w_0 \in (0, 1) \setminus \mathbb{Q} \quad \mathbf{C}(w_0, q_0) < \infty \quad (6.1a)$$

$$q_0 \in (0, \infty) \setminus \mathbb{Q} \quad 0 < \mathbf{h}_0 \leq 2^{-14} (\mathbf{C}(w_0, q_0))^{-1} \quad (6.1b)$$

Let $(k_n)_{n \in \mathbb{Z}}$ and $J : \mathbb{Z} \rightarrow \mathbb{Z}$ (Era Pointer) and $N : \mathbb{Z} \rightarrow \mathbb{Z}$ (Era Counter) and $(w_j)_{j \in \mathbb{Z}}$, $(q_j)_{j \geq 0}$, $(\mathbf{h}_j)_{j \geq 0}$ be just as in Proposition 4.4. Introduce the sequence $(\mathbf{f}_j)_{j \geq 0}$ by

$$\mathbf{f}_j = (\mathbf{h}_j, w_j, q_j) \in (0, \infty)^3$$

Introduce sequences $(\mathbf{H}_j)_{j \geq 0}$ and $(K_j)_{j \geq 0}$ by

$$\begin{aligned} \mathbf{H}_j &= \mathbf{h}_0 \frac{1+w_j}{1+w_0} \prod_{\ell=0}^{N(j)-1} w_{J(\ell)} w_{J(\ell-1)} &> 0 \\ K_j &= \max\{k_{N(j)-2}, k_{N(j)-1}, k_{N(j)}, k_{N(j)+1}\} &\geq 1 \end{aligned}$$

Suppose:

- (a) $\mathbf{H}_j < 2^{-21} (K_j)^{-2}$ for all $j \geq 0$.
(b) $2^{71} \left(\frac{1}{\mathbf{H}_j} \right)^2 (K_j)^5 \exp \left(- \frac{1}{\mathbf{H}_j} 2^{-21} (K_j)^{-2} \right) < 1$ for all $j \geq 0$.
(c) The sequence $(\delta_j)_{j \geq 0}$ given by

$$\delta_j = \sum_{\ell=j+1}^{\infty} \left\{ \prod_{k=j+1}^{\ell-1} 2^{28} (K_k)^3 \right\} 2^{71} \left(\frac{1}{\mathbf{H}_\ell} \right)^2 (K_\ell)^5 \exp \left(- \frac{1}{\mathbf{H}_\ell} 2^{-21} (K_\ell)^{-2} \right) > 0$$

satisfies $\delta_j \leq 2^{-4} \mathbf{H}_j < \infty$.

- (d) $\pi_0 \in S_3$ and $(\pi_j)_{j \geq 0}$ is the unique sequence in S_3 that satisfies $\pi_{j-1} = \mathcal{P}_L(\pi_j, \mathbf{f}_j)$ for all $j \geq 1$.
(e) $\sigma_* \in \{-1, +1\}^3$.

Then $(\delta_j, \mathbf{f}_j) \in B_2\mathcal{F}$ for all $j \geq 0$ and there exists a sequence $(\mathbf{g}_j)_{j \geq 0}$ with $\mathbf{g}_j \in B[\delta_j, \mathbf{f}_j] \subset \mathcal{F}$ such that for all $j \geq 1$:

$$\mathbf{g}_{j-1} = \Pi[\pi_j, \sigma_*](\mathbf{g}_j) \quad \text{and} \quad \pi_{j-1} = \mathcal{P}_L(\pi_j, \mathbf{g}_j)$$

Proof. By Proposition 4.4 and by hypotheses (a), (c) in Theorem 6.2, for all $j \geq 0$:

$$2^{-2}\mathbf{H}_j \leq \mathbf{h}_j \leq 2^2\mathbf{H}_j \quad (6.2a)$$

$$\max\left\{\frac{1}{w_j}, w_j, \frac{1}{q_j}, \frac{1}{|q_j-1|}, q_j\right\} \leq 2^4 K_j \quad (6.2b)$$

$$2^{-4}(K_j)^{-1} \leq \min\{w_j, q_j, |q_j - 1|\}$$

$$2\delta_j \leq 2^{-1} \min\{w_j, q_j, |q_j - 1|, \mathbf{h}_j\}$$

Hence, $B[2\delta_j, \mathbf{f}_j] \subset (0, \infty)^3$ for every $j \geq 0$. Furthermore, for all $j \geq 0$ and all $(\mathbf{h}', w', q') \in B[2\delta_j, \mathbf{f}_j] \subset (0, \infty)^3$, we have $q' \neq 1$ and

$$2^{-3}\mathbf{H}_j \leq 2^{-1}\mathbf{h}_j \leq \mathbf{h}_j - 2\delta_j \leq \mathbf{h}' \leq \mathbf{h}_j + 2\delta_j \leq 2\mathbf{h}_j \leq 2^3\mathbf{H}_j \quad (6.3a)$$

and

$$\begin{aligned} & \max\left\{\frac{1}{w'}, w', \frac{1}{q'}, \frac{1}{|q'-1|}, q'\right\} \\ & \leq \max\left\{\frac{1}{w_j - 2\delta_j}, w_j + 2\delta_j, \frac{1}{q_j - 2\delta_j}, \frac{1}{|q_j - 1| - 2\delta_j}, q_j + 2\delta_j\right\} \\ & \leq 2 \max\left\{\frac{1}{w_j}, w_j, \frac{1}{q_j}, \frac{1}{|q_j - 1|}, q_j\right\} \leq 2^5 K_j \end{aligned} \quad (6.3b)$$

The last two estimates (6.3) imply $\tau_*(\mathbf{h}', w', q') \geq 2^{-11}(K_j)^{-2}$ and

$$\mathbf{K}(\mathbf{h}', w', q') \leq 2^{71} \left(\frac{1}{\mathbf{H}_j}\right)^2 (K_j)^5 \exp\left(-\frac{1}{\mathbf{H}_j} 2^{-21}(K_j)^{-2}\right) < 1$$

The last inequality is hypothesis (b) in Theorem 6.2. Furthermore,

$$\mathbf{h}' \leq 2^3\mathbf{H}_j < 2^{-18}(K_j)^{-2} \leq 2^{-7}\tau_*(\mathbf{h}', w', q')$$

The second inequality is hypothesis (a) in Theorem 6.2. These estimates are true for all $(\mathbf{h}', w', q') \in B[2\delta_j, \mathbf{f}_j]$, and therefore $B[2\delta_j, \mathbf{f}_j] \subset \mathcal{F}$ for all $j \geq 0$, in particular $\mathbf{f}_j \in \mathcal{F}$ (see Definition 3.19). In other words, $(\delta_j, \mathbf{f}_j) \in B_2\mathcal{F}$.

The last result and the fact that $\mathbf{f}_{j-1} = \mathcal{Q}_L(\mathbf{f}_j)$ for all $j \geq 1$ (see Proposition 4.4) imply that Theorem 6.2 follows from Theorem 6.1, if we can show that $\delta_j \leq \delta_j$ for all $j \geq 0$, where δ_j is given as in Theorem 6.1. The inequality $\delta_j \leq \delta_j$ is a consequence of $W(\mathbf{f}_\ell) \leq 2^4 K_\ell$ and $2^{-2}\mathbf{H}_\ell \leq \mathbf{h}_\ell \leq 2^2\mathbf{H}_\ell$, where $j, \ell \geq 0$. $\square$

Theorem 6.3 (Main Theorem 3). Fix constants $\mathbf{D} \geq 1$, $\gamma \geq 0$. Suppose the vector $\mathbf{f}_0 = (\mathbf{h}_0, w_0, q_0) \in (0, \infty)^3$ satisfies

- (i) $w_0 \in (0, 1) \setminus \mathbb{Q}$ and $q_0 \in (0, \infty) \setminus \mathbb{Q}$.
- (ii) $k_n \leq \mathbf{D} \max\{1, n\}^\gamma$ for all $n \geq -2$, with $(k_n)_{n \in \mathbb{Z}}$ as in Proposition 4.4, that is

$$(1 + q_0)^{-1} = \langle k_0, k_{-1}, k_{-2}, \dots \rangle \quad w_0 = \langle k_1, k_2, k_3, \dots \rangle$$

- (iii) $0 < \mathbf{h}_0 < \mathbf{A}^\#$ where $\mathbf{A}^\# = \mathbf{A}^\#(\mathbf{D}, \gamma) = 2^{-56}\mathbf{D}^{-4}(4(\gamma + 1))^{-4(\gamma + 1)}$.

Then

- The assumptions (6.1) and (a), (b), (c) of Theorem 6.2 hold.
- Set $\rho_+ = \frac{1}{2}(1 + \sqrt{5})$. The sequence $(\delta_j)_{j \geq 0}$ in Theorem 6.2 satisfies for all $j \geq 0$:

$$\delta_j \leq \exp\left(-\frac{1}{\mathbf{h}_0}\mathbf{A}^\#\rho_+^{N(j)}\right) \quad \text{and} \quad N(j) \geq (\mathbf{D}^{-1}j)^{1/(\gamma+1)} \quad (6.4)$$

where $N : \mathbb{Z} \rightarrow \mathbb{Z}$ (Era Counter) is the map in Proposition 4.4.

If $\gamma > 1$ and $\mathbf{D} > \frac{1}{\log 2} \frac{\gamma}{\gamma-1}$, then the set of all vectors $\mathbf{f}_0 \in (0, \infty)^3$ that satisfy (i), (ii), (iii) has positive Lebesgue measure.

Proof. Preliminaries. The following facts will be used without further comment:

- $a^{-x}x^b \leq \left(\frac{b}{e \log a}\right)^b$ for all real numbers $a > 1, b > 0, x \geq 0$ where $e = \exp(1)$.
- $a^b \leq c^d$ for all real numbers $1 \leq a \leq c$ and $0 \leq b \leq d$.
- $1 < \rho_+ < 2$ and $1 < e \log \rho_+ < 2$ where $e = \exp(1)$ and $\rho_+ = \frac{1}{2}(1 + \sqrt{5})$.

Fix $\mathbf{D} \geq 1$ and $\gamma \geq 0$ as in Theorem 6.3. For all 5-tuples of real numbers $s = (s_1, s_2, s_3, s_4, s_5) \geq (0, 0, 0, 1, 0)$, set $\mathbf{A}(s) = 2^{-s_1-s_2\gamma} \mathbf{D}^{-s_3} (s_4(\gamma+1))^{-s_5(\gamma+1)}$. Observe that $0 < \mathbf{A}(s) \leq 2^{-s_1} \leq 1$ and $\mathbf{A}(s) \leq \mathbf{A}(s')$ if $s \geq s'$.

Basic smallness assumptions. $k_n \leq \mathbf{D} \max\{1, n\}^\gamma$ for all $n \geq -2$ and $\mathbf{h}_0 < \mathbf{A}(\kappa)$. The vector $\kappa = (\kappa_1, \kappa_2, \kappa_3, \kappa_4, \kappa_5) \geq (0, 0, 0, 1, 0)$ will be fixed during the proof.

Estimates 1. Recall Proposition 4.4 and $\rho_+ = \frac{1}{2}(1 + \sqrt{5})$. For all $j \geq 0, n \geq 0$:

$$\begin{aligned}
\mathbf{C}(w_0, q_0) &= \sup_{n \geq 0} (n+1) \rho_+^{-2n} k_n \max\{k_{n-1}, k_{n-2}\} \\
&\leq 2\mathbf{D}^2 \sup_{n \geq 0} \rho_+^{-2n} \max\{1, n\}^{2(\gamma+1)} \\
&\leq 2\mathbf{D}^2 (\gamma+1)^{2(\gamma+1)} = \mathbf{A}(1, 0, 2, 1, 2)^{-1} \\
J(n) &= \sum_{\ell=1}^n k_\ell \leq \mathbf{D} \sum_{\ell=1}^n \ell^\gamma \leq \mathbf{D} n^{\gamma+1} \\
j &\leq J(N(j)) \leq \mathbf{D} N(j)^{\gamma+1} \\
N(j) &\geq (\mathbf{D}^{-1} j)^{1/(\gamma+1)} \\
\mathbf{H}_j &\leq 2^4 \mathbf{h}_0 \rho_+^{-2N(j)} \quad \text{see (4.19)} \\
\mathbf{H}_j &\geq 2^{-1} \mathbf{h}_0 \prod_{\ell=0}^{N(j)-1} (k_\ell + 1)^{-1} (k_{\ell+1} + 1)^{-1} \\
&\geq 2^{-1} \mathbf{h}_0 \prod_{\ell=0}^{N(j)-1} (2\mathbf{D}(\ell+1)^\gamma)^{-2} \geq 2^{-1} \mathbf{h}_0 \max\{1, 2\mathbf{D}N(j)^\gamma\}^{-2N(j)} \\
K_j &\leq \mathbf{D}(N(j)+1)^\gamma \leq \mathbf{D} 2^\gamma \max\{1, N(j)\}^\gamma \\
\mathbf{H}_j K_j^2 &\leq 2^{4+2\gamma} \mathbf{D}^2 \mathbf{h}_0 \rho_+^{-2N(j)} \max\{1, N(j)\}^{2\gamma} \\
&\leq 2^{4+2\gamma} \mathbf{D}^2 \mathbf{h}_0 \sup_{n \geq 0} \rho_+^{-2n} \max\{1, n\}^{2(\gamma+1)} \\
&\leq 2^{4+2\gamma} \mathbf{D}^2 \mathbf{h}_0 (\gamma+1)^{2(\gamma+1)} = \mathbf{h}_0 \mathbf{A}(4, 2, 2, 1, 2)^{-1}
\end{aligned}$$

Require $\kappa \geq (25, 2, 2, 1, 2)$. Then $\mathbf{H}_j < 2^{-21} (K_j)^{-2}$ and $\mathbf{h}_0 \leq 2^{-14} (\mathbf{C}(w_0, q_0))^{-1}$.

Estimates 2. Let $(\delta_j)_{j \geq 0}$ be as in Theorem 6.2. We claim that with proper choice of κ :

- (A) $\delta_{J(n)} \leq 2^{-5} \mathbf{h}_0 (2\mathbf{D}(n+1)^\gamma)^{-2(n+1)} \exp(-\frac{1}{\mathbf{h}_0} \mathbf{A}(\kappa) \rho_+^{n+1})$ for all $n \geq 0$.
 (B) $\delta_j \leq 2^{-4} \mathbf{H}_j$ and $\delta_j \leq \exp(-\frac{1}{\mathbf{h}_0} \mathbf{A}(\kappa) \rho_+^{N(j)})$ for all $j \geq 0$.

We first check (A) $\implies$ (B). Note that $\delta_j \geq \delta_{j+1}$, $j \geq 0$. Fix any $j \geq 0$. Set $n = N(j+1)-1 \geq 0$. By (A), by $j \geq J(n)$ (see the line before (4.16)) and by $n+1 \geq N(j)$,

$$\begin{aligned}
\delta_j &\leq \delta_{J(n)} \leq 2^{-5} \mathbf{h}_0 (2\mathbf{D}(n+1)^\gamma)^{-2(n+1)} \exp(-\frac{1}{\mathbf{h}_0} \mathbf{A}(\kappa) \rho_+^{n+1}) \\
&\leq \left(2^{-5} \mathbf{h}_0 \max\{1, 2\mathbf{D}N(j)^\gamma\}^{-2N(j)}\right) \exp(-\frac{1}{\mathbf{h}_0} \mathbf{A}(\kappa) \rho_+^{N(j)})
\end{aligned}$$

See the second bullet in the preliminaries. On the right hand side, both factors are ≤ 1 (use $\mathbf{h}_0 < \mathbf{A}(\kappa) \leq 1$). By the lower bound on $\mathbf{H}_j$ derived above, claim (B) follows.

We now check (A). For all $n \geq 0$:

$$\begin{aligned}
& \delta_{J(n)} \\
&= \sum_{m=n}^{\infty} \sum_{\ell=J(m)+1}^{J(m+1)} \left\{ \prod_{k=J(n)+1}^{\ell-1} 2^{28} (K_k)^3 \right\} 2^{71} \left(\frac{1}{\mathbf{H}_\ell} \right)^2 (K_\ell)^5 \\
&\quad \times \exp \left(- (2^{21} \mathbf{H}_\ell K_\ell^2)^{-1} \right) \\
&\leq \sum_{m=n}^{\infty} \sum_{\ell=J(m)+1}^{J(m+1)} (2^{15} \max_{1 \leq k \leq \ell} K_k)^{3\ell+2} \left(\frac{1}{2\mathbf{H}_\ell} \right)^2 \exp \left(- (2^{21} \mathbf{H}_\ell K_\ell^2)^{-1} \right) \\
&\leq \sum_{m=n}^{\infty} k_{m+1} (2^{15+\gamma} \mathbf{D}(m+1)^\gamma)^{3J(m+1)+2} \left(\frac{1}{\mathbf{h}_0} \right)^2 (2\mathbf{D}(m+1)^\gamma)^{4(m+1)} \\
&\quad \times \exp \left(- 2^{-25-2\gamma} \mathbf{D}^{-2} \frac{1}{\mathbf{h}_0} \rho_+^{2(m+1)} (m+1)^{-2\gamma} \right) \\
&\leq \left(\frac{1}{\mathbf{h}_0} \right)^2 \sum_{m=n+1}^{\infty} (2^{15+\gamma} \mathbf{D} m^\gamma)^{(10\mathbf{D} m^{\gamma+1})} \exp \left(- 2^{-25-2\gamma} \mathbf{D}^{-2} \frac{1}{\mathbf{h}_0} \rho_+^{2m} m^{-2\gamma} \right)
\end{aligned}$$

Since $2^5 \frac{1}{\mathbf{h}_0} (2\mathbf{D}(n+1)^\gamma)^{2(n+1)} \leq \frac{1}{\mathbf{h}_0} (2^6 \mathbf{D} m^\gamma)^{2m}$ for all $m \geq n+1$, we have

$$\begin{aligned}
\mathbf{S}(n) &\stackrel{\text{def}}{=} \delta_{J(n)} 2^5 \frac{1}{\mathbf{h}_0} (2\mathbf{D}(n+1)^\gamma)^{2(n+1)} \\
&\leq \left(\frac{1}{\mathbf{h}_0} \right)^3 \sum_{m=n+1}^{\infty} (2^{15+\gamma} \mathbf{D} m^\gamma)^{(12\mathbf{D} m^{\gamma+1})} \exp \left(- 2^{-25-2\gamma} \mathbf{D}^{-2} \frac{1}{\mathbf{h}_0} \rho_+^{2m} m^{-2\gamma} \right) \\
&\leq \left(\frac{1}{\mathbf{h}_0} \right)^3 \sum_{m=n+1}^{\infty} \exp \left(12\mathbf{D} m^{\gamma+1} \log (2^{15+\gamma} \mathbf{D} m^\gamma) - 2^{-25-2\gamma} \mathbf{D}^{-2} \frac{1}{\mathbf{h}_0} \rho_+^{2m} m^{-2\gamma} \right) \\
&\leq \left(\frac{1}{\mathbf{h}_0} \right)^3 \sum_{m=n+1}^{\infty} \exp \left(2^9 \mathbf{D}^2 (\gamma+1) m^{\gamma+2} - 2^{-25-2\gamma} \mathbf{D}^{-2} \frac{1}{\mathbf{h}_0} \rho_+^{2m} m^{-2\gamma} \right)
\end{aligned}$$

The second term in the argument of the exponential dominates the first term, if we require $\kappa \geq (35, 2, 4, \frac{3}{2}, 3)$. More precisely, the absolute value of the second term is at least twice the absolute value of the first term. In fact,

$$\begin{aligned}
& 2^{35+2\gamma} \mathbf{D}^4 (\gamma+1) \sup_{m \geq 1} \rho_+^{-2m} m^{3\gamma+2} \\
& \leq 2^{35+2\gamma} \mathbf{D}^4 \left(\frac{3}{2} (\gamma+1) \right)^{3(\gamma+1)} = \mathbf{A}(35, 2, 4, \frac{3}{2}, 3)^{-1} \leq \mathbf{A}(\kappa)^{-1} \leq \frac{1}{\mathbf{h}_0}
\end{aligned}$$

Therefore,

$$\mathbf{S}(n) \leq \left(\frac{1}{\mathbf{h}_0} \right)^3 \sum_{m=n+1}^{\infty} \exp \left(- 2^{-26-2\gamma} \mathbf{D}^{-2} \frac{1}{\mathbf{h}_0} \rho_+^{2m} m^{-2\gamma} \right)$$

Moreover, $2^{26+2\gamma} \mathbf{D}^2 \sup_{m \geq 1} \rho_+^{-m} m^{2\gamma} \leq 2^{26+2\gamma} \mathbf{D}^2 (2(\gamma+1))^{2(\gamma+1)} = 2^{-2} \mathbf{A}_*^{-1}$, where $\mathbf{A}_* = \mathbf{A}(28, 2, 2, 2, 2)$. Require $\kappa \geq (28, 2, 2, 2, 2)$. Then $\mathbf{h}_0 \leq \mathbf{A}_*$ and

$$\begin{aligned}
\mathbf{S}(n) &\leq \left(\frac{1}{\mathbf{h}_0} \right)^3 \sum_{m=n+1}^{\infty} \exp \left(- 4 \frac{1}{\mathbf{h}_0} \mathbf{A}_* \rho_+^m \right) \\
&\leq \exp \left(- \frac{1}{\mathbf{h}_0} \mathbf{A}_* \rho_+^{n+1} \right) \left(\left(\frac{1}{\mathbf{h}_0} \right)^3 \exp \left(- \frac{1}{\mathbf{h}_0} \mathbf{A}_* \right) \right) \sum_{m=1}^{\infty} \exp \left(- 2 \rho_+^m \right)
\end{aligned}$$

We have $\sum_{m=1}^{\infty} \exp \left(- 2 \rho_+^m \right) \leq \frac{1}{2} \sum_{m=1}^{\infty} \rho_+^{-m} = \frac{1}{2} (\rho_+ - 1)^{-1} = \frac{1}{2} \rho_+ \leq 1$. Require $\kappa \geq (56, 4, 4, 2, 4)$. Then $\mathbf{h}_0 \leq \mathbf{A}(\kappa) \leq \mathbf{A}(56, 4, 4, 2, 4) = \mathbf{A}_*^2$, and

$$\left(\frac{1}{\mathbf{h}_0} \right)^3 \exp \left(- \frac{1}{\mathbf{h}_0} \mathbf{A}_* \right) \leq \left(\frac{1}{\mathbf{h}_0} \right)^3 \exp \left(- \left(\frac{1}{\mathbf{h}_0} \right)^{1/2} \right) \leq 8! \mathbf{h}_0 \leq 2^{16} \mathbf{h}_0 \leq 1$$

Since $\mathbf{A}_* \geq \mathbf{A}(\kappa)$, we have $\mathbf{S}(n) \leq \exp(-\frac{1}{\mathbf{h}_0} \mathbf{A}(\kappa) \rho_+^{n+1})$. Fix $\kappa = (56, 4, 4, 2, 4)$. All the inequalities for κ hold, and claim (A) is proved. Let $\mathbf{A}^\sharp = \mathbf{A}(56, 0, 4, 4, 4)$, as in the statement of Theorem 6.3. Since $\mathbf{A}^\sharp \leq \mathbf{A}(\kappa)$, the condition $\mathbf{h}_0 < \mathbf{A}^\sharp$ in the statement of Theorem 6.3 implies the condition $\mathbf{h}_0 < \mathbf{A}(\kappa)$ used in this proof.

So far, we have verified the estimate (6.4), and we have verified the assumptions Theorem 6.2 (a), (c) and (6.1). In the assumption Theorem 6.2 (b), the cases $j \geq 1$ follow from Theorem 6.2 (a), (c). Since $\mathbf{H}_0 = \mathbf{h}_0$ and $K_0 \leq \mathbf{D}$, the remaining $j = 0$ case in Theorem 6.2 (b) follows from

$$\begin{aligned} 2^{71} \left(\frac{1}{\mathbf{H}_0}\right)^2 (K_0)^5 \exp\left(-\frac{1}{\mathbf{H}_0} 2^{-21} (K_0)^{-2}\right) &\leq 2^{71} \left(\frac{1}{\mathbf{h}_0}\right)^2 \mathbf{D}^5 \exp\left(-\frac{1}{\mathbf{h}_0} 2^{-21} \mathbf{D}^{-2}\right) \\ &\leq 2^{71} \left(\frac{1}{\mathbf{h}_0}\right)^2 \mathbf{D}^5 8! (\mathbf{h}_0 2^{21} \mathbf{D}^2)^8 \leq 2^{255} \mathbf{D}^{21} \mathbf{h}_0^6 \leq (\mathbf{h}_0 / \mathbf{A}^\sharp)^6 < 1 \end{aligned}$$

Lebesgue measure of the set of admissible $\mathbf{f}_0$. The set of all $\mathbf{f}_0 = (\mathbf{h}_0, w_0, q_0) \in (0, \infty)^3$ that satisfy (i), (ii), (iii) is a product $(0, \mathbf{A}^\sharp) \times F_w \times F_q$ (depending on $\mathbf{D}$ and γ), where $F_w \subset (0, 1) \setminus \mathbb{Q}$ and $F_q \subset (0, \infty) \setminus \mathbb{Q}$. Both $(0, \mathbf{A}^\sharp)$ and F_q have positive measure, because $\mathbf{A}^\sharp > 0$ and $(\frac{1}{2}, \frac{2}{3}) \setminus \mathbb{Q} \subset F_q$. In fact, if $q_0 \in (\frac{1}{2}, \frac{2}{3})$, then $1/(1+q_0) = 1/(1+1/(1+1/(1+x)))$ with $x = (2q_0 - 1)/(1 - q_0) \in (0, 1)$, that is $k_0 = k_{-1} = k_{-2} = 1 \leq \mathbf{D}$. Suppose $\gamma > 1$ and $\mathbf{D} > (\log 2)^{-1} \gamma / (\gamma - 1)$. Let $G(x) = \frac{1}{x} - \lfloor \frac{1}{x} \rfloor$ be the Gauss map from $(0, 1) \setminus \mathbb{Q}$ to itself. We have $k_{n+1} = \lfloor 1/G^n(w_0) \rfloor$ for all $n \geq 0$. For all $n \geq 0$, set

$$X_n = \{w_0 \in (0, 1) \setminus \mathbb{Q} \mid G^n(w_0) < \mathbf{D}^{-1}(n+1)^{-\gamma}\} = G^{-n}((0, \mathbf{D}^{-1}(n+1)^{-\gamma}) \setminus \mathbb{Q})$$

where G^{-n} is the n -th inverse image of sets. Let μ_G be the probability measure on $(0, 1) \setminus \mathbb{Q}$ with density $(\log 2)^{-1} (1+x)^{-1}$ (with respect to the Lebesgue measure). It is well-known that $\mu_G(X) = \mu_G(G^{-1}(X))$ for all measurable $X \subset (0, 1) \setminus \mathbb{Q}$. Therefore,

$$\mu_G(X_n) = \mu_G((0, \mathbf{D}^{-1}(n+1)^{-\gamma}) \setminus \mathbb{Q}) = \frac{1}{\log 2} \log \left(1 + \frac{1}{\mathbf{D}(n+1)^\gamma}\right) \leq \frac{1}{\log 2} \frac{1}{\mathbf{D}(n+1)^\gamma}$$

Let X_n^c be the complement of X_n in $(0, 1) \setminus \mathbb{Q}$. Then $\bigcap_{n \geq 0} X_n^c \subset F_w$, since $w_0 \in X_n^c$ implies $k_{n+1} = \lfloor 1/G^n(w_0) \rfloor \leq 1/G^n(w_0) \leq \mathbf{D}(n+1)^\gamma$. We have

$$\begin{aligned} \mu_G(F_w) &\geq \mu_G(\bigcap_{n \geq 0} X_n^c) = 1 - \mu_G(\bigcup_{n \geq 0} X_n) \geq 1 - \sum_{n \geq 0} \mu_G(X_n) \\ &\geq 1 - \frac{1}{\mathbf{D} \log 2} \sum_{n \geq 0} \frac{1}{(n+1)^\gamma} \geq 1 - \frac{1}{\mathbf{D} \log 2} \left(1 + \int_1^\infty x^{-\gamma} dx\right) = 1 - \frac{1}{\mathbf{D} \log 2} \frac{\gamma}{\gamma-1} > 0 \end{aligned}$$

Consequently, also the Lebesgue measure of F_w is positive. $\square$

7. Causal structure and particle horizons

In this section we show that the spatially homogeneous vacuum spacetimes corresponding to those solutions of (1.1) that are obtained by combining Theorems 6.2 and 6.3 and Propositions 3.1 and 3.3, have “particle horizons” (see [Mis] for this notion), contradicting a conjecture in [Mis].

Theorem 7.1. *Let $\mathbf{D}, \gamma, \mathbf{f}_0 = (\mathbf{h}_0, w_0, q_0)$ be as in Theorem 6.3. Let $\mathbf{f}_j = (\mathbf{h}_j, w_j, q_j)$, π_j, σ_* and $\mathbf{g}_j$ be as in Theorem 6.2. Adopt the remaining notation of Theorems 6.2 and*

6.3. Denote the components of $\mathbf{g}_j \in \mathcal{F}$ by $(\mathbf{h}'_j, w'_j, q'_j)$. Recall that $\mathbf{h}'_j \in (0, 1)$. Fix a constant $\lambda_0 > 0$. Set $\tau_0 = 0$. Introduce sequences $(\lambda_j)_{j \geq 0}$ and $(\tau_j)_{j \geq 0}$ by

$$\begin{aligned} \lambda_j &= \lambda_{j-1} \{A[\pi_j, \sigma_*](\mathbf{g}_j)\}^{-1} \in (0, \lambda_0] && \text{for all } j \geq 1 \\ \tau_j &= \tau_{j-1} + (\mathbf{h}'_j \lambda_j)^{-1} \{\tau_{1+}(\mathbf{g}_j) - \tau_{2-}[\pi_j, \sigma_*](\mathbf{g}_j)\} && \text{for all } j \geq 1 \end{aligned}$$

Then:

- (a) $\tau_j > \tau_{j-1}$ for all $j \geq 1$ and $\lim_{j \rightarrow \infty} \tau_j = +\infty$.
- (b) The solution to (1.1) with initial data $\Phi(0) = \lambda_0 \Phi_*(\pi_0, \mathbf{g}_0, \sigma_*)$ exists for all $\tau \geq 0$, that is $\Phi = \alpha \oplus \beta : [0, \infty) \rightarrow \mathcal{D}(\sigma_*)$, and $\Phi(\tau_j) = \lambda_j \Phi_*(\pi_j, \mathbf{g}_j, \sigma_*)$ for all $j \geq 0$.
- (c) For all $j \geq 1$ we have the bound

$$\mathbf{M}_j \stackrel{\text{def}}{=} \sup_{\tau \in (\tau_{j-1}, \tau_j)} \max_{(\mathbf{i}, \mathbf{j}, \mathbf{k}) \in \mathcal{C}} \alpha_{\mathbf{j}, \mathbf{k}}[\Phi](\tau) \leq -2^{-2} \lambda_j \min\{(w'_j)^2, (w'_j)^{-1}\}$$

Set $\zeta_{\mathbf{i}}(\tau) = -\frac{1}{2} \int_0^\tau ds \alpha_{\mathbf{i}}(s)$ for $\mathbf{i} = 1, 2, 3$ (see Proposition 2.1) and for all $s \geq 0$ set

$$\mathbf{L}(s) \stackrel{\text{def}}{=} \int_s^\infty d\tau \max_{(\mathbf{i}, \mathbf{j}, \mathbf{k}) \in \mathcal{C}} \exp(-\zeta_{\mathbf{j}} - \zeta_{\mathbf{k}})$$

(see the right hand side of (2.1) in Proposition 2.2). Then $\mathbf{L}(s) \leq \mathbf{L}(0) < \infty$ and $\lim_{s \rightarrow \infty} \mathbf{L}(s) = 0$.

Proof. Proposition 3.3 implies $\frac{1}{2} \leq \mathbf{h}'_j \lambda_j (\tau_j - \tau_{j-1}) \leq 2^2$ and $\tau_j \geq \tau_{j-1} + (2\lambda_0)^{-1}$, which implies (a). Theorems 6.2, 6.3, and Propositions 3.1, 3.3 imply (b). Proposition 3.3 (e.5) implies (c). Estimate

$$\begin{aligned} \mathbf{L}(0) &\leq \sum_{\ell=1}^\infty \int_{\tau_{\ell-1}}^{\tau_\ell} d\tau \exp\left(\frac{1}{2} \int_0^\tau d\tau' \max_{(\mathbf{i}, \mathbf{j}, \mathbf{k}) \in \mathcal{C}} \alpha_{\mathbf{j}, \mathbf{k}}[\Phi](\tau')\right) \\ &\leq \sum_{\ell=1}^\infty \int_{\tau_{\ell-1}}^{\tau_\ell} d\tau \exp\left(\frac{1}{2} \sum_{m=1}^{\ell-1} \int_{\tau_{m-1}}^{\tau_m} d\tau' \max_{(\mathbf{i}, \mathbf{j}, \mathbf{k}) \in \mathcal{C}} \alpha_{\mathbf{j}, \mathbf{k}}[\Phi](\tau')\right) \\ &\leq \sum_{\ell=1}^\infty (\tau_\ell - \tau_{\ell-1}) \exp\left(\frac{1}{2} \sum_{m=1}^{\ell-1} (\tau_m - \tau_{m-1}) \mathbf{M}_m\right) \\ &\leq 2^2 \sum_{\ell=1}^\infty (\mathbf{h}'_\ell \lambda_\ell)^{-1} \exp\left(-2^{-4} \sum_{m=1}^{\ell-1} (\mathbf{h}'_m)^{-1} \min\{(w'_m)^2, (w'_m)^{-1}\}\right) \end{aligned}$$

By Theorem 6.2, we have $(\delta_j, \mathbf{f}_j) \in B_2 \mathcal{F}$ and $\mathbf{g}_j \in B[\delta_j, \mathbf{f}_j]$ for all $j \geq 0$. Hence, $\frac{1}{2} \mathbf{h}_j \leq \mathbf{h}'_j \leq 2\mathbf{h}_j$ and $\frac{1}{2} w_j \leq w'_j \leq 2w_j$. Proposition 3.3 (b) implies $A[\pi_j, \sigma_*](\mathbf{g}_j) \leq 1 + \lambda_L(\mathbf{g}_j) \leq 3 + w'_j \leq 3(1 + w_j)$ and $\lambda_\ell^{-1} \leq \lambda_0^{-1} \prod_{k=1}^\ell 3(1 + w_k)$. Therefore,

$$\begin{aligned} \mathbf{L}(0) &\leq 2^3 (\lambda_0)^{-1} \sum_{\ell=1}^\infty (\mathbf{h}_\ell)^{-1} \left(\prod_{k=1}^\ell 3(1 + w_k) \right) \\ &\quad \times \exp\left(-2^{-7} \sum_{k=1}^{\ell-1} (\mathbf{h}_k)^{-1} \min\{(w_k)^2, (w_k)^{-1}\}\right) \\ &\leq 2^5 (\lambda_0)^{-1} \sum_{m=0}^\infty \sum_{\ell=J(m)+1}^{J(m+1)} (\mathbf{H}_\ell)^{-1} (2^7 \max_{1 \leq k \leq \ell} K_k)^\ell \\ &\quad \times \exp\left(-2^{-17} (1 - \delta_{\ell 1}) (\mathbf{H}_{\ell-1})^{-1} (K_{\ell-1})^{-2}\right) \quad (7.1) \end{aligned}$$

where $\mathbf{H}_j$ and K_j are as in Theorem 6.2, and $\delta_{\ell 1}$ is a Kronecker delta. See (6.2). In the exponential, we have bounded the sum over $k = 1, \dots, \ell-1$ from below by its $k = \ell-1$ summand if $\ell \geq 2$ and by zero otherwise. The sum over $\ell = J(m) + 1, \dots, J(m+1)$ has $k_{m+1} \leq \mathbf{D}(m+1)^\gamma$ many terms. By the proof of Theorem 6.3, for every $m \geq 0$, the following estimates, uniformly in $\ell = J(m) + 1, \dots, J(m+1)$, hold:

- $(\mathbf{H}_\ell)^{-1} \leq 2(\mathbf{h}_0)^{-1}(2\mathbf{D}(m+1)^\gamma)^{2(m+1)}$
- $(2^7 \max_{1 \leq k \leq \ell} K_k)^\ell \leq (2^{7+\gamma} \mathbf{D}(m+1)^\gamma)^{\mathbf{D}(m+1)^{\gamma+1}}$
- $(\mathbf{H}_{\ell-1})^{-1} \geq 2^{-4}(\mathbf{h}_0)^{-1} \rho_+^{2m}$ where $\rho_+ = \frac{1}{2}(1 + \sqrt{5})$
- $(K_{\ell-1})^{-2} \geq \mathbf{D}^{-2} 2^{-2\gamma} (m+1)^{-2\gamma}$

By these estimates, in particular the fact that $(\mathbf{H}_{\ell-1})^{-1}$ grows at least exponentially in m , the right hand side of (7.1) is finite, and $\mathbf{L}(0) < \infty$. $\square$

A. Bounds for a particular product of continued fractions

This appendix is entirely self-contained, the notation is completely local. Its single purpose is to prove Proposition A.1 below, which is used in the proof of Lemma 4.4.

Definition A.1. For all integers m and n and all sequences $(x_i)_{i \in \mathcal{I}}$ where $\mathcal{I} \subset \mathbb{Z}$, define $x_{m:n}$ to be the ordered sequence $x_m, x_{m+1}, \dots, x_{n-1}, x_n$ if $m \leq n$ and the empty sequence if $m > n$. In the first case, it is required that $[m, n] \cap \mathbb{Z} \subset \mathcal{I}$. Similarly, define $x_{m::n}$ to be the ordered sequence $x_m, x_{m-1}, \dots, x_{n+1}, x_n$ if $m \geq n$ and the empty sequence if $m < n$. In the first case, it is required that $[n, m] \cap \mathbb{Z} \subset \mathcal{I}$.

Definition A.2 (Continued fractions). For every integer $n \geq 0$ and every finite sequence of strictly positive integers $(k_i)_{1 \leq i \leq n}$ set recursively

$$\langle k_{1:n} \rangle = \begin{cases} 0 & n = 0 \\ (k_1 + \langle k_{2:n} \rangle)^{-1} & n \geq 1 \end{cases} \in [0, 1] \cap \mathbb{Q}$$

For every infinite sequence $(k_i)_{i \geq 1}$ of strictly positive integers, set

$$\langle k_1, k_2, \dots \rangle = \lim_{n \rightarrow \infty} \langle k_{1:n} \rangle \in (0, 1) \setminus \mathbb{Q}$$

Example A.1. $\langle \rangle = \langle k_{1:0} \rangle = 0$ and $\langle k_1 \rangle = \langle k_{1:1} \rangle = 1/k_1$ and $\langle k_1, k_2 \rangle = \langle k_{1:2} \rangle = 1/(k_1 + 1/k_2)$.

Definition A.3 (Fibonacci numbers). $F_1 = F_2 = 1$ and $F_n = F_{n-1} + F_{n-2}$, $n \geq 3$.

Proposition A.1. For every two-sided sequence of strictly positive integers $(k_i)_{i \in \mathbb{Z}}$, define two-sided sequences $(v_i)_{i \in \mathbb{Z}}$ and $(w_i)_{i \in \mathbb{Z}}$ by $v_i = \langle k_i, k_{i-1}, k_{i-2}, \dots \rangle$ and $w_i = \langle k_i, k_{i+1}, k_{i+2}, \dots \rangle$. Then, for all integers $M < N$:

- (a) $\frac{1}{2} \leq \prod_{i=M+1}^N (v_i/w_i) \leq 2$
- (b) $\prod_{i=M+1}^N w_i \leq (F_{N-M+1})^{-1} \leq (\frac{1}{2}(\sqrt{5} - 1))^{N-M-1}$

The proof of Proposition A.1 is given at the end of this appendix.

Definition A.4. Let $P_0(\cdot) = 1$ and $P_1(x_1) = x_1$ and for all $n \geq 2$, set

$$P_n(x_{1:n}) = x_1 P_{n-1}(x_{2:n}) + P_{n-2}(x_{3:n}) \quad (\text{A.1})$$

Example A.2. $P_2(x_{1:2}) = 1 + x_1 x_2$ and $P_3(x_{1:3}) = x_1 + x_3 + x_1 x_2 x_3$ and $P_4(x_{1:4}) = 1 + x_1 x_2 + x_3 x_4 + x_1 x_4 + x_1 x_2 x_3 x_4$.

Lemma A.1. Recall Definition A.4. For all integers $n \geq 0$, we have:

- (a) P_n is a polynomial of degree n , jointly in its n arguments, with coefficients in $\{0, 1\}$
 (b) P_n is a polynomial of degree 1, separately in each of its n arguments
 (c) $P_n(1, \dots, 1) = F_{n+1}$
 (d) $P_n(x_{1:n}) = P_n(x_{n::1})$ for all $x_1, \dots, x_n \in \mathbb{R}$
 (e) $\langle k_{1:n} \rangle = P_{n-1}(k_{2:n})/P_n(k_{1:n})$ for all strictly positive integers $(k_i)_{1 \leq i \leq n}$, $n \geq 1$

Proof. (a) through (e) are all shown by induction, using (A.1). To show (d), observe that (d)₀, (d)₁, (d)₂ and (d)₃ hold. For the induction step, let $n \geq 4$ and suppose (d)₀ through (d)_{n-1} hold. Then, using only (A.1) and the induction hypothesis,

$$\begin{aligned} & P_n(x_{1:n}) - P_n(x_{n::1}) \\ &= x_1 P_{n-1}(x_{2:n}) + P_{n-2}(x_{3:n}) - x_n P_{n-1}(x_{n-1::1}) - P_{n-2}(x_{n-2::1}) \\ &= x_1 P_{n-1}(x_{n::2}) + P_{n-2}(x_{n::3}) - x_n P_{n-1}(x_{1:n-1}) - P_{n-2}(x_{1:n-2}) \\ &= x_1 (x_n P_{n-2}(x_{n-1::2}) + P_{n-3}(x_{n-2::2})) + (x_n P_{n-3}(x_{n-1::3}) + P_{n-4}(x_{n-2::3})) \\ &\quad - x_n (x_1 P_{n-2}(x_{2:n-1}) + P_{n-3}(x_{3:n-1})) - (x_1 P_{n-3}(x_{2:n-2}) + P_{n-4}(x_{3:n-2})) \end{aligned}$$

Verify that all the terms cancel, by the induction hypothesis. This implies (d)_n. To show (e), observe that (e)₁ holds. Let $n \geq 2$ and suppose (e)_{n-1} holds. Then,

$$\langle k_{1:n} \rangle = (k_1 + \langle k_{2:n} \rangle)^{-1} = \left(k_1 + \frac{P_{n-2}(k_{3:n})}{P_{n-1}(k_{2:n})} \right)^{-1} = \frac{P_{n-1}(k_{2:n})}{k_1 P_{n-1}(k_{2:n}) + P_{n-2}(k_{3:n})}$$

Now, (A.1) implies (e)_n. $\square$

Lemma A.2. For all integers $m-1 \leq M < N \leq n$ and all $x_m, \dots, x_n \in [1, \infty)$,

$$2 P_{M-m+1}(x_{m:M}) P_{N-M}(x_{M+1:N}) - P_{N-m+1}(x_{m:N}) P_{n-N}(x_{N+1:n}) \geq 0 \quad (\text{A.2})$$

Moreover, if $m = M+1$, then the factor 2 on the left hand side can be dropped, that is,

$$P_{n-M}(x_{M+1:n}) - P_{N-M}(x_{M+1:N}) P_{n-N}(x_{N+1:n}) \geq 0 \quad (\text{A.3})$$

Proof. In this proof, we use the recursion relation (A.1) and the reflected recursion relation that is obtained by applying Lemma A.1 (d) to all three terms of (A.1). Fix M and N . Inequality (A.2) is proved by induction over m and n , where $m \leq M+1$ and $n \geq N$. Denote the left hand side of (A.2) by $Q_{m,n}$. Then,

$$\begin{aligned} Q_{M+1,N} &= P_{N-M}(x_{M+1:N}) \geq 0 \\ Q_{M+1,N+1} &= 2 P_{N+1-M}(x_{M+1:N+1}) - P_{N-M}(x_{M+1:N}) x_{N+1} \\ &= P_{N-M}(x_{M+1:N}) x_{N+1} + 2 P_{N-1-M}(x_{M+1:N-1}) \geq 0 \\ Q_{M,N} &= 2 x_M P_{N-M}(x_{M+1:N}) - P_{N-M+1}(x_{M:N}) \\ &= x_M P_{N-M}(x_{M+1:N}) - P_{N-M-1}(x_{M+2:N}) \\ &\geq x_M x_{M+1} P_{N-M-1}(x_{M+2:N}) - P_{N-M-1}(x_{M+2:N}) \geq 0 \\ Q_{M,N+1} &= 2 x_M P_{N+1-M}(x_{M+1:N+1}) - P_{N-M+1}(x_{M:N}) x_{N+1} \\ &= 2 x_M x_{N+1} P_{N-M}(x_{M+1:N}) + 2 x_M P_{N-1-M}(x_{M+1:N-1}) \\ &\quad - x_M P_{N-M}(x_{M+1:N}) x_{N+1} - P_{N-M-1}(x_{M+2:N}) x_{N+1} \\ &\geq x_M x_{N+1} P_{N-M}(x_{M+1:N}) - P_{N-M-1}(x_{M+2:N}) x_{N+1} \\ &\geq x_M x_{N+1} x_{M+1} P_{N-M-1}(x_{M+2:N}) - P_{N-M-1}(x_{M+2:N}) x_{N+1} \geq 0 \end{aligned}$$

These four cases and the two recursion relations

- $Q_{m,n} = x_m Q_{m+1,n} + Q_{m+2,n}$ when $m \leq M-1$ and $n \geq N$
- $Q_{m,n} = x_n Q_{m,n-1} + Q_{m,n-2}$ when $m \leq M+1$ and $n \geq N+2$

imply (A.2). Inequality (A.3) is shown in an entirely similar way. $\square$

Proof (of Proposition A.1). Recall (d), (e) in Lemma A.1. Let $m-1 \leq M < N \leq n$.

$$\begin{aligned} \prod_{i=M+1}^N \frac{\langle k_{i::m} \rangle}{\langle k_{i:n} \rangle} &= \prod_{i=M+1}^N \frac{P_{i-m}(k_{i-1::m})}{P_{i-m+1}(k_{i::m})} \cdot \frac{P_{n-i+1}(k_{i:n})}{P_{n-i}(k_{i+1:n})} \\ &= \frac{P_{M-m+1}(k_{M::m})}{P_{N-m+1}(k_{N::m})} \cdot \frac{P_{n-M}(k_{M+1:n})}{P_{n-N}(k_{N+1:n})} = \frac{P_{M-m+1}(k_{m:M})}{P_{N-m+1}(k_{m:N})} \cdot \frac{P_{n-M}(k_{M+1:n})}{P_{n-N}(k_{N+1:n})} \end{aligned}$$

The right hand side is $\geq \frac{1}{2}$, by inequality (A.2). Now, let $m \rightarrow -\infty$ and $n \rightarrow +\infty$ to obtain $\prod_{i=M+1}^N (v_i/w_i) \geq \frac{1}{2}$. By symmetry, we also have $\prod_{i=M+1}^N (w_i/v_i) \geq \frac{1}{2}$. This implies (a) in Proposition A.1. Similarly, using (A.3),

$$\prod_{i=M+1}^N \langle k_{i:n} \rangle = \frac{P_{n-N}(k_{N+1:n})}{P_{n-M}(k_{M+1:n})} \leq \frac{1}{P_{n-M}(k_{M+1:n})} \leq \frac{1}{P_{n-M}(1, \dots, 1)}$$

Let $n \rightarrow +\infty$ to obtain $\prod_{i=M+1}^N w_i \leq 1/F_{N-M+1}$. $\square$

B. The modulus of continuity of the map $\mathcal{Q}_L$ introduced in Definition 3.16

Lemma B.1. Let $\mathcal{Q}_L : (0, \infty)^3 \rightarrow (0, \infty)^2 \times \mathbb{R}$ be the map introduced in Definition 3.16. For all $\mathbf{f}_i = (\mathbf{h}_i, w_i, q_i) \in (0, \infty)^3$ with $0 < \mathbf{h}_i \leq 1$, $i = 1, 2$, with $\mathbf{f}_1 \neq \mathbf{f}_2$, such that q_1 and q_2 are either both < 1 or both > 1 ,

$$\frac{\|\mathcal{Q}_L(\mathbf{f}_2) - \mathcal{Q}_L(\mathbf{f}_1)\|_{\mathbb{R}^3}}{\|\mathbf{f}_2 - \mathbf{f}_1\|_{\mathbb{R}^3}} \leq \begin{cases} 2^{12} q_{\min}^{-2} \log(2 + w_{\max}) & \text{if } q_1, q_2 < 1 \\ 2^{11} q_{\max} & \text{if } q_1, q_2 > 1 \end{cases}$$

Here, $w_{\max} = \max\{w_1, w_2\}$ and $q_{\max} = \max\{q_1, q_2\}$ and $q_{\min} = \min\{q_1, q_2\}$.

Proof. We prove the following claim, which implies the Lemma: *Each of the nine partial derivatives of $\mathcal{Q}_L : \mathbf{f} = (\mathbf{h}, w, q) \mapsto \mathcal{Q}_L(\mathbf{f})$ is bounded in absolute value by*

$$\begin{cases} 2^{10} q^{-2} \log(2 + w) & \text{if } \mathbf{f} \in (0, 1] \times (0, \infty) \times (0, 1) \\ 2^9 q & \text{if } \mathbf{f} \in (0, 1] \times (0, \infty) \times (1, \infty) \end{cases} > 1 \quad (\text{B.1})$$

Let $0 < \mathbf{h} \leq 1$ and $q \neq 1$. Let $(\mathbf{h}_L, w_L, q_L) = \mathcal{Q}_L(\mathbf{f})$ and let $\text{num}1_L, \text{num}2_L, \text{den}_L$ be as in Definition 3.16. We first estimate the partial derivatives of $q_L = \text{num}1_L/\text{den}_L$ and $\mathbf{h}_L = \text{num}2_L/\text{den}_L$. Each of $\text{num}1_L, \text{num}2_L, \text{den}_L$ is of the form

$$L_1(w, q) + L_2(w, q)q + L_3(w, q)\mathbf{h} + L_4(w, q)\mathbf{h} \log \lambda_L(\mathbf{f})$$

with $\lambda_L(\mathbf{f}) = 1 + 1/w_L(\mathbf{f})$ as in Definition 3.16 and with $L_i(w, q) = a_i(q)w + b_i(q)$ where $a_i(q)$ and $b_i(q)$ are constant separately for $q < 1$ and for $q > 1$ and satisfy $-3 \leq a_i(q), b_i(q) \leq 3$, where $i = 1, 2, 3, 4$. Let $k = 1, 2$ and $\text{num}k_L = L_1 + L_2q +$

$L_3 \mathbf{h} + L_4 \mathbf{h} \log \lambda_L$ and $\text{den}_L = L'_1 + L'_2 q + L'_3 \mathbf{h} + L'_4 \mathbf{h} \log \lambda_L$ with $L_i = a_i w + b_i$ and $L'_i = a'_i w + b'_i$ (Warning: the prime does *not* denote a derivative). Then

$$\begin{aligned} & \left(\frac{\partial}{\partial x} \text{num} k_L \right) \text{den}_L - \left(\frac{\partial}{\partial x} \text{den}_L \right) \text{num} k_L \\ = & \begin{cases} (L_3 + L_4 \log \lambda_L)(L'_1 + L'_2 q) - (L'_3 + L'_4 \log \lambda_L)(L_1 + L_2 q) & \text{if } x = \mathbf{h} \\ \begin{aligned} & + (a_1 + a_2 q + a_3 \mathbf{h} + a_4 \mathbf{h} \log \lambda_L)(b'_1 + b'_2 q + b'_3 \mathbf{h} + b'_4 \mathbf{h} \log \lambda_L) \\ & - (a'_1 + a'_2 q + a'_3 \mathbf{h} + a'_4 \mathbf{h} \log \lambda_L)(b_1 + b_2 q + b_3 \mathbf{h} + b_4 \mathbf{h} \log \lambda_L) \\ & + \mathbf{h} \left\{ L_4 (L'_1 + L'_2 q + L'_3 \mathbf{h}) - L'_4 (L_1 + L_2 q + L_3 \mathbf{h}) \right\} \frac{\partial}{\partial w} \log \lambda_L \end{aligned} & \text{if } x = w \\ L_2 (L'_1 + L'_3 \mathbf{h} + L'_4 \mathbf{h} \log \lambda_L) - L'_2 (L_1 + L_3 \mathbf{h} + L_4 \mathbf{h} \log \lambda_L) & \text{if } x = q \end{cases} \end{aligned}$$

Recall that $|\mathbf{h}| \leq 1$ and $|a_i|, |a'_i|, |b_i|, |b'_i| \leq 3$ and $|L_i|, |L'_i| \leq 3(1+w)$ and $\log \lambda_L \geq 0$.

- If $q < 1$, then $|\frac{\partial}{\partial w} \log \lambda_L| \leq (1+w)^{-1}$ and

$$\begin{aligned} & \left| \left(\frac{\partial}{\partial x} \text{num} k_L \right) \text{den}_L - \left(\frac{\partial}{\partial x} \text{den}_L \right) \text{num} k_L \right| \\ & \leq \begin{cases} 36(1+w)^2(1+\log \lambda_L) & \text{if } x = \mathbf{h} \\ 18(3+\log \lambda_L)^2 + 54(1+w) & \text{if } x = w \\ 18(1+w)^2(2+\log \lambda_L) & \text{if } x = q \end{cases} \leq 2^{10}(1+w)^2 \log(2+w) \end{aligned}$$

For the second inequality, use $\frac{1}{2} \leq \log \lambda_L \leq 1+w$ and $(\log \lambda_L)^2 \leq 1+w$.

- If $q > 1$, then $|\log \lambda_L| \leq 1$ and $|\frac{\partial}{\partial w} \log \lambda_L| \leq (1+w)^{-2}$ and $a'_2 = b'_2 = 0$ and

$$\begin{aligned} & \left| \left(\frac{\partial}{\partial x} \text{num} k_L \right) \text{den}_L - \left(\frac{\partial}{\partial x} \text{den}_L \right) \text{num} k_L \right| \\ & \leq \begin{cases} 72(1+w)^2 q & \text{if } x = \mathbf{h} \\ 270 q & \text{if } x = w \\ 27(1+w)^2 & \text{if } x = q \end{cases} \leq 2^9 (1+w)^2 q \end{aligned}$$

To finish the proof, observe that $\text{den}_L \geq (1+w) \min\{1, q\} > 0$. Each partial derivative of $q_L = \text{num} 1_L / \text{den}_L$ and $\mathbf{h}_L = \text{num} 2_L / \text{den}_L$ is bounded in absolute value by (B.1). And so are the partial derivatives of w_L , because $\partial w_L / \partial \mathbf{h} = \partial w_L / \partial q = 0$, and because $\partial w_L / \partial w = -(1+w)^{-2}$ if $q < 1$ and $\partial w_L / \partial w = 1$ if $q > 1$. $\square$

References

- Be. Béguin, F., arXiv 1004.2984 (2010)
 BKL1. Belinskii, V.A., Khalatnikov, I.M., and Lifshitz, E.M., Adv. Phys. **19**, (1970) 525-573
 HU1. Heinzle, J.M. and Uggla, C., arXiv 0901.0776 (2009)
 HU2. Heinzle, J.M. and Uggla, C., arXiv 0901.0806 (2009)
 LHWG. Liebscher, S., Härterich, J., Webster, K. and Georgi, M., arXiv 1004.1989 (2010)
 Mis. Misner, C.W., Phys. Rev. Lett. **22**, (1969) 1071-1074
 Ren. Rendall, A., Class. Quantum Grav. **14**, (1997) 2341-2356
 Ri1. Ringström, H., Class. Quantum Grav. **17** (2000) 713
 Ri2. Ringström, H., Annales Henri Poincaré **2** (2001) 405